



# Hierarchical spatial modelling for applied population and community ecology

---

Jeffrey W. Doser, Marc Kéry,  
Gesa von Hirschheydt

24-27 June 2024







# Multi-season spatial occupancy models

---

Jeffrey W. Doser  
24-27 June 2024



Why use multi-season occupancy models?

# Why use multi-season occupancy models?

- Understand how species distributions are shifting over space and time

# Why use multi-season occupancy models?

- Understand how species distributions are shifting over space and time
- Assess occupancy trends over time
  - Detection-nondetection data are easier to collect than count data
  - Occupancy-abundance relationship
  - Exact interpretation of occupancy trends depends on how data are collected (see Steenweg et al. 2018 Ecology)

# Why use multi-season occupancy models?

- Understand how species distributions are shifting over space and time
- Assess occupancy trends over time
  - Detection-nondetection data are easier to collect than count data
  - Occupancy-abundance relationship
  - Exact interpretation of occupancy trends depends on how data are collected (see Steenweg et al. 2018 Ecology)
- From a statistical perspective, having multiple seasons of data can improve our ability to estimate spatial random effects.

# Multi-season detection-nondetection data

- Data follow the **robust design**

# Multi-season detection-nondetection data

- Data follow the **robust design**
- A set of  $J$  sites are sampled across a set of  $T$  seasons (e.g., years)



# Multi-season detection-nondetection data

- Data follow the **robust design**
- A set of  $J$  sites are sampled across a set of  $T$  seasons (e.g., years)
- Within each season  $t$ , each site  $j$  is sampled  $K_{jt}$  times.

# Multi-season detection-nondetection data

- Data follow the **robust design**
- A set of  $J$  sites are sampled across a set of  $T$  seasons (e.g., years)
- Within each season  $t$ , each site  $j$  is sampled  $K_{jt}$  times.
- Multi-season occupancy models can easily handle "imbalanced" data sets
  - Each site may not be sampled each season
  - Each sampled site may not be sampled the same number of times within a season

# Multi-season detection-nondetection data

- Data follow the **robust design**
- A set of  $J$  sites are sampled across a set of  $T$  seasons (e.g., years)
- Within each season  $t$ , each site  $j$  is sampled  $K_{jt}$  times.
- Multi-season occupancy models can easily handle "imbalanced" data sets
  - Each site may not be sampled each season
  - Each sampled site may not be sampled the same number of times within a season
- Seasons are sometimes referred to as **primary replicates** and repeat visits within a season as **secondary replicates**

# Multi-season detection-nondetection data

- Recall the **closure assumption**.
- For multi-season models, we assume closure within a primary time period (i.e., season), but do not assume closure across primary time periods (i.e., across seasons)
- In other words, we assume the occupancy status does not change within a season, but that sites can change from occupied to unoccupied across seasons.



# Multi-species detection-nondetection data

- Example: 6 sites, 2 seasons, 3 surveys within a season

Season 1

Site	Survey 1	Survey 2	Survey 3
1	1	0	0
2	0	0	0
3	1	1	0
4	1	NA	0
5	0	1	1
6	0	0	0

Season 2

Site	Survey 1	Survey 2	Survey 3
1	0	1	NA
2	1	0	0
3	1	1	0
4	1	1	0
5	NA	NA	NA
6	0	0	1

# Lots of interesting design questions in multi-season occupancy models



**GfÖ**

GfÖ Ecological Society of Germany,  
Austria and Switzerland

Basic and Applied Ecology 67 (2023) 61–69

**Basic and  
Applied Ecology**

[www.elsevier.com/locate/baaec](http://www.elsevier.com/locate/baaec)

RESEARCH PAPER

## “Mixed” occupancy designs: When do additional single-visit data improve the inferences from standard multi-visit models?

Gesa von Hirschheydt<sup>a,\*</sup>, Silvia Stofer<sup>a</sup>, Marc Kéry<sup>b</sup>

<sup>a</sup>Swiss Federal Research Institute for Forest, Snow and Landscape Research WSL, 8903 Birmensdorf, Switzerland

<sup>b</sup>Swiss Ornithological Institute, 6204 Sempach, Switzerland

Received 27 May 2022; accepted 29 January 2023  
Available online 31 January 2023



**Methods in Ecology and Evolution**



RESEARCH ARTICLE | [Open Access](#) |

## “Fractional replication” in single-visit multi-season occupancy models: Impacts of spatiotemporal autocorrelation on identifiability

Jeffrey W. Doser Sara Stoudt

First published: 26 December 2023 | <https://doi.org/10.1111/2041-210X.14275>

Jeffrey W. Doser and Sara Stoudt contributed equally to this work.

**Handling Editor:** Xingfeng Si

# Terminology

## **Multi-season occupancy model**

Dynamic  
occupancy  
model

Stacked  
occupancy  
model

Spatio-temporal  
occupancy  
model

# Dynamic occupancy models

- MacKenzie et al. (2003)
- Explicitly estimates colonization and extinction (or survival)
- Arguably the most mechanistic form of multi-season occupancy model
- More data hungry and often harder to fit
- Can fit in `unmarked` (frequentist) and `ubms` (Bayesian)



# Stacked occupancy models

- Fit a single-season occupancy model, but now your "sites" are really "site-seasons" (e.g., "site-years", combinations of site and year).
- Less mechanistic, but far less data hungry than dynamic models.
- Often a season trend is of interest and included in the model.
- Can account for pseudoreplication by including a random site effect.
- Can fit stacked models in `spOccupancy`

# Spatio-temporal occupancy models

- A form of stacked occupancy model, but now includes explicit components to account for spatial and/or temporal autocorrelation.
- Less mechanistic than dynamic models.
- Better at predicting distributions (and changes) than basic stacked models
- Lots of different flavors.
- Some nice examples:
  - Rushing et al. (2019) Scientific Reports
  - Wright et al. (2021) Ecology and Evolution
  - Hepler et al. (2023) R Journal (the `multiocc` package)

# Terminology

## **Multi-season occupancy model**

Dynamic  
occupancy  
model

Stacked  
occupancy  
model

Spatio-temporal  
occupancy  
model

# Multi-season occupancy model

$j = 1, \dots, J$  (site)

$t = 1, \dots, T$  (season)

## Occupancy (ecological) sub-model

$k = 1, \dots, K_{j,t}$  (replicate)

$$z_{j,t} \sim \text{Bernoulli}(\psi_{j,t})$$

$$\text{logit}(\psi_{j,t}) = \mathbf{x}_{j,t}\boldsymbol{\beta} + w_j + \eta_t$$

$z_{j,t}$  True occurrence of the species at site  $j$  in season  $t$

$\psi_{j,t}$  Occurrence probability at site  $j$  in season  $t$

$\mathbf{x}_{j,t}$  Site and/or season-varying covariates

$w_j$  Site-level random effect

$\eta_t$  Season-level (temporal) random effect



# Multi-season occupancy model

$j = 1, \dots, J$  (site)

$t = 1, \dots, T$  (season)

## Occupancy (ecological) sub-model

$k = 1, \dots, K_{j,t}$  (replicate)

$$z_{j,t} \sim \text{Bernoulli}(\psi_{j,t})$$

$$\text{logit}(\psi_{j,t}) = \mathbf{x}_{j,t}\boldsymbol{\beta} + \boxed{w_j + \eta_t}$$

$z_{j,t}$  True occurrence of the species at site  $j$  in season  $t$

$\psi_{j,t}$  Occurrence probability at site  $j$  in season  $t$

$\mathbf{x}_{j,t}$  Site and/or season-varying covariates

$w_j$  Site-level random effect

$\eta_t$  Season-level (temporal) random effect

# Site-level random effects $w_j$

Two types:

1. **Unstructured** -> a typical random intercept with the form:

$$w_j \sim \text{Normal}(0, \sigma^2)$$

2. **Spatial NNGP** -> same as before. This is a "spatial multi-season occupancy model" or "spatio-temporal occupancy model".

# Unstructured site-level random effects

- This is the standard approach in stacked occupancy models.
- Random site effect accounts for non-independence between occupancy probability at a site over the T seasons (i.e., pseudoreplication).
- Does not explicitly account for spatial autocorrelation
- Often reasonable when focus is on inference, but spatial effects are often much better for prediction.

$$w_j \sim \text{Normal}(0, \sigma^2)$$

# Spatial NNGP site-level random effects

- Account for spatial autocorrelation in occupancy probability.
- Nothing new here from previous spatial models.

$$\mathbf{w} \sim \text{Normal}(\mathbf{0}, \tilde{\mathbf{C}}(d, \phi, \sigma^2))$$



# Temporal random effects $\eta_t$

- Account for correlation in occupancy probability over time

# Temporal random effects $\eta_t$

- Account for correlation in occupancy probability over time
- Alternative view: account for non-linear variation in occupancy probability when estimating a trend

# Temporal random effects $\eta_t$

- Account for correlation in occupancy probability over time
- Alternative view: account for non-linear variation in occupancy probability when estimating a trend
- Two types:
  1. Unstructured -> a typical random intercept with the form:

$$\eta_t \sim \text{Normal}(0, \sigma_T^2)$$

# Temporal random effects $\eta_t$

- Account for correlation in occupancy probability over time
- Alternative view: account for non-linear variation in occupancy probability when estimating a trend
- Two types:
  1. Unstructured -> a typical random intercept with the form:

$$\eta_t \sim \text{Normal}(0, \sigma_T^2)$$

2. AR(1) -> random temporal effects follow an autoregressive structure. Covariance between two time points is:

$$\sigma_T^2 \rho^{|t-t'|}$$

# Multi-season occupancy model

$$z_{j,t} \sim \text{Bernoulli}(\psi_{j,t})$$

$$\text{logit}(\psi_{j,t}) = \mathbf{x}_{j,t}\boldsymbol{\beta} + \boxed{w_j + \eta_t}$$

- In the statistics literature, this is known as a **separable** model, because the spatial random effects are independent from the temporal random effects.

# Multi-season occupancy model

$$z_{j,t} \sim \text{Bernoulli}(\psi_{j,t})$$

$$\text{logit}(\psi_{j,t}) = \mathbf{x}_{j,t}\boldsymbol{\beta} + \boxed{w_j + \eta_t}$$

- In the statistics literature, this is known as a **separable** model, because the spatial random effects are independent from the temporal random effects.
- Nonseparable models allow different types of interactions between the spatial and temporal effects.
  - Examples include Wright et al. (2021) and Hepler et al. (2021).

# Multi-season occupancy model

$$z_{j,t} \sim \text{Bernoulli}(\psi_{j,t})$$

$$\text{logit}(\psi_{j,t}) = \mathbf{x}_{j,t}\boldsymbol{\beta} + \boxed{w_j + \eta_t}$$

- In the statistics literature, this is known as a **separable** model, because the spatial random effects are independent from the temporal random effects.
- Nonseparable models allow different types of interactions between the spatial and temporal effects.
  - Examples include Wright et al. (2021) and Hepler et al. (2021).
- We will shortly discuss applications of spatially-varying coefficient models in `spOccupancy` for looking at spatial variation in occupancy trends over time.

# Multi-season occupancy model

$j = 1, \dots, J$  (site)

$t = 1, \dots, T$  (season)

$k = 1, \dots, K_{j,t}$  (replicate)

## Detection (observation) sub-model

$$y_{j,t,k} \sim \text{Bernoulli}(p_{j,t,k} \cdot z_{j,t})$$

$$\text{logit}(p_{j,t,k}) = \mathbf{v}_{j,t,k} \cdot \boldsymbol{\alpha}$$

$y_{j,t,k}$  Detection-nondetection data at site  $j$  during replicate  $k$  and season  $t$

$p_{j,t,k}$  Detection probability at site  $j$  during replicate  $k$  and season  $t$

$\mathbf{v}_{j,t,k}$  Covariates affecting detection at site  $j$  during replicate  $k$  and season  $t$



# Fitting multi-season occupancy models in `spOccupancy`

- `tPGOcc()` : non-spatial multi-season occupancy models (temporal Pólya-Gamma occupancy model)
- `stPGOcc()` : spatio-temporal Pólya-Gamma occupancy models
- All multi-season models require the use of an adaptive Metropolis-Hastings sampler, and so we specify the number of batches and batch length as with previous spatial models.
- `tMsPGOcc()` and `stMsPGOcc()` for multi-species models.



# Fitting multi-season occupancy models in `spOccupancy`

Site Effect	Temporal Effect	<code>spOccupancy</code>
None	None	<code>tPGOcc()</code>
None	Unstructured	<code>tPGOcc()</code> with random time intercept
None	AR(1)	<code>tPGOcc()</code> with <code>ar1 = TRUE</code>
Unstructured	None	<code>tPGOcc()</code> with random site intercept
Unstructured	Unstructured	<code>tPGOcc()</code> with random time and site intercept
Unstructured	AR(1)	<code>tPGOcc()</code> with random site intercept and <code>ar1 = TRUE</code>
Spatial (NNGP)	None	<code>stPGOcc()</code>
Spatial (NNGP)	Unstructured	<code>stPGOcc()</code> with random time intercept
Spatial (NNGP)	AR(1)	<code>stPGOcc()</code> with <code>ar1 = TRUE</code>

Different ways to model the site-level and temporal random effects in multi-season occupancy models in `spOccupancy`.



# Exercise: Estimating bat distributions in the Western USA

---

7-bat-multi-season-occ.R

