

The genetic architecture of target-site resistance to DDT and pyrethroids in the malaria vectors *Anopheles gambiae* and *Anopheles coluzzii*

DRAFT

Chris S. Clarkson¹, Alistair Miles^{2,1}, Nicholas J. Harding², @@TODO¹,
Dominic Kwiatkowski^{1,2}, Martin Donnelly^{3,1}, and The *Anopheles gambiae*
1000 Genomes Consortium⁴

¹Sanger @@TODO

²Oxford @@TODO

³Liverpool @@TODO

⁴MalariaGEN @@TODO

18th February 2017

Abstract

TODO

Introduction

The malaria vectors *Anopheles gambiae* and *Anopheles coluzzii* are evolving insecticide resistance. Asdlkj dsalkj daslkjd aslkjdas lkadsj lkadsj adslkj adslkja dslkadsj lkadsj lkasd jlkadsj lkads jlkads alksdj asdlk jasdlk adslk jadslkj adslkj adslkj adslkj adslkj adslkj adslk jasd.

This is the second paragraph of the introduction. Asdlkjio weipo ewrpoi ewrpoi rwepoi werpoiwe poi rewpoi rwepoi erwpoi rewpoi rwepoi rwepoi rwepoi rwepoi rewpo iwrepoi rwepoi wrepoi wrepoi rwepo irwpo irwepoi rwepo ipoewi rpow ierwe.

Third paragraph zcx,m ncxz,mxczn ,mxczn,mxcz n,mcnxz ,mxczn ,mz ncx,mzcxn.

Fourth paragraph asdlkj dsakljdsalkj dsalkj daslkj daslkj dsalkj daslkj sda [1].

TODO

Results

Let's add some results qweoi qewoiewqoip peqwpoi ewqpoi ewqpoieqwipo ewqipo eqwpio eqwipo ewqoip ewqoip ewqioip eqw.

Isn't Figure 1 interesting! Table 2 is pretty interesting too.

Mutation				Population allele frequency (%)									LD (D')	
Position ¹		Ag^2	Md^3	AOAc	BFAc	GNAg	BFAg	CMAg	GAAG	UGAg	KE	GW	L995F	L995S
2,390,177	G>A	R254K	NA	0	0	0	0	32	21	0	0	0	NA	NA
2,391,228	G>C	V402L	NA	0	7	0	0	0	0	0	0	0	NA	NA
2,391,228	G>T	V402L	NA	0	7	0	0	0	0	0	0	0	NA	NA
2,399,997	G>C	D466H	NA	0	0	0	0	7	0	0	0	0	NA	NA
2,400,071	G>A	M490I	NA	0	0	0	0	0	0	0	18	0	NA	NA
2,400,071	G>T	M490I	NA	0	0	0	0	0	0	0	0	0	NA	NA
2,416,980	C>T	T791M	NA	0	1	13	14	0	0	0	0	0	NA	NA
2,422,651	T>C	L995S	NA	0	0	0	0	15	64	100	76	0	NA	NA
2,422,652	A>T	L995F	NA	86	85	100	100	53	36	0	0	0	NA	NA
2,424,384	C>T	A1125V	NA	9	0	0	0	0	0	0	0	0	NA	NA
2,425,077	G>A	V1254I	NA	0	0	0	0	0	0	0	0	5	NA	NA
2,429,617	T>C	I1527T	NA	0	14	0	0	0	0	0	0	0	NA	NA
2,429,745	A>T*	N1570Y	NA	0	26	10	22	6	0	0	0	0	NA	NA
2,429,897	A>G	E1597G	NA	0	0	6	4	0	0	0	0	0	NA	NA
2,429,915	A>C	K1603T	NA	0	5	0	0	0	0	0	0	0	NA	NA
2,430,424	G>T	A1746S	NA	0	0	11	13	0	0	0	0	0	NA	NA
2,430,817	G>A	V1853I	NA	0	0	8	5	0	0	0	0	0	NA	NA
2,430,863	T>C	I1868T	NA	0	0	18	25	0	0	0	0	0	NA	NA
2,430,880	C>T	P1874S	NA	0	21	0	0	0	0	0	0	0	NA	NA
2,430,881	C>T	P1874L	NA	0	7	45	26	0	0	0	0	0	NA	NA
2,431,061	C>T	A1934V	NA	0	12	0	0	0	0	0	0	0	NA	NA
2,431,079	T>C	I1940T	NA	0	4	0	0	7	0	0	0	0	NA	NA

Table 1. Non-synonymous mutations in the voltage-gated sodium channel gene. All mutations are at 5% frequency or above in one or more of the 9 Ag1000G phase 1 populations, with the exception of 2,400,071 G>T which is only found in the CMAg population at 0.4% frequency but is included because another mutation (2,400,071 G>A) is found at the same position causing the same amino acid substitution (M490I). Substitutions marked with an asterisk (*) failed conservative variant filters applied genome-wide in the Ag1000G phase 1 AR3 callset, but appeared sound on manual inspection of read alignments.

¹Position relative to AgamP3 reference sequence, chromosome arm 2L.

²Codon numbering according to transcript AGAP004707-RA in geneset AgamP4.4.

³Codon numbering according to @@TODO.




Figure 1. Demo figure.

TODO

Foo	Bar	Baz
1	a	True
2	b	False

Table 2. This is a table.

Discussion

TODO

Methods

TODO

References

- [1] Nandita R. Garud et al. ‘Recent Selective Sweeps in North American *Drosophila melanogaster* Show Signatures of Soft Sweeps’. In: *PLoS Genetics* 11.2 (2015), pp. 1–32. ISSN: 15537404. arXiv: 1303.0906.