

Basic math for machine learning



Departament de Teoria
del Senyal i Comunicacions



CONTENTS

1. Random variables
2. Matrix algebra: operators, norms and eigenvectors
3. Optimization with restrictions
4. Derivation of real variable functions

Definition of a random variable

Definition: A random variable X is an application from the result of a random experiment to a real value:

$$X : \Omega \rightarrow \mathbb{R}$$

Characterization of a random variable

The density function $f(x)$ characterizes X :


Property 1. It is non-negative.

Property 2. The area under an interval is the probability that X takes values within that interval.

$$\Pr(a \leq x \leq b) = \int_a^b f_x(x) dx$$

Property 3. The joint density of several independent variables is the product of density functions.

$$f_{xy}(x, y) = f_x(x) f_y(y)$$



Example 1. $f(x)$ for a real-valued Gaussian random variable

$$f_x(x) = \frac{1}{\sqrt{2\pi\sigma_x^2}} \exp\left(-\frac{|x - m_x|^2}{2\sigma_x^2}\right) \quad \text{if } x \in \mathbb{R}$$


The characterization is simpler if it is built from the statistical moments:

Mean $m_x = E\{x\} = \int_{-\infty}^{\infty} x f_x(x) dx$ Average value

Power $P_x = E\{|x|^2\} = \int_{-\infty}^{\infty} |x|^2 f_x(x) dx$

Cross-correlation $r_{xy} = E\{xy\} = \int_{-\infty}^{\infty} xy f_{xy}(x, y) dx dy$

Variance $\sigma_x^2 = E\{|x - m_x|^2\} = \int_{-\infty}^{\infty} |x - m_x|^2 f_x(x) dx$ Dispersion around the mean



Cross-covariance

$$c_{xy} = E\{(x - E\{x\})(y - E\{y\})\} = \int_{-\infty}^{\infty} (x - E\{x\})(y - E\{y\}) f_{xy}(x, y) dx dy$$




If random variables are independent:

$$\begin{aligned} c_{xy} &= \int_{-\infty}^{\infty} (x - E\{x\})(y - E\{y\}) f_x(x) f_y(y) dx dy = \\ &= \int_{-\infty}^{\infty} (x - E\{x\}) f_x(x) dx \int_{-\infty}^{\infty} (y - E\{y\}) f_y(y) dy = 0 \end{aligned}$$


then, they are uncorrelated. The reverse implication is only valid for Gaussian random variables.

Histogram equalisation

Example 2: Each pixel is interpreted as an observation of a random variable



Cumulative function



$$F_X(x) = \int_{-\infty}^x f_X(\lambda) d\lambda$$


Characterization of a vector random variable

Arrange the random variables in a vector and define a joint density function.

$$\mathbf{x} = \begin{bmatrix} x(1) \\ x(2) \\ \vdots \\ x(d) \end{bmatrix}$$

Example 3: Joint Gaussian probability density function for the elements of \mathbf{x}

$$d=2 \quad f_x(\mathbf{x}) = \frac{1}{\sqrt{(2\pi)^d \det(\mathbf{C})}} \exp\left(-\frac{1}{2}(\mathbf{x} - \mathbf{m})^T \mathbf{C}^{-1} (\mathbf{x} - \mathbf{m})\right) \quad \text{if } \mathbf{x} \in \mathbb{R}^{d \times 1}$$



The first and second order moments are now defined as

Mean $\mathbf{m}_x = E\{\mathbf{x}\} = \begin{bmatrix} E\{x(1)\} \\ E\{x(2)\} \\ \vdots \\ E\{x(d)\} \end{bmatrix}$ Average value

Correlation matrix

$$\mathbf{R}_x = E\{\mathbf{x}\mathbf{x}^T\} = \begin{bmatrix} E\{|x(1)|^2\} & E\{x(1)x(2)\} & \cdots & E\{x(1)x(N)\} \\ E\{x(2)x(1)\} & E\{|x(2)|^2\} & \cdots & E\{x(2)x(N)\} \\ \vdots & \vdots & \ddots & \vdots \\ E\{x(N)x(1)\} & E\{x(N)x(2)\} & \cdots & E\{|x(N)|^2\} \end{bmatrix}$$

Contains all possible correlations and cross-correlations between the elements of vector \mathbf{x}


Covariance matrix

$$\begin{aligned}\mathbf{C}_x &= E\{(\mathbf{x} - \mathbf{m})(\mathbf{x} - \mathbf{m})^T\} = \\ &= \begin{bmatrix} E\{|x(1) - m(1)|^2\} & E\{(x(1) - m(1))(x(2) - m(2))\} & \cdots \\ E\{(x(2) - m(2))(x(1) - m(1))\} & E\{|x(2) - m(2)|^2\} & \cdots \\ \vdots & \vdots & \ddots & \vdots \\ E\{(x(N) - m(N))(x(1) - m(1))\} & \cdots & E\{|x(N) - m(N)|^2\} \end{bmatrix} = \\ &= \mathbf{R}_x - \mathbf{m}\mathbf{m}^T\end{aligned}$$

Contains all possible covariances and cross-covariances between the elements of vector \mathbf{x}

Correlation function


$$\mathbf{R}_x = E\{\mathbf{x}\mathbf{x}^H\} = \begin{bmatrix} E\{|x(1)|^2\} & E\{x(1)x(2)\} & \cdots & E\{x(1)x(N)\} \\ E\{x(2)x(1)\} & E\{|x(2)|^2\} & \cdots & E\{x(2)x(N)\} \\ \vdots & \vdots & \ddots & \vdots \\ E\{x(N)x(1)\} & E\{x(N)x(2)\} & \cdots & E\{|x(N)|^2\} \end{bmatrix}$$


$$\mathbf{r}_x = \begin{bmatrix} r_x(0) \\ r_x(1) \\ \vdots \\ r_x(N-1) \end{bmatrix}$$

Property 1. Hermitian symmetry $r_x(k) = r_x^*(-k)$

Property 2. Its maximum is in $k = 0$, and it is the power of $x(n)$

Cross-correlation

$$\mathbf{R}_{xy} = E\{\mathbf{xy}^T\} = \begin{bmatrix} E\{x(1)y(1)\} & \cdots & E\{x(1)y(k)\} & \cdots & E\{x(1)y(N)\} \\ E\{x(2)y(1)\} & \cdots & E\{x(2)y(k)\} & \cdots & E\{x(2)y(N)\} \\ \vdots & \cdots & \vdots & \ddots & \vdots \\ E\{x(N)y(1)\} & \cdots & E\{x(N)y(k)\} & \cdots & E\{x(N)y(N)\} \end{bmatrix}$$

$$\mathbf{r}_{xy} = E\{\mathbf{xy}(k)\} = \begin{bmatrix} E\{x(1)y(k)\} \\ E\{x(2)y(k)\} \\ \vdots \\ E\{x(N)y(k)\} \end{bmatrix}$$

Cross-correlation is a measure of similarity between random variables: the larger it is, the lower is the error.

$$\begin{aligned} MSE &= E\{|y(n) - x(n+k)|^2\} = \\ &= r_x(0) + r_y(0) - 2E\{x(n+k)y(n)\} = \\ &= r_x(0) + r_y(0) - 2r_{xy}(k) \geq 0 \end{aligned}$$

CONTENTS

1. Random variables
2. Matrix algebra: operators, norms and eigenvectors
3. Optimization with restrictions
4. Derivation of real variable functions

Notation (I)

x : Random variable

$x(n)$: Temporal sequence

$X(f)$: Fourier transform of a temporal sequence

\mathbf{x} , $\underline{\mathbf{x}}$: Column vector

\mathbf{x}^T , $\underline{\mathbf{x}}^T$: Row vector, transpose of \mathbf{x}



$$\mathbf{x} \in C^N ; \mathbf{x} = \begin{bmatrix} x(1) \\ x(2) \\ \vdots \\ x(N) \end{bmatrix}$$

\mathbf{X} , $\underline{\mathbf{X}}$: Matrix

\mathbf{X}^T , $\underline{\mathbf{X}}^T$: Transpose matrix \mathbf{X}

$\mathbf{x}^T \mathbf{y}$: Scalar product between vectors \mathbf{x} and \mathbf{y} $\mathbf{x}^T \mathbf{y} = \sum_{i=0}^{N-1} x(i)y(i)$

Notation (II)

$\mathbf{x} \mathbf{y}^T$: Outer product between vectors \mathbf{x} and \mathbf{y}

$$\mathbf{x} \mathbf{y}^T = \begin{bmatrix} x(1)y(1) & x(1)y(2) & \cdots & x(1)y(N) \\ x(2)y(1) & x(2)y(2) & \cdots & x(2)y(N) \\ \vdots & \vdots & \ddots & \vdots \\ x(N)y(1) & x(N)y(2) & \cdots & x(N)y(N) \end{bmatrix}$$

$\mathbf{C} \mathbf{y}$: Matrix-vector product

$$\mathbf{C} \mathbf{y} = \begin{bmatrix} \mathbf{c}_0^T \\ \mathbf{c}_1^T \\ \vdots \\ \mathbf{c}_{N-1}^T \end{bmatrix} \mathbf{y} = \begin{bmatrix} \sum_{i=0}^{N-1} c(0,i)y(i) \\ \sum_{i=0}^{N-1} c(1,i)y(i) \\ \vdots \\ \sum_{i=0}^{N-1} c(N-1,i)y(i) \end{bmatrix}$$

Norms and Schwarz's inequality

The norm is defined using the scalar product $\langle \cdot, \cdot \rangle$

- Given a scalar product $\langle \cdot, \cdot \rangle$, the norm is $\|\mathbf{v}\| = \sqrt{\langle \mathbf{v}, \mathbf{v} \rangle}$

- Examples... $\langle \mathbf{x}, \mathbf{y} \rangle = \mathbf{y}^T \mathbf{x}$, $\|\mathbf{x}\|_2 = \sqrt{\mathbf{x}^T \mathbf{x}} = \sqrt{\sum_{i=1}^N |x_i|^2}$

$$\langle \mathbf{X}, \mathbf{Y} \rangle = \text{Tr}(\mathbf{Y}^T \mathbf{X}), \quad \|\mathbf{X}\| = \sqrt{\text{Tr}(\mathbf{X}^T \mathbf{X})} = \sqrt{\sum_{i=1}^N \sum_{j=1}^M |x_{i,j}|^2} = \|\mathbf{X}\|_F$$
$$\langle x(t), y(t) \rangle = \int x(t) y(t) dt, \quad \|x(t)\| = \sqrt{\int |x(t)|^2 dt} = \sqrt{E_x}$$
$$\langle X, Y \rangle = E\{XY\}, \quad \|X\| = \sqrt{E\{|X|^2\}} = \sqrt{\sigma_X^2 + m_X^2}$$

Schwarz's inequality (valid in all cases):

$$|\langle \mathbf{x}, \mathbf{y} \rangle|^2 \leq \|\mathbf{x}\|^2 \|\mathbf{y}\|^2$$

Equality condition is satisfied when...

$$|\langle \mathbf{x}, \mathbf{y} \rangle|^2 = \|\mathbf{x}\|^2 \|\mathbf{y}\|^2 \quad \Leftrightarrow \quad \exists k, \quad \mathbf{x} = k \cdot \mathbf{y}$$

Matrix operators

Trace $\mathbf{A} \in \mathbb{R}^{N \times N} \Rightarrow \text{Tr}(\mathbf{A}) = \sum_{i=1}^N [\mathbf{A}]_{i,i}$

$$\mathbf{A} \in \mathbb{R}^{N \times M}, \quad \mathbf{B} \in \mathbb{R}^{M \times N} \Rightarrow \text{Tr}(\mathbf{AB}) = \sum_{i=1}^N [\mathbf{AB}]_{i,i} = \text{Tr}(\mathbf{BA}) = \sum_{i=1}^M [\mathbf{BA}]_{i,i}$$

$$\text{Tr}(\mathbf{ABC}) = \text{Tr}(\mathbf{BCA}) = \text{Tr}(\mathbf{CAB})$$

Determinant

$$\mathbf{A} \in \mathbb{R}^{N \times N}, \quad \mathbf{B} \in \mathbb{R}^{N \times N} \Rightarrow \det(\mathbf{AB}) = \det(\mathbf{A}) \det(\mathbf{B})$$

$$\det(\mathbf{ABC}) = \det(\mathbf{A}) \det(\mathbf{B}) \det(\mathbf{C})$$

Frobenius' (or Euclidean) norm

$$\mathbf{x} \in \mathbb{R}^{N \times 1} \Rightarrow \|\mathbf{x}\|_F = \sqrt{\sum_{i=1}^N |x_i|^2} = \sqrt{\mathbf{x}^T \mathbf{x}} = \sqrt{\text{Tr}(\mathbf{x} \mathbf{x}^T)}$$

$$\mathbf{X} \in \mathbb{R}^{N \times M} \Rightarrow \|\mathbf{X}\|_F = \sqrt{\sum_{i=1}^N \sum_{j=1}^M |x_{i,j}|^2} = \sqrt{\text{Tr}(\mathbf{X}^T \mathbf{X})} = \sqrt{\text{Tr}(\mathbf{X} \mathbf{X}^T)}$$

Inverse of a matrix product

$$\mathbf{A} \in \mathbb{R}^{N \times N}, \quad \mathbf{B} \in \mathbb{R}^{N \times N} \quad \Rightarrow \quad (\mathbf{AB})^{-1} = \mathbf{B}^{-1} \mathbf{A}^{-1}$$

Inversion lemma (Woodbury's identity)

$$(\mathbf{A} + \mathbf{UCV})^{-1} = \mathbf{A}^{-1} - \mathbf{A}^{-1} \mathbf{U} (\mathbf{C}^{-1} + \mathbf{VA}^{-1} \mathbf{U})^{-1} \mathbf{VA}^{-1}$$

... particular case

$$(\mathbf{A} + k\mathbf{u}\mathbf{u}^H)^{-1} = \mathbf{A}^{-1} - \frac{\mathbf{A}^{-1} \mathbf{u} \mathbf{u}^H \mathbf{A}^{-1}}{\frac{1}{k} + \mathbf{u}^H \mathbf{A}^{-1} \mathbf{u}}$$

Eigenvalues and eigenvectors of a matrix

The eigenvectors of a matrix $\mathbf{A} \in \mathbb{R}^{N \times N}$ are those vectors whose norm is not altered after being transformed:

$$\begin{aligned}\mathbf{A}\mathbf{q} &= \lambda\mathbf{q} \\ \Downarrow \\ (\mathbf{A} - \lambda\mathbf{I})\mathbf{q} &= \mathbf{0} \\ \Downarrow \\ P(\lambda) &= \det(\mathbf{A} - \lambda\mathbf{I}) = 0\end{aligned}$$

This equation is a polynomial in λ (characteristic polynomial of \mathbf{A}), that has N roots. Hence, \mathbf{A} has N eigenvalues $\lambda_i \quad i=1,\dots,N$ (possibly multiple).

Not all matrices can be diagonalised using the eigenvectors. Only in those cases where each eigenvalue is associated to a vector space (generated by the eigenvectors) of dimension equal to the multiplicity of the eigenvalue.

Properties of the correlation and covariance matrices

P1. \mathbf{R}_x is symmetric $\mathbf{R} = \mathbf{R}^T$, $\mathbf{R} \in \mathbb{R}^{N \times N}$

P2. \mathbf{R}_x is positive semidefinite: $\mathbf{y}^T \mathbf{R}_x \mathbf{y} \geq 0 \quad \forall \mathbf{y}$

P3. The eigenvalues of \mathbf{R}_x are real and non-negative.

P4. The trace of a matrix is the sum of its eigenvalues.

P5. The eigenvalues of \mathbf{R}_x are bounded by the maximum and the minimum of the power spectral density of $x(n)$.

P6. The eigenvectors of \mathbf{R}_x are orthogonal, so we can write $\mathbf{R}_x = \mathbf{Q}\mathbf{\Lambda}\mathbf{Q}^T$

P7. $\mathbf{R}_x^{-1} = \mathbf{R}_x^{-T}$

Symmetric matrices

A square matrix is symmetric if $\mathbf{R} = \mathbf{R}^T$, $\mathbf{R} \in \mathbb{R}^{N \times N}$

Spectral decomposition theorem

A symmetric matrix \mathbf{R} can always diagonalize using a base of orthonormal eigenvectors with real eigenvalues.

$$\mathbf{R} = \mathbf{Q}\mathbf{\Lambda}\mathbf{Q}^T, \quad \lambda_i \mathbf{q}_i = \mathbf{R}\mathbf{q}_i, \quad \lambda_i \in \mathbb{R}, \quad \|\mathbf{q}_i\| = 1$$

$$\mathbf{Q} = [\mathbf{q}_1 \quad \mathbf{q}_2 \quad \cdots \quad \mathbf{q}_N] \in \mathbb{R}^{N \times N}, \quad \mathbf{q}_i \in \mathbb{R}^{N \times 1}, \quad \mathbf{Q}\mathbf{Q}^T = \mathbf{Q}^T\mathbf{Q} = \mathbf{I}, \quad \mathbf{Q}^{-1} = \mathbf{Q}^T$$

$$\mathbf{\Lambda} = \text{diag}[\lambda_1, \lambda_2, \dots, \lambda_N] \quad \text{with } \lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_N$$

Positive semidefinite matrices: $\mathbf{v}^T \mathbf{R} \mathbf{v} \geq 0, \quad \forall \mathbf{v} \neq \mathbf{0} \in \mathbb{R}^{N \times 1}$

- ✓ A symmetric matrix is positive semidefinite if all eigenvalues are positive: $\lambda_i \geq 0, \quad \forall i$

CONTENTS

1. Random variables
2. Matrix algebra: operators, norms and eigenvectors
3. Optimization with restrictions
4. Derivation of real variable functions

Constrained optimization

Maximization or minimization of $f(\mathbf{x})$ with restrictions on $\mathbf{x} \in \mathbb{R}^{M \times 1}$:

$$\text{optimize } f(\mathbf{x}) \text{ with } g_i(\mathbf{x}) \leq 0 \text{ and/or } g_j(\mathbf{x}) = 0 \quad i, j = 1, \dots, K$$

Computation of a solution based on the method of Lagrange multipliers:

1. Build the Lagrangian function:

$$L(\mathbf{x}, \{\mu_i\}) = f(\mathbf{x}) + \sum_{i=1}^K \mu_i g_i(\mathbf{x})$$


2. Equate the gradient to zero and solve the equation to obtain the possible values of \mathbf{x} , that depend on the Lagrange multipliers λ_i

$$\nabla_{\mathbf{x}} L(\mathbf{x}, \{\mu_i\}) = \nabla_{\mathbf{x}} f(\mathbf{x}) + \sum_{i=1}^K \mu_i \nabla_{\mathbf{x}} g_i(\mathbf{x}) = \mathbf{0}$$

3. Select the solution among the possible values of \mathbf{x} for which $f(\mathbf{x})$ is maximum/minimum and all restrictions are satisfied simultaneously.

Constrained optimization

The derivation of the Lagrangian provides the sufficient condition if the function is concave/convex. As an illustration: when only one restriction is set, in the optimum point, the gradient of $f(\mathbf{x})$ and $g(\mathbf{z})$ are parallel...



CONTENTS

1. Random variables
2. Matrix algebra: operators, norms and eigenvectors
3. Optimization with restrictions
4. Derivation of real variable functions

Derivation of scalar functions of real variables

In many optimization problems we are interested in determining the gradient of a real scalar function with a vector variable:

$$\nabla_{\mathbf{h}} f(\mathbf{h}) = \begin{bmatrix} \frac{\partial f(\mathbf{h})}{\partial h(1)} \\ \vdots \\ \frac{\partial f(\mathbf{h})}{\partial h(L)} \end{bmatrix}$$

Some useful cases are...

$$\nabla_{\mathbf{h}} \mathbf{a}^T \mathbf{h} = \mathbf{a}$$

$$\nabla_{\mathbf{h}} \mathbf{h}^T \mathbf{a} = \mathbf{a}$$

$$\nabla_{\mathbf{h}} \mathbf{h}^T \mathbf{h} = 2\mathbf{h}$$

$$\nabla_{\mathbf{h}} \mathbf{h}^T \mathbf{R} \mathbf{h} = \begin{cases} 2\mathbf{R} \mathbf{h} & \text{si } \mathbf{R}^T = \mathbf{R} \\ (\mathbf{R} + \mathbf{R}^T) \mathbf{h} & \text{si } \mathbf{R}^T \neq \mathbf{R} \end{cases}$$