


Linear Regression Algorithm:

Linear regression is a supervised machine learning algorithm used to find a linear relationship between a dependent (y) and one or more independent (x) variables, hence called linear regression



Types of Linear Regression

1. Simple Linear Regression:

If a single independent variable is used to predict the value of a numerical dependent variable, then such a Linear Regression algorithm is called Simple Linear Regression. $y = mx + c$

2. Multiple Linear regression:

If more than one independent variable is used to predict the value of a numerical dependent variable, then such a Linear Regression algorithm is called Multiple Linear Regression.

$y = m_1x_1 + m_2x_2 + \dots + m_Nx_N + c$
y = Dependent Variable (Target Variable)
x = Independent Variable (predictor Variable)
c = intercept of the line (Gives an additional degree of freedom)
m = Linear regression coefficient (scale factor to each input value).

Loss Function(Cost Function):

- The loss is the error in our predicted value of m and c. Our goal is to minimize this error to obtain the most accurate value of m and c.
- We will use the Mean Squared Error function to calculate the loss.

$$E = \frac{1}{n} \sum_{i=0}^n (y_i - \bar{y}_i)^2$$

$$E = \frac{1}{n} \sum_{i=0}^n (y_i - (mx_i + c))^2$$

Gradient Descent Algorithm:

- 1 • Gradient descent is an iterative optimization algorithm to find the minimize the
- 2 Loss Function.
- 3 • Gradient descent is a method of updating m and c values to minimize the cost
- 4 function (MSE) and to get the best fit line(regression line)
- 5 • Best Fit Line: When working with linear regression, our main goal is to find the
- 6 best fit line which means the error between predicted values and actual values
- 7 should be minimized. The best fit line will always have the least Mean squares
- 8 error.
- 9 • A regression model uses gradient descent to update the coefficients of the line
- 10 (m and c) by reducing the cost function by a random selection of coefficient
- 11 values and then iteratively updating the values to reach the minimum cost
- 12 function.
- 13 • To update m and c, we take gradients from the cost function. To find these
- 14 gradients, we take partial derivatives for m and c.

$$D_m = \frac{1}{n} \sum_{i=0}^n 2(y_i - (mx_i + c))(-x_i)$$

$$D_m = \frac{-2}{n} \sum_{i=0}^n x_i(y_i - \bar{y}_i)$$

- 1 • Calculate the partial derivative of the loss function with respect to m , and plug in the current values of x , y , m , and c in it to obtain the derivative value D .

$$D_c = \frac{-2}{n} \sum_{i=0}^n (y_i - \bar{y}_i)$$




○

$$m = m - L \times D_m$$

$$c = c - L \times D_c$$

○

- 1 • Global minima: It is a point that obtains the absolute lowest value of our function
- 2 • Learning Rate: It determines the size of the steps that are taken by the gradient descent algorithm
- 3 • If α (Learning Rate) is very small, it would take a long time to converge and become computationally expensive.
- 4 • If α is large, it may fail to converge and overshoot the minimum.
- 5 • The most commonly used rates are : 0.001, 0.003, 0.01, 0.03, 0.1, 0.3.



• Gradient Descent variants:

There are three types of gradient descent methods based on the amount of data used to calculate the gradient:

1. Batch gradient descent
2. Stochastic gradient descent
3. Mini-batch gradient descent

Assumptions of Linear Regression:

1. Linearity: Relationship between the independent and dependent variables to be linear.
2. No Multicollinearity (Independence): Observations are independent of each other.
3. Normality of Residual
4. Homoscedasticity: The variance of residual is the same for any value of X.

1. Linearity:

Relationship between the independent and dependent variables to be linear.

1.1 How to check Linearity:

1. Coefficient of correlation($R=?$)
2. Scatter Plot
3. Correlation matrix

1.2 How to Handle Linearity if get violated:

Apply a nonlinear transformation to the independent variable.

1. Log transformation
2. Square root transformation
3. cube.root transformation.
3. Reciprocal transformation

1.1.1 Coefficient of correlation(R): *{Pearson corr.}*

Correlation coefficients are used to measure how strong a relationship is between two variables. There are several types of correlation coefficients, but the most popular is Pearson's.

$$\text{Correlation Coefficient}(r) = \frac{\text{Covariance}(x,y)}{\text{Std dev}(x) * \text{Std dev}(y)}$$

$$r = \frac{\sum (x_i - \bar{x})(y_i - \bar{y})}{\sqrt{\sum (x_i - \bar{x})^2 \sum (y_i - \bar{y})^2}}$$

Where,




r = Pearson Correlation Coefficient

x_i = x variable samples

y_i = y variable sample

\bar{x} = mean of values in x variable

\bar{y} = mean of values in y variable



- 1 The range of R-Value is between -1 to +1
- 2 $R = 1$ >> indicates a strong positive relationship. $R = -1$ >> indicates a strong negative relationship.
- 3
- 4 $R = -1$ >> indicates a strong negative relationship. if one variable increases, other variable decreases
- 5
- 6 $R = 0$ >> It means there is no linear relationship. It doesn't mean that there is no relationship

2. No Multicollinearity:

- Multicollinearity (or collinearity) occurs when one independent variable in a regression model is linearly correlated with another independent variable.
- This means that an independent variable can be predicted from another independent variable in a regression model
- Multicollinearity can be a problem in a regression model because we would not be able to distinguish between the individual effects of the independent variables on the dependent
- $Y = M_1X_1 + M_2X_2 + C$
- Coefficient M_1 is the increase in Y for a unit increase in X_1 while keeping X_2 constant. But since X_1 and X_2 are highly correlated, changes in X_1 would also cause changes in X_2 and we would not be able to see their individual effect on Y .
- Multicollinearity may not affect the accuracy of the model as much. But we might lose reliability in determining the effects of individual features in your model – and that can be a problem when it comes to interpretability.

2.1 How to detect Multicollinearity:

1. VIF (Variable Inflation Factors):

The VIF score of an independent variable represents how well the variable is explained by other independent variables.

VIF = 1 → No correlation

VIF = 1 to 5 → Moderate correlation

VIF >10 → High correlation

2. Correlation matrix / Correlation plot

3. Scatter plots

2.1 How to handle Multicollinearity

1. Dropping variables
2. Combining variables

3. Normality of the residuals

Residuals: The difference between the actual y value and the estimated y value

Residuals = $(Y_a - Y_p)$

A normal distribution has some important properties:

1. The mean, median, and mode all represent the center of the distribution.
2. the distribution is a bell shape
3. ≈68% of the data falls within 1 standard deviation of the mean, ≈95% of the data falls within 2 STD of the mean and ≈99.7% of the data falls within 3 STD of the mean

3.1 How to check normality:

1. Graphs for Normality test:

1. Distribution curve, Histogram (sns. distplot, sns.kdeplot)
2. Q-Q or Quantile-Quantile Plot

2. Statistical Tests for Normality(Hypothesis Testing):

1. Shapiro-Wilk test
2. Kolmogorov-Smirnov test
3. D'Agostino's K-squared test

4. Homoscedasticity:

- Residuals have constant variance at every level of x . This is known as homoscedasticity. When this is not the case, the residuals are said to suffer from heteroscedasticity.
- When heteroscedasticity is present in a regression analysis, the results of the analysis become hard to trust

4.1 How to Check Homoscedasticity:

1. Scatter plot between fitted value and residual plot.

4.2 How to handle :

1. Transform the dependent variable(Y): log transformation of the dependent variable
2. Redefine the dependent variable
3. Use weighted regression: This type of regression assigns a weight to each data point based on the variance of its fitted value.

Advantages:

1. Simple to implement and easier to interpret the output coefficients.
2. When you know the dependent and independent variables have a linear relationship, this algorithm is the best to use because it's less complex as compared to other algorithms.
3. Linear Regression is prone to over-fitting but it can be avoided using some dimensionality reduction techniques, regularization (L1 and L2) techniques, and cross-validation.

❖ Disadvantages:

1. If the independent features are correlated it may affect performance.
2. it is only efficient for linear data(High Corr between x and Y)
3. Sometimes a lot of feature engineering is required.
4. Scaling is Required: predictors have a mean of 0.
5. It is often quite prone to noise and overfitting.
6. It is sensitive to missing values.
7. It is sensitive to Outliers

❖ Applications:

❖ Evaluation Metrics for Linear Regression:

1. Mean Absolute Error(MAE): It is most Robust to outliers.
2. Mean Squared Error(MSE)
3. Root Mean Squared Error(RMSE)
4. R-squared or Coefficient of Determination:
 - a. SSE(Sum of Squared Error
 - b. SSR(Sum of Squares due to Regression)
 - c. SST(Sum of Squares Total or Total Error)
5. Adjusted R Squared

$$\text{Variance} = \sum_{i=1}^n (y_i - \bar{y})^2 \rightarrow \text{SST (Sum of squares of total)}$$


$$\text{SSE} = \sum_{i=1}^n (y_i - \hat{y}_i)^2 \rightarrow \text{Sum of squares of Errors – Unexplained Variance}$$

$$\text{SSR} = \sum_{i=1}^n (\hat{y}_i - \bar{y})^2 \rightarrow \text{Sum of Squares of Regression – Explained Variance}$$

$$\text{SST} = \text{SSR} + \text{SSE}$$

y_i - Actual value of y
 \bar{y} - Mean value of y
 \hat{y}_i - Predicted value of y

$$R^2 = \frac{\text{SSR}}{\text{SST}} \quad \text{Or} \quad R^2 = 1 - \frac{\text{SSE}}{\text{SST}}$$



In []:

1