



University
of Glasgow


Virus evolution and molecular clocks

**WORLD
CHANGING
GLASGOW**

**A WORLD
TOP 100
UNIVERSITY**

Virus evolution

- Changes in the genetic sequences of viruses over time
- Can occur through various mechanisms (e.g. mutation & recombination), leading to the emergence of new virus variants
- Why should we care?






Molecular clocks

- Molecular sequence data from rapidly evolving organisms are often sampled at different points in time
- Sampling times can then be used for molecular clock calibration
- The root-to-tip (RTT) regression is an essential tool to assess the degree to which the data behave in a clock-like fashion
- Types of molecular clocks:
 - ✓ Strict vs relaxed
 - ✓ Local vs global



The strict molecular clock


- A strict molecular clock
 - ✓ The simplest molecular clock model
 - ✓ Assumes a constant evolutionary rate (rate of substitution per unit time)
- When the evolutionary rate is constant throughout a phylogenetic tree, the term ***global molecular clock*** is used



Under the strict clock model, evolution occurs at the same rate at every branch.

The strict molecular clock

- **A strict local clock** - different substitution rates apply to different monophyletic groups within a tree
 - ✓ sampling from different host populations, host species, or pathogen lineages
- Larger datasets are more likely to sample from distinct populations, making local clocks increasingly important




Under the fixed local clock model, a shift in evolutionary rate occurs at the TMRCA of each defined taxon set.



The relaxed molecular clock

- Multivariate statistical approach i.e. a different rate for each branch
- Computationally demanding




Under the uncorrelated relaxed clock model, the evolutionary rate at each branch is drawn from an underlying distribution.



Clockor2

- A scalable and accessible client-side web application for exploring the fit of local clocks, with results available in seconds to minutes
- Enables rapid inference of global and local strict molecular clocks from phylogenetic trees using root-to-tip regression (RTT)



Clockor2

Inferring global and local strict molecular clocks using root-to-tip regression



University
of Glasgow

Input

- A newick tree
- Sampling dates & group identifiers
(parsed from tip labels or separate files
on input)



The Best Fitting Root

- Based on the R^2 or residual mean square (RMS) of the RTT regression of a global clock model for the input tree
- The root position along the branch leading to the highest R^2 or RMS is selected
- The best-fitting root is inferred using a single global clock because it presents the most parsimonious model of the evolutionary rate for a given tree
- Clockor2 does not find the best-fitting root for local clock models because the search space of best-fitting roots and local clock configurations quickly becomes prohibitive and is possibly unidentifiable



Testing the fit of a clock

- Local clock and or global clock configurations can also be compared using an information criterion
 - ✓ Bayesian Information Criterion (BIC)
 - ✓ Akaike Information Criterion (AIC)
 - ✓ corrected Akaike Information Criterion (AICc)

Local Clock

AICc -585.68

AIC -592.32

BIC -567.18

CLOCK	#TIPS	RATE	R^2	RESIDUAL MEAN SQUARED	INTERCEPT
Bauchi	14	5.003×10^{-4}	0.913	8.234×10^{-7}	1945.074



University
of Glasgow

Let's put Clockor2 to work!

#UofGWorldChangers



@UofGlasgow