

24-02-22

Machine learning.
CSPE-65
Cycle test - 1

①
106119100
Rajneesh.

Question ①

(A) supervised.

1

12.5

Question ②

(i) values $\rightarrow 4, 7, 9, 8, 12, 80, 15$

Ascending order = $4, 7, 8, 9, 12, 15, 80$
 $\downarrow \quad \downarrow \quad \downarrow$
 $Q_1 \quad Q_2 \quad Q_3$

as, 1st quantile = 7

2nd quantile = 9

3rd quantile = 15.

smallest = 4

highest = 80.

Inter-Quantile Range

$$IQR = Q_3 - Q_1 = 15 - 7$$

$IQR = 8$


lower whisker

$$= Q_1 - 1.5 * IQR = 7 - \frac{3}{2} * 8$$


$= -5$

upper whisker

$$= Q_3 + 1.5 * IQR = 15 + \frac{3}{2} * 8$$
$$= 27.$$



(ii)



Data distribution → 25% (15 and 27)
 → 25% of data b/w 4-7
 → 50% of data b/w 7 and 15

Mean.

$$= \frac{4+7+9+8+12+80+15}{7} = 19.28.$$

Median = 9

∴, mean > Median

∴, the data is positively-skewed
 or Right-skewed.

$Q_1 = 7$

$Q_2 = 9$

$Q_3 = 15$

and $IQR = 8$

and

min = 4.

but max = 80

∴ data have outlier point.

Question (3)

(3)

(D) $A \rightarrow (iv)$; $B \rightarrow (iii)$; $C \rightarrow (ii)$; $D \rightarrow (i)$

Question (4)

(i)

Binning methods(a) unsupervised Binning

1) Equal width binning

2) frequency width binning

(b) supervised Binning

n Entropy based Binning.

(ii)

value : 15, 21, 45, 6, 11, 17, 45, 19, 12, 4, 9, 5

sorting : 4, 5, 6, 9, 11, 12, 15, 17, 19, 21, 45, 45.

(a) Apply Equal width binning :

Equal width binning.

 $\text{len}(\text{value}) = 12$

$$w = (\text{max} - \text{min}) / 3 = \frac{45 - 4}{3} = 13.67$$

bin 1 - range (4, 17.67)

bin 2 - range (17.67, 31.33)

bin 3 - range (31.33, 45).

Bin 1 : [4, 5, 6, 9, 11, 12, 15, 17]

Bin 2 : [19, 21]

Bin 3 : [45, 45]

106119100

(4)

(ii) Binning by frequency (equal frequency)

Bin-1 [4, 5, 6]

Bin size = 3

Bin 2 - [9, 11, 12]

Bin 3 - [15, 17, 19]

Bin-4 [21, 45, 45]

1.5

Question (5)

(i)

As we know, feature scaling. ~~Because~~,
require when data is missing, to remove
outlier, or pruned redundant rows

Yes, the feature "Age" in the
data set requires feature
scaling, Because

the other dataset are 25, 35, 30, 20
with differences of min 5 & max 15.

but the age value = 1000.

⑤

the difference between the values are much higher which is outlier point in the data.

we need to make the data in same scale so, that each feature is equally important.

(ii) using Pearson correlation

Speed (s)

Covariance

$$\bar{s} = \frac{\sum s}{n} = \frac{100+120+140+160+180}{5} = 140.$$

$$\begin{aligned} \sigma_s &= \sqrt{\frac{\sum (\bar{s} - s)^2}{n}} = \sqrt{\frac{(40)^2 + 20^2 + 20^2 + 20^2}{5}} \\ &= 10\sqrt{\frac{40}{5}} = \boxed{\sigma_s = 20\sqrt{2}} \end{aligned}$$

Acceleration (a)

$$\bar{a} = \frac{\sum a}{n} = \frac{60+70+75+85+100}{5} = \frac{390}{5} = 78.$$

$$\begin{aligned} \sigma_a &= \sqrt{\frac{18^2 + 8^2 + 3^2 + 7^2 + 22^2}{5}} = \sqrt{\frac{930}{5}} \\ &= \boxed{\sqrt{186} = \sigma_a} \end{aligned}$$

Covariance (s, a)

$$= \frac{1}{n} \sum_{i=1}^n (s_i - \bar{s}) * (a_i - \bar{a})$$

$$= \frac{1}{5} ((-40)(-18) + (-20)(-8) + 0(8) + 20(7) + 40(22))$$

$$= \frac{1900}{5} = 380 = \text{cov}(s, a)$$

Pearson correlation

$$P_{s,a} = \frac{\text{cov}(s, a)}{\sigma_s \sigma_a} = \frac{380}{20\sqrt{2} \times \sqrt{186}}$$

$$= \frac{380}{20\sqrt{4 \times 93}} = \frac{19}{2\sqrt{93}}$$

$$= 19/19.2873$$

$$P_{s,a} = 0.98510$$

\approx nearly 1.

Rough plot


→ since ρ is positive, they change in the same direction.

→ since $\rho \approx 1$, they are strongly correlated.

Rough plot

106119100

⑦



$p \approx 1$, so most point
on the line

check