

Final Exam – Linear Algebra III, 1MA026

Uppsala University
Department of Mathematics

2022-03-17
duration of the exam: 5 hours

Please write your answers in **English**. There are 8 problems in this exam, and each one is worth 5 points. The grade limits are: 18 points for grade 3, 25 points for grade 4 and 32 points for grade 5. **You need to motivate every step in your solution to get the full score on each question.** Good luck!

In the exercises below, F is an arbitrary field (unless otherwise specified) and V is a vector space over F .

1. Let $V = \mathbb{C}^3$, with the standard hermitian inner product. Let $U = \text{Span}(v_1, v_2)$, where

$$v_1 = (1, i, 2) \quad v_2 = (1, 0, -i).$$

- (a) Find a basis of the orthogonal complement U^\perp .
(b) Find a vector subspace $W \subset V$ such that $W \neq U^\perp$ and $W \oplus U = V$.

2. Consider the derivative operator $D \in \mathcal{L}(\mathcal{P}_n(F))$ defined by

$$D(a_0 + a_1 x + a_2 x^2 + \dots + a_n x^n) = a_1 + 2a_2 x + \dots + na_n x^{n-1}.$$

Note that this makes sense for every integer $n \geq 0$ and every field F .

- (a) Show that for every fixed $a \in F$, the vectors

$$v_0 = 1, v_1 = x - a, v_2 = (x - a)^2, \dots, v_n = (x - a)^n$$

form a basis B_a of the $(n+1)$ -dimensional vector space $\mathcal{P}_n(F)$.

- (b) Suppose that the characteristic of F , $\text{char}(F)$, is either 0 or greater than n . Show that the dual basis $\varphi_0, \dots, \varphi_n$ of $(\mathcal{P}_n(F))'$ that corresponds to the basis B_a is such that for every $0 \leq k \leq n$

$$\varphi_k(p(x)) = \frac{1}{k!} D^k(p(x))(a).$$

$(D^k(p(x)))(a)$ means: apply D to $p(x)$ k many times, then evaluate at $x = a$.)

3. Let V_1 and V_2 be the subsets of \mathbb{R}^3 given by

$$V_1 = \{(x, y, z) \in \mathbb{R}^3 \mid x - y + z = 0\}$$

and

$$V_2 = \{(x, y, z) \in \mathbb{R}^3 \mid x - y + z = 2\}.$$

- (a) Is V_1 a vector subspace of \mathbb{R}^3 ? If yes, then find an orthonormal basis of V_1 .
(b) Is V_2 a vector subspace of \mathbb{R}^3 ? If yes, then find an orthonormal basis of V_2 .
(c) Find the point in V_1 that is the closest to the point $(1, 0, 0)$.
(d) Find the point in V_2 that is the closest to the point $(1, 0, 0)$.

4. Consider the matrix

$$A = \begin{pmatrix} 2 & 1 & 0 \\ 0 & 1 & -1 \\ 0 & 2 & 4 \end{pmatrix}.$$

Find a Jordan matrix J and an invertible matrix C such that $J = C^{-1}AC$.

5. Let $V = \mathbb{R}^2$ and let $B : V \times V \rightarrow \mathbb{R}$ be the bilinear form given by

$$B((x_1, y_1), (x_2, y_2)) = 2x_1x_2 - 5y_1x_2 - x_1y_2 + 4y_1y_2.$$

- (a) Is B symmetric? Is B alternating? Justify.
 - (b) Let e_1, e_2 be the standard basis of \mathbb{R}^2 . Write the corresponding basis of $V \otimes V$.
 - (c) Is there a linear map $f : V \otimes V \rightarrow \mathbb{R}$ such that $f(v \otimes w) = B(v, w)$ for all $v, w \in \mathbb{R}^2$? Justify. If the answer is affirmative, then write the images under f of the vectors in a basis of $V \otimes V$ (for example, you can use the basis that you wrote in the previous answer).
6. Let $F = \mathbb{Z}/7\mathbb{Z}$ denote the field with 7 elements. Let $T \in \mathcal{L}(F^3, F^2)$ be given by $T(x, y, z) = (2x + 4y + 6z, x + y + z)$. Find a basis of the null space $\text{null}(T)$ and compute the number of elements in the quotient space $F^3/\text{null}(T)$.

7. Let $T \in \mathcal{L}(\mathbb{R}^3)$ be a linear operator whose matrix with respect to the standard basis of \mathbb{R}^3 is

$$A = \begin{pmatrix} -1 & 0 & 1 \\ 1 & -1 & 0 \\ 0 & 1 & -1 \end{pmatrix}.$$

Let $S \in \mathcal{L}(\mathbb{C}^3)$ be a linear operator whose matrix with respect to the standard basis of \mathbb{C}^3 is also A .

In each of the following questions, even if the answer is affirmative, you are not required to find the orthonormal basis or the corresponding diagonal matrix.

- (a) Consider the standard inner product in \mathbb{R}^3 . Can you find an ordered orthonormal basis of \mathbb{R}^3 with respect to which T is diagonal? Explain why.
- (b) Consider the standard inner product in \mathbb{C}^3 . Can you find an ordered orthonormal basis of \mathbb{C}^3 with respect to which S is diagonal? Explain why.

8. Let V be a finite dimensional vector space over F . The minimal polynomial of $T \in \mathcal{L}(V)$ is the monic polynomial $m_T(x) \in \mathcal{P}(F)$ of smallest possible degree, for which $m_T(T) = 0$. You may use the following two facts without proving them:

- Given a choice of ordered basis of V , let A be the matrix representing T with respect to that basis. Let $p(x) \in \mathcal{P}(F)$. Then, $p(T) = 0$ iff $p(A) = 0$.
- If $p(x) \in \mathcal{P}(F)$ is any polynomial such that $p(T) = 0$, then $p(x) = m_T(x) \cdot q(x)$ for some $q(x) \in \mathcal{P}(F)$ (equivalently: m_T divides p).

In this exercise, let $F = \mathbb{C}$.

- Show that every eigenvalue of T is a root of $m_T(x)$.
- Show that every root of $m_T(x)$ is an eigenvalue of T .
- Let $(V, \langle \cdot, \cdot \rangle)$ be a finite dimensional complex inner product space and let $T \in \mathcal{L}(T)$ be a normal operator. Show that

$$m_T(x) = (x - \lambda_1) \cdots (x - \lambda_m),$$

where $\lambda_1, \dots, \lambda_m$ are the distinct eigenvalues of T .