

Formula sheet for Statistical Machine Learning

Warning: This is not a complete list of formulas used in the course, some exam problems may require expressions not listed here. Furthermore, the formulas below are not self-explanatory, you need to be familiar with the expressions to interpret them.

The Gaussian distribution: The probability density function of the p -dimensional Gaussian distribution with mean vector $\boldsymbol{\mu}$ and covariance matrix $\boldsymbol{\Sigma}$ is

$$\mathcal{N}(\mathbf{x} | \boldsymbol{\mu}, \boldsymbol{\Sigma}) = \frac{1}{(2\pi)^{p/2} \sqrt{\det \boldsymbol{\Sigma}}} \exp \left(-\frac{1}{2} (\mathbf{x} - \boldsymbol{\mu})^\top \boldsymbol{\Sigma}^{-1} (\mathbf{x} - \boldsymbol{\mu}) \right), \quad \mathbf{x} \in \mathbb{R}^p.$$

Sum of identically distributed variables: For identically distributed random variables $\{z_i\}_{i=1}^n$ with mean μ , variance σ^2 and average correlation between distinct variables ρ , it holds that $\mathbb{E} \left[\frac{1}{n} \sum_{i=1}^n z_i \right] = \mu$ and $\text{Var} \left(\frac{1}{n} \sum_{i=1}^n z_i \right) = \frac{1-\rho}{n} \sigma^2 + \rho \sigma^2$.

Linear regression and regularization:

- The least-squares estimate of $\boldsymbol{\theta}$ in the linear regression model

$$y = \theta_0 + \sum_{j=1}^p \theta_j x_j + \epsilon$$

is given by the solution $\hat{\boldsymbol{\theta}}_{\text{LS}}$ to the normal equations $\mathbf{X}^\top \mathbf{X} \hat{\boldsymbol{\theta}}_{\text{LS}} = \mathbf{X}^\top \mathbf{y}$, where

$$\mathbf{X} = \begin{bmatrix} 1 & -\mathbf{x}_1^\top \\ 1 & -\mathbf{x}_2^\top \\ \vdots & \vdots \\ 1 & -\mathbf{x}_n^\top \end{bmatrix} \text{ and } \mathbf{y} = \begin{bmatrix} y_1 \\ y_2 \\ \vdots \\ y_n \end{bmatrix} \text{ from the training data } \mathcal{T} = \{\mathbf{x}_i, y_i\}_{i=1}^n$$

- Ridge regression uses the regularization term $\lambda \|\boldsymbol{\theta}\|_2^2 = \lambda \sum_{j=0}^p \theta_j^2$.
The ridge regression estimate is $\hat{\boldsymbol{\theta}}_{\text{RR}} = (\mathbf{X}^\top \mathbf{X} + \lambda \mathbf{I})^{-1} \mathbf{X}^\top \mathbf{y}$.
- LASSO uses the regularization term $\lambda \|\boldsymbol{\theta}\|_1 = \lambda \sum_{j=0}^p |\theta_j|$.

Maximum likelihood: The maximum likelihood estimate is given by

$$\hat{\boldsymbol{\theta}}_{\text{ML}} = \arg \max_{\boldsymbol{\theta}} \ln \ell(\boldsymbol{\theta})$$

where $\ln \ell(\boldsymbol{\theta}) = \sum_{i=1}^n \ln p(y_i | \mathbf{x}_i; \boldsymbol{\theta})$ is the log-likelihood function (the last equality holds when the n training data points are modeled to be independent).

Logistic regression: The logistic regression combines linear regression with the logistic function to model the class probability

$$p(y = 1 | \mathbf{x}_i) = \frac{e^{\boldsymbol{\theta}^\top \mathbf{x}_i}}{1 + e^{\boldsymbol{\theta}^\top \mathbf{x}_i}}.$$

For multi-class logistic regression we use the *softmax* function and model

$$p(y = m | \mathbf{x}_i) = \frac{e^{\boldsymbol{\theta}_m^\top \mathbf{x}_i}}{\sum_{j=1}^M e^{\boldsymbol{\theta}_j^\top \mathbf{x}_i}}.$$

Discriminant Analysis: The linear discriminant analysis (LDA) classifier models $p(y | \mathbf{x})$ using Bayes' theorem and the following assumptions

$$p(y = m | \mathbf{x}) = \frac{p(\mathbf{x} | m)p(y = m)}{\sum_{j=1}^M p(\mathbf{x} | j)p(y = j)} = \frac{\mathcal{N}(\mathbf{x} | \hat{\boldsymbol{\mu}}_m, \hat{\boldsymbol{\Sigma}}) \hat{\pi}_m}{\sum_{j=1}^M \mathcal{N}(\mathbf{x} | \hat{\boldsymbol{\mu}}_j, \hat{\boldsymbol{\Sigma}}) \hat{\pi}_j},$$

where

$$\begin{aligned} \hat{\pi}_m &= n_m / n \text{ for } m = 1, \dots, M \\ \hat{\boldsymbol{\mu}}_m &= \frac{1}{n_m} \sum_{i: y_i = m} \mathbf{x}_i \text{ for } m = 1, \dots, M \\ \hat{\boldsymbol{\Sigma}} &= \frac{1}{n - M} \sum_{m=1}^M \sum_{i: y_i = m} (\mathbf{x}_i - \hat{\boldsymbol{\mu}}_m)(\mathbf{x}_i - \hat{\boldsymbol{\mu}}_m)^\top. \end{aligned}$$

For quadratic discriminant analysis (QDA), the model is

$$p(y = m | \mathbf{x}) = \frac{\mathcal{N}(\mathbf{x} | \hat{\boldsymbol{\mu}}_m, \hat{\boldsymbol{\Sigma}}_m) \hat{\pi}_m}{\sum_{j=1}^M \mathcal{N}(\mathbf{x} | \hat{\boldsymbol{\mu}}_j, \hat{\boldsymbol{\Sigma}}_j) \hat{\pi}_j},$$

where $\hat{\boldsymbol{\mu}}_m$ and $\hat{\pi}_m$ are as for LDA, and

$$\hat{\boldsymbol{\Sigma}}_m = \frac{1}{n_m - 1} \sum_{i: y_i = m} (\mathbf{x}_i - \hat{\boldsymbol{\mu}}_m)(\mathbf{x}_i - \hat{\boldsymbol{\mu}}_m)^\top.$$

Classification trees: The cost function for tree splitting is $\sum_{\ell=1}^{|T|} n_{\ell} Q_{\ell}$ where T is the tree, $|T|$ the number of terminal nodes, n_{ℓ} the number of training data points falling in node ℓ , and Q_{ℓ} the impurity of node ℓ . Three common impurity measures for splitting classification trees are:

$$\begin{aligned} \text{Misclassification error:} \quad Q_{\ell} &= 1 - \max_m \hat{\pi}_{\ell m} \\ \text{Gini index:} \quad Q_{\ell} &= \sum_{m=1}^M \hat{\pi}_{\ell m} (1 - \hat{\pi}_{\ell m}) \\ \text{Entropy/deviance:} \quad Q_{\ell} &= - \sum_{m=1}^M \hat{\pi}_{\ell m} \log \hat{\pi}_{\ell m} \end{aligned}$$

where $\hat{\pi}_{\ell m} = \frac{1}{n_{\ell}} \sum_{i: \mathbf{x}_i \in R_{\ell}} \mathbb{I}(y_i = m)$

Loss functions for classification: For a binary classifier expressed as $\hat{y}(\mathbf{x}) = \text{sign}\{C(\mathbf{x})\}$, for some real-valued function $C(\mathbf{x})$, the margin is defined as $y \cdot C(\mathbf{x})$ (note the convention $y \in \{-1, 1\}$ here). A few common loss functions expressed in terms of the margin, $L(y, C(\mathbf{x}))$ are,

$$\begin{aligned} \text{Exponential loss:} \quad L(y, c) &= \exp(-yc). \\ \text{Hinge loss:} \quad L(y, c) &= \begin{cases} 1 - yc & \text{for } yc < 1, \\ 0 & \text{otherwise.} \end{cases} \\ \text{Binomial deviance:} \quad L(y, c) &= \log(1 + \exp(-yc)). \\ \text{Huber-like loss:} \quad L(y, c) &= \begin{cases} -yc & \text{for } yc < -1, \\ \frac{1}{4}(1 - yc)^2 & \text{for } -1 \leq yc \leq 0, \\ 0 & \text{otherwise.} \end{cases} \\ \text{Misclassification loss:} \quad L(y, c) &= \begin{cases} 1 & \text{for } yc < 0, \\ 0 & \text{otherwise.} \end{cases} \end{aligned}$$