

UPPSALA UNIVERSITY  
 Department of Mathematics  
 © Docent ÖS

Markov Processes, 1MS012  
 Spring semester 2024

## Lecture 10 Markov Processes, 1MS012

### 1 Markov processes (continuous time)

**Definition:** A random process  $(X_t)_{t \geq 0}$  with countable state space (w.l.o.g. a set of integers) is a (discrete) **Markov process** if, for any  $n$  and time-points  $t_1 < \dots < t_{n+1}$  and states  $j_1, \dots, j_{n+1}$

$$\begin{aligned} P(X_{t_{n+1}} = j_{n+1} \mid X_{t_n} = j_n, X_{t_{n-1}} = j_{n-1}, \dots, X_{t_1} = j_1) \\ = P(X_{t_{n+1}} = j_{n+1} \mid X_{t_n} = j_n). \end{aligned}$$

Thus the “future” is conditionally independent of the “past” given the present.

Time-homogeneous Markov process:

$$p_{ij}(t) = P(X_t = j \mid X_0 = i) = P(X_{t+s} = j \mid X_s = i),$$

for all  $t, s \geq 0$ .

Observe:  $p_{ii}(0) = 1$  and  $p_{ij}(0) = 0$  for all  $i \neq j$ , so if

$$\mathbf{P}(t) = \begin{pmatrix} \ddots & \vdots & \cdot \\ \dots & p_{ij}(t) & \dots \\ \cdot & \vdots & \ddots \end{pmatrix},$$

then  $\mathbf{P}(0) = I$  (the identity matrix). The matrices  $\mathbf{P}(t)$  are called the **matrices of transition probabilities**. These matrices are stochastic matrices i.e. they have non-negative entries and row-sums = 1.

**Example:** Poisson process  $(N_t)_{t \geq 0}$ , with intensity parameter  $\lambda > 0$ : Stochastic process on  $S = \{0, 1, 2, \dots\}$  starting at  $N_0 = 0$  with

$$P(N_{t_{n+1}} = j_{n+1} \mid N_{t_n} = j_n, N_{t_{n-1}} = j_{n-1}, \dots, N_{t_1} = j_1)$$

$\underbrace{\quad}_{\text{independent increments}} \stackrel{:=}{=} P(N_{t_{n+1}} - N_{t_n} = j_{n+1} - j_n)$

$$\underbrace{N_t - N_s \sim Po(\lambda(t-s))}_{\stackrel{:=}{=}}, 0 \leq s < t \quad e^{-\lambda(t_{n+1}-t_n)} \frac{(\lambda(t_{n+1}-t_n))^{j_{n+1}-j_n}}{(j_{n+1}-j_n)!},$$

if  $j_k \leq j_{k+1}$ , for all  $k = 1, 2, \dots, n$ .

( $P(N_{t_{n+1}} = j_{n+1} \mid N_{t_n} = j_n, N_{t_{n-1}} = j_{n-1}, \dots, N_{t_1} = j_1) = 0$ , otherwise.)

(With  $N_t - N_s \sim Po(\lambda(t-s))$  we thus mean that the increment  $N_t - N_s$  has the Poisson distribution with intensity parameter  $\lambda(t-s)$ .)

Thus  $(N_t)$  is a Markov process with

$$p_{ij}(t) = \begin{cases} e^{-\lambda t} \frac{(\lambda t)^{j-i}}{(j-i)!}, & j \geq i \\ 0, & j < i \end{cases}$$

Chapman-Kolmogorov equations:

$$\underbrace{P(X_{t+s} = k \mid X_0 = i)}_{p_{ik}(s+t)} = \sum_{j \in S} \underbrace{P(X_s = j \mid X_0 = i)}_{p_{ij}(s)} \underbrace{P(X_{t+s} = k \mid X_s = j, X_0 = i)}_{p_{jk}(t)}$$

If we express the Chapman-Kolmogorov equations in matrix form we get:

$$\underbrace{\begin{pmatrix} \ddots & \vdots & \cdot \\ \dots & p_{ik}(s+t) & \dots \\ \cdot & \vdots & \ddots \end{pmatrix}}_{\mathbf{P}(s+t)} = \underbrace{\begin{pmatrix} \ddots & \vdots & \cdot \\ p_{i1}(s) & p_{i2}(s) \dots & p_{i.}(s) \\ \cdot & \vdots & \ddots \end{pmatrix}}_{\mathbf{P}(s)} \underbrace{\begin{pmatrix} \ddots & p_{1k}(t) & \cdot \\ \dots & p_{2k}(t) & \dots \\ \cdot & p_{.k}(t) & \ddots \end{pmatrix}}_{\mathbf{P}(t)}$$

Note:

For Markov chains (with discrete time): If we know the transition matrix  $\mathbf{P}$ , then we can calculate  $\mathbf{P}^n$ , for all  $n$ .

For Markov processes (with continuous time): If we know the matrix  $\mathbf{P}(t)$ , for some  $t$ , then we cannot calculate  $\mathbf{P}(t)$  for all  $t$ . (only for  $\mathbf{P}(t), \mathbf{P}(2t), \mathbf{P}(3t), \dots$ )

If we let  $t \rightarrow 0$  and note that

$$\frac{p_{ij}(t) - p_{ij}(0)}{t} = \begin{cases} \frac{p_{ij}(t)}{t} & i \neq j \\ \frac{p_{ij}(t)-1}{t} & i = j, \end{cases}$$

we get

$$p_{ij}(t) \approx \begin{cases} tp'_{ij}(0) & i \neq j \\ 1 + tp'_{ij}(0) & i = j, \end{cases}$$

for small  $t$ . (provided these derivatives exist)

It therefore seems plausible that the matrix of derivatives  $(p'_{ij}(t))$  at time  $t = 0$  for a Markov process would play a similar role as the transition matrix  $\mathbf{P}$  for a Markov chain in characterizing the process.

**Remark:** It can be proved that the derivatives  $(p'_{ij}(t))$  exist for Markov processes with right-continuous trajectories satisfying the property that only a finite number of jumps can be made in each finite time-interval.

**Definition:** Let  $(X_t)$  be a (discrete, time-homogeneous) Markov process. Suppose there exists  $q_{ij} \geq 0$  for  $j \neq i$  and  $q_{ii} \leq 0$  such that

$$P(X_t = j \mid X_0 = i) = q_{ij}t + o(t),$$

and

$$P(X_t \neq i \mid X_0 = i) = q_i t + o(t),$$

where  $q_i = \sum_{j \neq i} q_{ij} =: -q_{ii}$ , and  $o(t)$ , small ordo of  $t$ , is a function satisfying  $\lim_{t \rightarrow 0} o(t)/t = 0$ . The numbers  $q_{ij}$  are called the **transition intensities** from state  $i$  to state  $j$  and the array

$$\mathbf{Q} = \left( \begin{array}{ccc} \ddots & \vdots & \cdot \\ \cdots & q_{ij} & \cdots \\ \cdot & \vdots & \ddots \end{array} \right)_{i,j \in S}$$

is called the **intensity matrix, or the infinitesimal generator** of the Markov process.

**Remark:** Note that

$$\sum_j q_{ij} = \sum_{j \neq i} q_{ij} + q_{ii} = -q_{ii} + q_{ii} = 0,$$

for any state  $i$ , so row sums in  $\mathbf{Q}$  are zero by definition.

**Remark:** Note that

$$P(X_t = i \mid X_0 = i) = 1 - q_i t + o(t),$$

for any state  $i$ , since  $P(X_t = i \mid X_0 = i) + P(X_t \neq i \mid X_0 = i) = 1$ .

**Remark:** Note that if  $S$  is finite, then  $\mathbf{P} := \frac{1}{\max q_i} \mathbf{Q} + I$  is a stochastic matrix. This makes it possible to translate some properties for discrete time Markov chains to the continuous time case, see e.g. Lawler, exercise 3.4.

**Theorem:** Let  $(X_t)_{t \geq 0}$  be a (discrete, time-homogeneous) Markov process with intensity matrix  $\mathbf{Q}$ . If  $T_1$  denotes the time for the first jump of the process, then:

- (i)  $P(T_1 \leq t \mid X_0 = i) = 1 - e^{-q_i t}$  i.e. if the process starts in state  $i$  then  $T_1$  is exponentially distributed with expectation  $1/q_i$ .
- (ii)  $P(X_{T_1} = j \mid X_0 = i) = \frac{q_{ij}}{q_i}$ , for  $j \neq i$  i.e. if the process starts in state  $i$  then the probability that the next state is  $j$  is  $q_{ij}/q_i$ .
- (iii)  $T_1$  and  $X_{T_1}$  are conditionally independent given that  $X_0 = i$ .

Sketch of proof: By the Markov property and homogeneity of the chain it follows that  $T_1$  has the “lack of memory” property (see the appendix below), and therefore  $T_1$  must be exponentially distributed. The remaining part of the proof follows from properties of exponentially distributed random variables stated in the appendix.

**Construction of a Markov process,  $(X_t)_{t \geq 0}$ , from an intensity matrix  $\mathbf{Q}$ :**

1. Choose a starting point  $Z_0$ .
2. If  $Z_0 = j$  then the process stays in state  $j$  a random time interval  $U_0 \sim \text{Exp}(q_j)$ . ( $U_0$  is called the first holding time.) (We will typically only consider cases when  $0 < q_j < \infty$ . If  $q_j = 0$  then the process will never leave state  $j$ , and if  $q_j = \infty$  then the process leaves  $j$  immediately.)
3. Next choose a new state,  $Z_1$ , according to the matrix  $R = (r_{ij})$ , with  $r_{ij} = q_{ij}/q_i$ ,  $i \neq j$ , and  $r_{ii} = 0$ .
4. Repeat inductively steps 2 and 3. If  $Z_j = k$  let  $U_j \sim \text{Exp}(q_k)$  (be holding times independent of  $U_0, \dots, U_{j-1}$ )

If  $J_n = \sum_{i=0}^{n-1} U_i$  and  $J_n \rightarrow \infty$  then we can define  $X_t$  for any  $t \geq 0$  by  $X_t = Z_m$  if  $J_m \leq t < J_{m+1}$ .




Figure 1: Trajectory of a Markov process with state space  $S = (1, 2, 3)$

**Theorem:** The process  $(X_t)$  defined as above is a (homogeneous) Markov process.

**Remark:** The Markov chain  $(Z_n)$  with transition matrix  $R$  is called the jump-chain corresponding to  $(X_t)$ . Note that the definition of  $R$  above presumes that  $q_i > 0$ . To avoid ambiguity we define  $r_{ij} = 0$ , for  $i \neq j$  and  $r_{ii} = 1$  in case  $q_i = 0$ .

**Example:** Construction of the Poisson process:

$q_j = \lambda$  for all  $j$  (for some parameter  $\lambda > 0$ )

$r_{i,i+1} = 1$  for any  $i \geq 0$ .

## 2 Appendix: Some properties of exponentially distributed random variables

The exponential distribution may be viewed as a continuous counterpart to the geometric distribution.

We say that a random variable  $X$  is exponentially distributed with parameter  $\lambda > 0$  (notation  $X \sim \text{Exp}(\lambda)$ ) if its density function is given by  $f_X(x) = \lambda e^{-\lambda x}$ ,  $x \geq 0$ ,  $f_X(x) = 0$ , otherwise.

Distribution function:  $F_X(x) = 1 - e^{-\lambda x}$ ,  $x \geq 0$ .

If  $X \sim \text{Exp}(\lambda)$  then  $E(X) = \int_0^\infty P(X > x) dx = \int_0^\infty e^{-\lambda x} = \frac{1}{\lambda}$ .

Exponentially distributed random variables have many interesting properties:

**Theorem:** (Lack of memory property.) Let  $X \sim \text{Exp}(\lambda)$ . Then

$$P(X > x + y \mid X > y) = P(X > x), \text{ for all } x, y > 0.$$

The exponential distribution is the only non-negative continuous distribution having the lack of memory property.

If  $X_1 \sim \text{Exp}(\lambda_1)$  and  $X_2 \sim \text{Exp}(\lambda_2)$  are independent then

$$\begin{aligned} P(X_1 < X_2) &= \int_{\mathbb{R}} P(X_1 < X_2 \mid X_1 = s) f_{X_1}(s) dt \\ &= \int_0^\infty \underbrace{P(X_2 > s)}_{e^{-\lambda_2 s}} \underbrace{f_{X_1}(s)}_{\lambda_1 e^{-\lambda_1 s}} ds \\ &= \lambda_1 \int_0^\infty e^{-(\lambda_1 + \lambda_2)s} ds = \frac{\lambda_1}{\lambda_1 + \lambda_2}, \end{aligned} \quad (1)$$

and if  $T = \min(X_1, X_2)$  then  $T \sim \text{Exp}(\lambda_1 + \lambda_2)$  since

$$\begin{aligned} P(T > t) &= P(X_1 > t, X_2 > t) \\ &\stackrel{\text{by indep.}}{=} P(X_1 > t)P(X_2 > t) = e^{-(\lambda_1 + \lambda_2)t}. \end{aligned}$$

The events  $\{\min(X_1, X_2) > t\}$  and  $\{X_1 < X_2\}$  are also independent, since a slight generalization of (1) gives  $P(t < X_1 < X_2) = \lambda_1 e^{-(\lambda_1 + \lambda_2)t} / (\lambda_1 + \lambda_2)$ .

More generally:

**Theorem:** Let  $X_1 \sim \text{Exp}(\lambda_1), X_2 \sim \text{Exp}(\lambda_2), \dots$  be independent exponentially distributed random variables with  $0 < \sum_i \lambda_i < \infty$ . Let  $X = \inf_k X_k$ .

Then this infimum is attained at a unique random value  $K$  of  $k$  (with probability 1). Moreover  $X$  and  $K$  are independent and

$$X \sim \text{Exp}\left(\sum_i \lambda_i\right),$$

and

$$P(K = k) = \frac{\lambda_k}{\sum_i \lambda_i}.$$

### 3 Matlab

#### 3.1 Program for simulating a Markov processes with finite state space and given generator $Q$ (Figure 1)

```

Q=[-12 10 2; 1 -2 1; 2 2 -4];           % Choose intensity matrix
Tmax=3;                                % The process will be simulated up to time Tmax
z(1) = 3;                               % Choose initial state in 1,2,...,m (in case Q has size m × m)
                                         % (We store  $Z_n$  in z(n+1) and  $J_n$  in J(n+1).)

for i=1:length(Q)                       % corresponding jump-chain
    for j=1:length(Q)
        if i==j
            R(i,j)=0;
        else
            R(i,j)=-Q(i,j)/Q(i,i);
        end
    end
end
J(1) = 0;                               % start time at t=0
i = 1;
while J(i) < Tmax,
    J(i+1) = J(i) + log(rand)/Q(z(i),z(i)); % Next jump-time
    u = rand;                             % Next state is chosen according to R
    j = 1;
    s = R(z(i),1);
    while ((u > s) & (j < length(Q) )),
        j=j+1;
        s=s+R(z(i),j);
    end
    z(i+1) = j;
    i=i+1;
end
cla;
hold on                                % A trajectory plot consists of many...
for i=1:(length(J)-1),
    plot([J(i) J(i+1)], [z(i) z(i)]);    % ... line-segments
end
axis([0 Tmax 0 (length(Q)+1)]);
xlabel('time');
ylabel('state');

```



## 4 Suggested exercises

Basic exercises:

18, 20

Exercises Lawler:

3.1, 3.3

## Lecture 11 Markov Processes, 1MS012

### 5 Forward & Backward equations

Recall:

A (well behaved) Markov process gives rise to an intensity matrix or generator  $\mathbf{Q}$ . Conversely given a matrix  $\mathbf{Q} = (q_{jk})$  where  $q_{jk}$ , for all  $j, k \in S$  satisfy

(a)  $q_{jk} \geq 0, j \neq k$ .

(b)  $-\infty < q_{jj} \leq 0$ .

(c)  $\sum_{k \in S} q_{jk} = 0, j \in S$ ,

we can construct a Markov process (at least if  $S$  is finite), with holding times, exponentially distributed with parameters  $-q_{jj} = q_j < \infty$ , and jump chain also specified by the entries in  $\mathbf{Q}$ .

The matrix  $\mathbf{Q}$  gives all information about the Markov process in the sense that for small time steps  $h$

$$p_{ij}(h) = P(X_{t+h} = j \mid X_t = i) = q_{ij}h + o(h), i \neq j$$

$$p_{jj}(h) = P(X_{t+h} = j \mid X_t = j) = 1 + q_{jj}h + o(h) = 1 - q_jh + o(h)$$

Markov processes in discrete time:

$P(X_n = k)$  is "easy" to find if we know the distribution of  $X_0$  and the transition matrix  $\mathbf{P}$ .

Markov processes continuous time:

Not obvious how to use  $\mathbf{Q}$  to find  $P(X_t = k)$ . We can derive equations for  $p_{ij}(t) = P(X_t = j \mid X_0 = i)$  that hopefully can be solved:

Forward equations: ( $S$  finite)

We study the derivatives of  $p_{ij}(t)$ .

$$\begin{aligned} \lim_{h \rightarrow 0} \frac{p_{ij}(t+h) - p_{ij}(t)}{h} & \stackrel{\text{CK-equations}}{=} \lim_{h \rightarrow 0} \frac{\sum_{k \in S} p_{ik}(t) p_{kj}(h) - p_{ij}(t)}{h} \\ & \stackrel{S \text{ finite}}{=} \sum_{k \in S \setminus \{j\}} p_{ik}(t) \underbrace{\lim_{h \rightarrow 0} \frac{p_{kj}(h)}{h}}_{q_{kj}} + p_{ij}(t) \underbrace{\lim_{h \rightarrow 0} \frac{p_{jj}(h) - 1}{h}}_{q_{jj}} \\ & = \sum_{k \in S} p_{ik}(t) q_{kj}. \end{aligned}$$

The equations

$$p'_{ij}(t) = \sum_{k \in S} p_{ik}(t) q_{kj}$$

are called **Kolmogorov's forward equations**.

Forward equations expressed in matrix form:  $\mathbf{P}'(t) = \mathbf{P}(t)\mathbf{Q}$

$$\underbrace{\begin{pmatrix} \ddots & \vdots & \cdot \\ \cdots & p'_{ij}(t) & \cdots \\ \cdot & \vdots & \ddots \end{pmatrix}}_{\mathbf{P}'(t)} = \underbrace{\begin{pmatrix} \ddots & \vdots & \cdot \\ p_{i1}(t) & p_{i2}(t) \cdots & p_{i|S|(t)} \\ \cdot & \vdots & \ddots \end{pmatrix}}_{\mathbf{P}(t)} \underbrace{\begin{pmatrix} \ddots & q_{1j} & \cdot \\ \cdots & q_{2j} & \cdots \\ \cdot & q_{|S|j} & \ddots \end{pmatrix}}_{\mathbf{Q}}$$

(Actually we just proved that the right-derivative exists but a similar argument shows that also the left derivative exists.)

**Example:** A machine works an exponential time with mean  $\frac{1}{\mu}$ . It takes an exponential time with mean  $\frac{1}{\lambda}$  to repair it. If the machine works at time 0, what is the probability that it will not work at time  $t$ ?

We model the state of the machine with a Markov process. Let  $(X_t)$  be a Markov process with state space  $S = (0, 1)$  where  $X_t$  represents the state of the process at time  $t$ : Let

$X_t = 1$  mean that the machine works at time  $t$ , and

$X_t = 0$  mean that it doesn't work at time  $t$ .

When  $X_t$  enters state 1 it stays there an  $\text{Exp}(\mu)$ -distributed time and then goes to 0 and stays there an  $\text{Exp}(\lambda)$ -distributed time and then goes to state 1 and so on.

We have  $q_0 = \lambda = -q_{00}$ , and  $q_1 = \mu = -q_{11}$ . Thus we get the intensity matrix

$$\mathbf{Q} = \begin{pmatrix} -\lambda & \lambda \\ \mu & -\mu \end{pmatrix},$$

(since row sums of  $\mathbf{Q}$  are zero).

We want to find  $p_{10}(t) = P(X_t = 0 | X_0 = 1)$ .

The forward equations gives:

$$\begin{aligned} p'_{10}(t) &= -\lambda p_{10}(t) + \mu p_{11}(t) = -\lambda p_{10}(t) + \mu(1 - p_{10}(t)) \\ &= \mu - (\lambda + \mu)p_{10}(t), \end{aligned}$$

with the boundary condition  $p_{10}(0) = 0$ .

The equation

$$p'_{10}(t) + (\lambda + \mu)p_{10}(t) = \mu,$$




Figure 2:  $(X_t)$  with parameters  $\lambda = 1$ , and  $\mu = 2$ .

$$\lim_{t \rightarrow \infty} p_{10}(t) = \mu/(\lambda + \mu) = 2/3.$$

can be solved by the standard trick of equivalently writing it as

$$\underbrace{p'_{10}(t)e^{(\lambda+\mu)t} + (\lambda + \mu)e^{(\lambda+\mu)t}p_{10}(t)}_{\frac{d}{dt}(p_{10}(t)e^{(\lambda+\mu)t})} = \mu e^{(\lambda+\mu)t}.$$

By integrating both sides we get

$$p_{10}(t)e^{(\lambda+\mu)t} = \frac{\mu}{\lambda + \mu}e^{(\lambda+\mu)t} + c,$$

where  $c$  is a constant. The boundary condition  $p_{10}(0) = 0 \Rightarrow c = \frac{-\mu}{\lambda + \mu}$ .  
Thus

$$p_{10}(t) = \frac{\mu}{\lambda + \mu}(1 - e^{-(\lambda+\mu)t}).$$

## 5.1 Kolmogorov's backward equations

If  $S$  is infinite then the Kolmogorov forward equations may fail to hold.

To avoid such problems we derive another set of equations.

By conditioning on the state of the process at time  $h$  we get

$$\begin{aligned} \lim_{h \rightarrow 0} \frac{p_{ij}(t+h) - p_{ij}(t)}{h} &\stackrel{\text{CK-equations}}{=} \lim_{h \rightarrow 0} \frac{\sum_{k \in S} p_{ik}(h)p_{kj}(t) - p_{ij}(t)}{h} \\ &\stackrel{\text{Can be justified}}{=} \sum_{k \in S \setminus \{i\}} \underbrace{\lim_{h \rightarrow 0} \frac{p_{ik}(h)}{h}}_{q_{ik}} p_{kj}(t) + p_{ij}(t) \underbrace{\lim_{h \rightarrow 0} \frac{p_{ii}(h) - 1}{h}}_{q_{ii}} \\ &= \sum_{k \in S} q_{ik} p_{kj}(t). \end{aligned}$$

We have thus derived **Kolmogorov's backward equations**

$$p'_{jk}(t) = \sum_{r \in S} q_{jr} p_{rk}(t)$$

expressed in matrix form as:

$$\mathbf{P}'(t) = \mathbf{Q}\mathbf{P}(t).$$

Thus if the forward equation also holds then

$$\mathbf{P}'(t) = \mathbf{Q}\mathbf{P}(t) = \mathbf{P}(t)\mathbf{Q}.$$

**Theorem:** If  $(X_t)_{t \geq 0}$  is a Markov process with finite state space, then

$$\mathbf{P}(t) = e^{t\mathbf{Q}} := \sum_{k=0}^{\infty} \mathbf{Q}^k t^k / k! = \mathbf{I} + t\mathbf{Q} + \frac{t^2 \mathbf{Q}^2}{2!} + \frac{t^3 \mathbf{Q}^3}{3!} + \dots,$$

is the unique solution of the forward and backward equations, with initial condition  $\mathbf{P}(0) = \mathbf{I}$  where  $\mathbf{I}$  is the identity matrix.

This explicitly explains why  $\mathbf{Q}$  is called the “generator” of the process.

If  $S$  is finite and  $\mathbf{Q}$  can be diagonalized, with  $\mathcal{P}^{-1}\mathbf{Q}\mathcal{P} = \mathbf{D}$ , for some invertible matrix  $\mathcal{P}$ , where  $\mathbf{D}$  is a diagonal matrix with diagonal entries  $d_i$ , then

$$\mathbf{P}(t) = \mathcal{P}e^{t\mathbf{D}}\mathcal{P}^{-1},$$

where  $e^{t\mathbf{D}}$  is the diagonal matrix with diagonal entries  $e^{d_i t}$ .

If  $S$  infinite then it is possible to prove that if there exists a solution,  $\mathbf{P}(t)$  to the forward and backward equations with  $\sum_{j \in S} p_{ij}(t) = 1$ , for all  $t$  and  $i \in S$ , then this is the unique solution.

In general it is hard to explicitly solve the forward and backward equations if  $S$  has many states and  $\mathbf{Q}$  has no particular structure. Typically the forward and backward equations have to be solved by numerical methods.

## 6 Matlab

### 6.1 Program for Figure 2

```

lambda=1;      % parameter values
mu=2;          %
x0=0;          % starting point
[t,y]=ode45(@hogerled,[0,5],x0,[],lambda,mu);
               % Numerically solves ODEs of the form  $y'(t)=f(t,y)$ 
               % with given starting values, on a specified interval
               % with specified parameters.
               % The function  $f(t,y)$  is specified in the m-file
               % hogerled.m below.

plot(t,y)
legend('p_{10}(t)')

% In the file hogerled.m
function fty=hogerled(t,y,lambda,mu);
fty=mu-(lambda+mu)*y;

```

## 7 Suggested exercises

Basic exercises:

19

Extra problems:

c2

Exercises Lawler:

3.5

## Lecture 12 Markov Processes, 1MS012

### 8 Class structure

**Definition:** State  $k$  is **accessible** from state  $j$  ( $j \rightarrow k$ ) if  $p_{jk}(t) > 0$  for some  $t \geq 0$ .

$k$  and  $j$  are intercommunicating ( $j \leftrightarrow k$ ) if  $j \rightarrow k$  and  $k \rightarrow j$

**Theorem:** Let  $(X_t)$  be a discrete Markov process with intensity matrix  $\mathbf{Q}$ . Let  $j$  and  $k$  be two (distinct) states. Then

$$j \rightarrow k$$

$$\Leftrightarrow$$

there exists an admissible path from  $j$  to  $k$  with respect to the matrix  $\mathbf{Q}$  i.e. if either  $q_{jk} > 0$  or there exists states  $j_1, \dots, j_n$  such that  $q_{jj_1} > 0, q_{j_1j_2} > 0, \dots, q_{j_nk} > 0$ .

$$\Leftrightarrow$$

$$p_{jk}(t) > 0, \text{ for all } t > 0,$$

The last equivalence can be seen since holding times of arbitrary small length have positive probability.

Conclusion:

Don't need to bother about aperiodicity for Markov processes in continuous time.

Closed sets and irreducible sets/processes are defined in analogy with the discrete time case.

The definition of recurrent and transient states has to be presented with care since all states are re-visited an infinite amount of time during the first holding time;

**Definition:** We say that  $j$  is recurrent if  $j$  is recurrent for the jump-chain,

otherwise  $j$  is transient.

A state  $j$  is recurrent if, starting at  $j$  the process visits  $j$  at arbitrary large time-points (with probability 1). A state  $j$  is transient otherwise.

Let  $T_{jj}$  be the first time the process returns to  $j$  after the first jump. Thus if  $X_0 = j$  then  $T_{jj} = \inf(t > J_1 : X_t = j)$ , where  $J_1$  is the time for the first jump. If  $j$  is a recurrent state, and  $ET_{jj} = \infty$ , then  $j$  is said to be null-recurrent. Otherwise we call  $j$  non-null, (or positive) recurrent.

If  $q_j = 0$  then we may define  $j$  to be non-null (positive) recurrent.

**Theorem:**  $j$  is recurrent iff  $\int_0^\infty p_{jj}(t)dt = \infty$ .

Similarly as for Markov chains in discrete time it can also be proved that

- if  $i$  and  $j$  are two states with  $i \leftrightarrow j$  then  $i$  is positive recurrent iff  $j$  is positive recurrent.
- All states are positive recurrent for a finite irreducible Markov process.

## 8.1 Stationary distributions and the long run

Suppose  $S = \{1, 2, \dots, n\}$  is finite and

$$p_{jk}(t) \rightarrow \pi_k \quad \text{as } t \rightarrow \infty, \quad (2)$$

for all  $j$  and  $k$ , i.e.

$$\mathbf{P}(t) = \begin{pmatrix} p_{11}(t) & \cdot & \cdot & p_{1n}(t) \\ p_{21}(t) & \cdot & \cdot & p_{2n}(t) \\ \cdot & \cdot & \cdot & \cdot \\ p_{n1}(t) & \cdot & \cdot & p_{nn}(t) \end{pmatrix} \rightarrow \begin{pmatrix} \pi_1 & \pi_2 & \cdot & \pi_n \\ \pi_1 & \pi_2 & \cdot & \pi_n \\ \pi_1 & \pi_2 & \cdot & \pi_n \\ \pi_1 & \pi_2 & \cdot & \pi_n \end{pmatrix}$$

Then,  $\pi = (\pi_1, \pi_2, \dots, \pi_n)$ , has the following properties:

1. If  $\boldsymbol{\mu}(t) = (\mu_1(t), \dots, \mu_n(t)) := (P(X_t = 1), \dots, P(X_t = n))$  is the  $1 \times n$  vector representing the probability distribution of  $X_t$  then

$$\boldsymbol{\mu}(t) = \boldsymbol{\mu}(0)\mathbf{P}(t) \rightarrow \pi,$$

as  $t \rightarrow \infty$ . This can be seen from (2) since

$$\begin{aligned} \mu_k(t) &= P(X_t = k) = \sum_{j \in S} P(X_t = k | X_0 = j)P(X_0 = j) = \sum_{j \in S} p_{jk}(t)\mu_j(0) \\ &\rightarrow \sum_{j \in S} \pi_k \mu_j(0) = \pi_k, \quad \text{as } t \rightarrow \infty. \end{aligned}$$



2.

$$\pi = \pi \mathbf{P}(s),$$

for all  $s$ , i.e.  $\pi$  is a stationary (or equilibrium) distribution.

This can be seen by letting  $t \rightarrow \infty$  in the Chapman-Kolmogorov identity since

$$\underbrace{\boldsymbol{\mu}(0)\mathbf{P}(t+s)}_{\rightarrow \pi} = \underbrace{\boldsymbol{\mu}(0)\mathbf{P}(t)}_{\rightarrow \pi} \mathbf{P}(s)$$

(Thus if  $X_0 \stackrel{d}{\sim} \pi$  then  $(X_t)$  forms a stationary process.)

3.

$$\pi \mathbf{Q} = \mathbf{0}$$

This can be seen since  $\lim_{t \rightarrow \infty} p_{jk}(t) = \pi_k \Rightarrow \lim_{t \rightarrow \infty} p'_{jk}(t) = 0$  and since by the forward equations,  $\mathbf{P}'(t) = \mathbf{P}(t)\mathbf{Q}$ ,

$$\underbrace{\pi \mathbf{P}'(t)}_{\rightarrow \mathbf{0}} = \underbrace{\pi \mathbf{P}(t)}_{\pi} \mathbf{Q}.$$

4. The distribution

$$\nu = \frac{1}{\sum \pi_k q_k} (\pi_1 q_1, \dots, \pi_n q_n),$$

is a stationary distribution for the jump chain, i.e.  $\nu \mathbf{R} = \nu$ , where

$$\mathbf{R} = \begin{pmatrix} r_{11} & \cdot & \cdot & r_{1n} \\ r_{21} & \cdot & \cdot & r_{2n} \\ \cdot & \cdot & \cdot & \cdot \\ r_{n1} & \cdot & \cdot & r_{nn} \end{pmatrix}$$

is defined by  $r_{ij} = q_{ij}/q_i$  if  $i \neq j$ , and  $r_{ii} = 0$  for any  $i$ . (Recall the implicit assumption that  $q_i > 0$  for all  $i$ , so  $r_{ij}$  is well defined.)

Check:

Let  $\nu_j = \frac{\pi_j q_j}{\sum \pi_k q_k}$ . The distribution  $\nu = (\nu_1, \dots, \nu_n)$  is stationary, since

$$\nu_j = \frac{\pi_j q_j}{\sum \pi_k q_k} = \frac{-\pi_j q_{jj}}{\sum \pi_k q_k} = \frac{\sum_{i \neq j} \pi_i q_{ij}}{\sum \pi_k q_k} = \sum_{i \neq j} \frac{\pi_i q_i}{(\sum \pi_k q_k)} \frac{q_{ij}}{q_i} = \sum_i \nu_i r_{ij},$$

for each  $j$ , where in the third equality we used that  $\pi \mathbf{Q} = \mathbf{0}$ .

5. The stationary distribution  $\pi$  satisfies

$$\pi_k = \frac{1}{q_k E(T_{kk})},$$

i.e.  $\pi_k$  is the proportion of time spent in state  $k$ .

The proof of this uses the renewal theorem.

(Interpretation: On average it takes a time of length  $E(T_{kk})$  between arrivals to the state  $k$ . During this time we spend on average a time of length  $1/q_k$  in the state  $k$  and the rest of the time in other states.)

**Theorem:** If  $(X_t)$  is an irreducible (discrete, time-homogeneous) Markov process such that all states are positive recurrent then

$$p_{jk}(t) \rightarrow \pi_k = \frac{1}{q_k E(T_{kk})},$$

as  $t \rightarrow \infty$  for all  $j$  and  $k$ , and  $\pi = (\pi_j)_{j \in S}$ , is uniquely stationary for the process, and the unique solution to the equation  $\pi \mathbf{Q} = \mathbf{0}$ , with  $\sum_{i \in S} \pi_i = 1$ , and  $\pi_i \geq 0$ , for any  $i$ .

How can we check if an irreducible Markov process is positive recurrent?

**Theorem:** Let  $(X_t)$  be an irreducible Markov process with intensity matrix  $\mathbf{Q}$  such that there is a unique solution to the forward and backward equations. Then

$(X_t)$  is positive recurrent

$\Leftrightarrow$

there exists a solution to  $\pi \mathbf{Q} = \mathbf{0}$ , with  $\sum_{i \in S} \pi_i = 1$ , and  $\pi_i \geq 0$ , for any  $i$ .

**Remark:** If  $S$  is finite then  $P(t) = e^{\mathbf{Q}t}$  is the unique solution to the forward and backward equations.

**Example:** Let  $(X_t)_{t \geq 0}$  be a Markov process on  $S = (0, 1, 2)$  with intensity matrix

$$\mathbf{Q} = \begin{pmatrix} q_{00} & q_{01} & q_{02} \\ q_{10} & q_{11} & q_{12} \\ q_{20} & q_{21} & q_{22} \end{pmatrix} = \begin{pmatrix} -1 & 1 & 0 \\ 2 & -3 & 1 \\ 0 & 2 & -2 \end{pmatrix}.$$

Suppose we wish to find the limit

$$\lim_{t \rightarrow \infty} P(X_t = 2 \mid X_0 = 0).$$

Since the Markov process  $(X_t)$  is irreducible with a finite state space it follows from the convergence theorem that

$$\lim_{t \rightarrow \infty} P(X_t = 2 \mid X_0 = 0) = \pi_2,$$

where  $\pi = (\pi_0, \pi_1, \pi_2)$  is the unique probability vector solving the equation

$$\pi \mathbf{Q} = \mathbf{0}.$$

This system has solution  $\pi = (4/7, 2/7, 1/7)$ .

Thus

$$\lim_{t \rightarrow \infty} P(X_t = 2 \mid X_0 = 0) = 1/7.$$

**Example:** (continued)

Consider the same Markov process as in the previous example.

Suppose, starting at state 0, we wish to find the expected time it takes to reach state 2.

Let  $T_{i2} = \inf(t > J_1 : X_t = 2 \mid X_0 = i)$  be the first passage time from state  $i$  to state 2,  $i = 0, 1, 2$ .

In order to find  $E(T_{02})$  we will use the idea of “first step analysis” but since time is continuous the first “interesting time-step” is naturally the time when the first jump occurs. More precisely

$$E(T_{02}) = \sum_{k=0}^2 E(T_{02} \mid X_{J_1} = k, X_0 = 0) \underbrace{P(X_{J_1} = k \mid X_0 = 0)}_{r_{0k}},$$

where  $r_{ik} = -q_{ik}/q_{ii}$ ,  $i \neq j$  are the transition probabilities of the jump chain, i.e.

$$\mathbf{R} = \begin{pmatrix} r_{00} & r_{01} & r_{02} \\ r_{10} & r_{11} & r_{12} \\ r_{20} & r_{21} & r_{22} \end{pmatrix} = \begin{pmatrix} 0 & 1 & 0 \\ 2/3 & 0 & 1/3 \\ 0 & 1 & 0 \end{pmatrix}.$$

Since  $T_{02}$  is the time it takes to leave state 0 plus the time it takes to reach state 2 from the new state reached after the first jump from 0, it follows from the Markov property that

$$E(T_{02} \mid X_{J_1} = k, X_0 = 0) = E(J_1 \mid X_0 = 0) + E(T_{k2}), \quad k = 0, 1,$$

$$E(T_{02} \mid X_{J_1} = 2, X_0 = 0) = E(J_1 \mid X_0 = 0).$$

From the intensity matrix we see that  $J_1 \mid (X_0 = 0) \sim \text{Exp}(q_0)$  and  $J_1 \mid (X_0 = 1) \sim \text{Exp}(q_1)$ , where  $q_0 = -q_{00} = 1$ , and  $q_1 = -q_{11} = 3$ . Thus  $E(J_1 \mid X_0 = 0) = 1$  and  $E(J_1 \mid X_0 = 1) = 1/3$ .

Therefore

$$E(T_{02}) = \underbrace{E(J_1 \mid X_0 = 0)}_{=1} + E(T_{12})r_{01} = 1 + E(T_{12}).$$

Similarly

$$E(T_{12}) = (1/q_1) + r_{10}E(T_{02}) = 1/3 + (2/3) \cdot E(T_{02}).$$

Thus  $E(T_{02}) = 1 + E(T_{12}) = 1 + 1/3 + (2/3)E(T_{02})$ , i.e.  $E(T_{02}) = 4$ .

## 9 Matlab

### 9.1 Code for solving $\pi Q = 0 \Leftrightarrow Q' \pi' = 0'$ for probability vectors $\pi$ .

```
Q=[-1 1 0; 2 -3 1; 0 2 -2]
```

```
pi= null(transpose(Q))/ sum(null(transpose(Q)))
```

## 10 Suggested exercises

Basic exercises:  
21–22, 27

Extra problems:  
b3, B3, B4

Exercises Lawler:  
3.8

## Lecture 13 Markov Processes, 1MS012

### 11 Birth-processes

**Definition:** A birth-process on  $S = (0, 1, 2, \dots)$  is a Markov process with generator

$$\mathbf{Q} = \begin{pmatrix} -\lambda_0 & \lambda_0 & 0 & 0 & \cdot & \cdot \\ 0 & -\lambda_1 & \lambda_1 & 0 & \cdot & \cdot \\ 0 & 0 & -\lambda_2 & \lambda_2 & 0 & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \end{pmatrix}.$$

The  $\lambda_j$ 's are called birthrates.

A birth-process is thus a Markovprocess with (trivial) jump chain  $r_{j,j+1} = 1$  and holding times  $\lambda_j$  for all states  $j$ .

Birth-processes can be used as models for the size of a growing population, where new individuals are born with intensity  $\lambda_i$  if the size of the population is  $i$ . The Poisson process corresponds to the special case when  $\lambda_i = \lambda$  i.e. the case when the birthrates do not depend on the population size.

#### 11.1 Transition probabilities for a birth-process

Forward equations  $\mathbf{P}'(t) = \mathbf{P}(t)\mathbf{Q}$ :

$$p'_{ij}(t) = \lambda_{j-1}p_{i,j-1}(t) - \lambda_j p_{i,j}(t), \quad j \geq 1.$$

Since  $p_{ij}(t) = 0$ , if  $j < i$ , we get

(I):

$$p'_{ii}(t) = -\lambda_i p_{ii}(t)$$

(II):

$$p'_{ij}(t) = \lambda_{j-1}p_{i,j-1}(t) - \lambda_j p_{i,j}(t), \quad j > i$$

Since  $p_{ii}(0) = 1$ , (I) has solution  $p_{ii}(t) = e^{-\lambda_i t}$ , for all  $i \geq 0$ .

We now solve (II):

$$\underbrace{e^{\lambda_j t}(p'_{ij}(t) + \lambda_j p_{i,j}(t))}_{\frac{d}{dt}(e^{\lambda_j t} p_{i,j}(t))} = (\lambda_{j-1} p_{i,j-1}(t)) e^{\lambda_j t}$$

By integrating both sides and using  $p_{ij}(0) = 0$ , for  $j > i$ , we finally get the recursive equations

$$p_{ij}(t) = e^{-\lambda_j t} \lambda_{j-1} \int_0^t p_{i,j-1}(s) e^{\lambda_j s} ds \quad (3)$$

for  $j > i$ .

**Remark:** It is an exercise to explicitly solve the recursive equations (3) in the special case  $\lambda_i = \lambda$  (corresponding to the Poisson process), to obtain the solution

$$p_{ij}(t) = e^{-\lambda t} (\lambda t)^{j-i} / (j-i)!, \quad j \geq i,$$

in that case.

## 11.2 Honest and dishonest processes

Let  $J_n$  be the time for the  $n$ :th jump. Let  $J_\infty = \lim_{n \rightarrow \infty} J_n$ .

If  $J_\infty < \infty$  then the process makes an infinite number of jumps just before time  $J_\infty$ . The process is then explosive and  $\mathbf{Q}$  does not define the process for all times in this case. This occurs if  $\lambda_i$  grows sufficiently fast.

Since  $U_i := (J_{i+1} - J_i) \sim \text{Exp}(\lambda_i)$  and thus

$$E J_\infty = E \left( \sum_{i=0}^{\infty} U_i \right) = \sum_{i=0}^{\infty} E(U_i) = \sum_{i=0}^{\infty} 1/\lambda_i,$$

we have:

**Theorem:** (Feller-Lundberg)

$$\sum_{i=0}^{\infty} 1/\lambda_i < \infty \Rightarrow P(J_\infty < \infty) = 1 \quad (\text{explosion always})$$

It is also possible to prove

$$\sum_{i=0}^{\infty} 1/\lambda_i = \infty \Rightarrow P(J_\infty = \infty) = 1 \quad (\text{no explosion})$$

**Example:** Poisson process  $\lambda_i = \lambda$ , does not explode since  $\sum_{i=0}^{\infty} 1/\lambda = \infty$ .

**Definition:** A Markov process  $(X_t)_{t \geq 0}$  is said to be

honest if  $P(J_\infty = \infty) = 1$   
dishonest if  $P(J_\infty = \infty) < 1$

We thus have a simple criteria for telling if a birth-process is honest or not.  
For more general Markov processes it is much harder:

**Theorem:** The Markov process  $(X_t)_{t \geq 0}$  with generator  $\mathbf{Q}$  is honest if one of the following conditions hold.

- $\sup_j q_j < \infty$
- The state space  $S$  is finite
- The process starts in a recurrent state.

Note that these are only sufficient conditions for a Markov process to be honest.

**Theorem:** If  $(X_t)_{t \geq 0}$  is irreducible with generator  $\mathbf{Q}$  then:

all states are positive recurrent  $\Leftrightarrow (X_t)_{t \geq 0}$  is honest and  $\pi \mathbf{Q} = \mathbf{0}$   
for some probability distribution  $\pi$ .

Obs:

Birth-processes are not irreducible, and stationary distributions do not exist for such processes.

## 12 Birth-death processes

**Definition:** A birth death process is a Markov process on  $S = (0, 1, 2, \dots)$  with generator

$$\mathbf{Q} = \begin{pmatrix} -\lambda_0 & \lambda_0 & 0 & 0 & \cdot & \cdot \\ \mu_1 & -(\lambda_1 + \mu_1) & \lambda_1 & 0 & \cdot & \cdot \\ 0 & \mu_2 & -(\lambda_2 + \mu_2) & \lambda_2 & 0 & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \end{pmatrix}$$

Thus  $i \mapsto i + 1$  (birth) and  $i \mapsto i - 1$  (death) are the only possible transitions.

$\lambda_0, \lambda_1, \dots$  birthrates/arrival rates  
 $\mu_1, \mu_2, \dots$  deathrates/departure rates

(the latter notation if regarded as the number of people in a system)

The process will be irreducible if  $\lambda_n > 0$ , and  $\mu_n > 0$  for all  $n$ .

In order to find stationary distributions we solve  $\pi \mathbf{Q} = \mathbf{0}$ .

$$-\lambda_0 \pi_0 + \mu_1 \pi_1 = 0 \quad (\text{I})$$

$$\lambda_{n-1} \pi_{n-1} - (\lambda_n + \mu_n) \pi_n + \mu_{n+1} \pi_{n+1} = 0, \quad n \geq 1 \quad (\text{II})$$

Thus

$$\mu_{n+1} \pi_{n+1} - \lambda_n \pi_n \underbrace{=}_{(\text{II})} \mu_n \pi_n - \lambda_{n-1} \pi_{n-1} \underbrace{=}_{(\text{II})} \dots = \mu_1 \pi_1 - \lambda_0 \pi_0 \underbrace{=}_{(\text{I})} 0, \quad n \geq 0,$$

i.e.

$$\mu_{n+1} \pi_{n+1} = \lambda_n \pi_n, \quad n \geq 0, \quad (\text{detailed balance equations})$$

so

$$\pi_n = \frac{\lambda_{n-1}}{\mu_n} \pi_{n-1} = \dots = \frac{\lambda_{n-1} \lambda_{n-2} \dots \lambda_0}{\mu_n \mu_{n-1} \dots \mu_1} \pi_0, \quad n \geq 1.$$

If  $\pi$  is a probability distribution then

$$\sum_{n=0}^{\infty} \pi_n = 1 \Leftrightarrow \left(1 + \sum_{n=1}^{\infty} \frac{\lambda_{n-1} \lambda_{n-2} \dots \lambda_0}{\mu_n \mu_{n-1} \dots \mu_1}\right) \pi_0 = 1$$

This partly proves

**Theorem:**

Let  $(X_t)_{t \geq 0}$  be a birth-death process with birth intensities  $\lambda_i > 0, i \geq 0$ , and death intensities  $\mu_i > 0, i \geq 1$ .

If

$$\sum_{k=1}^{\infty} \frac{\lambda_0 \lambda_1 \dots \lambda_{k-1}}{\mu_1 \mu_2 \dots \mu_k} < \infty$$

and

$$\sum_{k=1}^{\infty} \frac{\mu_1 \mu_2 \dots \mu_k}{\lambda_1 \lambda_2 \dots \lambda_k} = \infty \quad (4)$$

then a unique stationary distribution  $\pi$ , exists and

$$p_{ij}(t) = P(X_t = j | X_0 = i) \rightarrow \pi_j,$$



as  $t \rightarrow \infty$ , where

$$\pi_k = \frac{\lambda_0 \lambda_1 \cdots \lambda_{k-1}}{\mu_1 \mu_2 \cdots \mu_k} \pi_0, \quad \text{and} \quad \pi_0 = \left(1 + \sum_{k=1}^{\infty} \frac{\lambda_0 \lambda_1 \cdots \lambda_{k-1}}{\mu_1 \mu_2 \cdots \mu_k}\right)^{-1}.$$

**Remark:** It can be proved that (4) is a condition for recurrence of the state 0.

**Remark:** If  $\lambda_{n_0} = 0$ , for some  $n_0$ , then we regard  $S = (0, \dots, n_0)$  as the state space and if the process is irreducible (regarded as a process on  $S$ ), then

$$p_{ij}(t) = P(X_t = j | X_0 = i) \rightarrow \pi_j,$$

as  $t \rightarrow \infty$ , where

$$\pi_k = \frac{\lambda_0 \lambda_1 \cdots \lambda_{k-1}}{\mu_1 \mu_2 \cdots \mu_k} \pi_0, \quad \text{and} \quad \pi_0 = \left(1 + \sum_{k=1}^{n_0} \frac{\lambda_0 \lambda_1 \cdots \lambda_{k-1}}{\mu_1 \mu_2 \cdots \mu_k}\right)^{-1}.$$

## 12.1 Population models

Let  $X_t$  = number of individuals in a population at time  $t$ .

Suppose  $X_t = i$ , at some fixed time  $t$ . Suppose  $B \sim \text{Exp}(\lambda_i)$  is the time until next birth, and  $D \sim \text{Exp}(\mu_i)$  is the time until the next death, and  $B$  and  $D$  are independent. If  $D < B$  then the next jump will be  $i \mapsto i - 1$ . By properties of the exponential distribution it follows that  $P(D < B) = \frac{\mu_i}{\lambda_i + \mu_i}$ . If  $J_1$  is the time for the next jump, then  $J_1 = \min(B, D) \sim \text{Exp}(\lambda_i + \mu_i)$ .

It follows from the memoryless property of the exponential distribution that conditional on  $X_{J_1} = j$ , the process starts afresh from  $j$  at time  $J_1$ . Thus  $(X_t)_{t \geq 0}$  is a birth-death process with birthrates  $\lambda_i$ , and deathrates  $\mu_i$ .

**Example:** Each individual produces new individuals with rate  $\lambda$  and dies with rate  $\mu$ .

This is a birth-death process with  $\lambda_n = n\lambda$ ,  $\mu_n = n\mu$ .

## 12.2 Some common queueing models

$X_t$  = number of people on line for service.

Independent interarrival times  $\sim \text{Exp}(\lambda)$ ,  $\lambda > 0$ , and service times  $\sim \text{Exp}(\mu)$ ,  $\mu > 0$ .

**Example:**  $M/M/1$  queue

One server.

This is a birth-death process with  $\lambda_n = \lambda$ ,  $\mu_n = \mu$ .

(The two  $M$ :s refer to the fact that both inter-arrival and service times are exponential and thus memoryless, and hence arrival and service times are Markovian.)

If  $\lambda < \mu$  then  $\pi_k = (1 - \frac{\lambda}{\mu})(\frac{\lambda}{\mu})^k$ , is the unique stationary distribution.

(Shifted geometric distribution with parameter  $1 - \frac{\lambda}{\mu}$ .)

**Example:**  $M/M/k$  queue

$k$  servers.

This is a birth-death process with  $\lambda_n = \lambda, \mu_n = \begin{cases} n\mu & \text{if } n \leq k \\ k\mu & \text{if } n > k \end{cases}$ .

If  $\lambda < k\mu$  then there is a unique stationary distribution.

**Example:**  $M/M/\infty$  queue

infinitely many servers.

This is a birth-death process with  $\lambda_n = \lambda, \mu_n = n\mu$ .

The distribution  $\pi_k = e^{-\lambda/\mu} \frac{(\lambda/\mu)^k}{k!}$ , is the unique stationary distribution.

(Poisson distribution with parameter  $\lambda/\mu$ .)

## 13 Suggested exercises

Basic exercises:

23–26, 28

Extra problems:

c1, B2

Exercises Lawler:

3.12