
Global Convergence Newton

Raffael Colonnello
University of Basel
Raffael.Colonnello@unibas.ch

Fynn Gohlke
University of Basel
Fynn.Gohlke@stud.unibas.ch

Benedikt Heuser
University of Basel
ben.heuser@unibas.ch

Abstract

1 The abstract paragraph should be indented 1/2 inch (3 picas) on both the left- and
2 right-hand margins. Use 10 point type, with a vertical spacing (leading) of 11 points.
3 The word **Abstract** must be centered, bold, and in point size 12. Two line spaces
4 precede the abstract. The abstract must be limited to one paragraph.

5 1 Introduction

6 In this paper we consider problems of the

$$\min_{x \in \mathbb{R}^d} f(\mathbf{x}) \tag{1}$$

7 where $f : \mathbb{R}^d \rightarrow \mathbb{R}$ is a twice-differentiable function. First-order optimization methods are widely
8 used for such problems due to their low per-iteration computational cost and their suitability for
9 parallelization they often suffer from slow convergence for ill-conditioned objective functions
10 [TODO:REFERENCE]. Newton’s method is a popular optimization algorithm that is commonly used
11 to solve optimization problems. It is a second-order optimization algorithm since it uses second-order
12 information of the objective function. Newton’s method is known to have fast local convergence
13 guarantees for convex functions. However, the global convergence properties of Newton’s method
14 are still an active area of research [TODO: REFERENCE]. The purpose of this project is to survey
15 and analyze various strategies to achieve global convergence. In contrast to first-order methods
16 such as gradient descent, second-order methods such as Newton’s method can achieve much faster
17 convergence when presented with ill conditioned Hessians by transferring the problem into a more
18 isotropic optimization problem at the cost of an increase to cubic run time. Newton’s method yields
19 local quadratic convergence if f is twice differentiable (or we have suitable regularity conditions),
20 which degrades to sublinear convergence outside of the local regions.

21 In this paper, we explore the theoretical foundations of several Newton-type methods that achieve
22 different global convergence guarantees, compare their performance in a classification-type problem
23 for two Loss function on four different datasets. Finally we will propose two modifications of the
24 algorithms to achieve an increase in runtime, by either coupling the Newton-type method with a
25 conjugate gradient method for Hessian vector multiplication or Strassens algorithm for fast matrix
26 inversion.

27 This sentence will be cited by the sources so i can test if bibtex is working properly [1]

28 **2 Background**

29 **2.1 Loss function and Datasets**

30 Let $X = \begin{bmatrix} \dots x_1^\top \dots \\ \vdots \\ \dots x_i^\top \dots \\ \vdots \\ \dots x_n^\top \dots \end{bmatrix} \in \mathbb{R}^{n \times d}$ be the set of data for n datapoints with d features, i.e. $x_i \in \mathbb{R}^d$
 31 and labels $y^\top = [y_1, \dots, y_n]$
 32 For $\sigma(z) := \frac{\exp(z)}{1+\exp(z)}$ the loss functions w.r.t. weights ω is given by

$$L_1(\omega) = -\frac{1}{n} \sum_{i=1}^n \left(y_i \log \hat{y}_i + (1 - y_i) \log(1 - \hat{y}_i) \right), \quad \hat{y}_i = \sigma(z_i^\top \omega) \quad (2)$$

$$L_2(\omega) = \frac{1}{n} \sum_{i=1}^n \log(1 + \exp(-y_i z_i^\top \omega)) + r(\omega), \quad r(\omega) = \lambda \sum_{j=1}^d \frac{\alpha \omega_j^2}{1 + \alpha \omega_j^2} \quad (3)$$

33 which yields the two optimization problems

$$\min_{\omega} L_1(\omega) \quad (5)$$

$$\min_{\omega} L_2(\omega) \quad (6)$$

34 The corresponding gradients of L_i are

$$\nabla L_1(x) = \frac{1}{n} X^\top (\hat{y} - y) \quad (7)$$

$$\nabla L_2(x) = -\frac{1}{n} X^\top (y \odot \sigma(-y \odot (X\omega))) + \nabla r(x) \quad (8)$$

$$\text{with } \nabla r(\omega)^\top = \lambda \cdot \left[\frac{2\alpha\omega_1}{(1 + \alpha\omega_1^2)^2}, \dots, \frac{2\alpha\omega_d}{(1 + \alpha\omega_d^2)^2} \right] \quad \text{and } \sigma(\cdot) \text{ applied elementwise}$$

and \odot denotes the entrywise multiplication of vectors.

35 Differentiating again yields the Hessians

$$\nabla^2 L_1(\omega) = \frac{1}{n} X^\top D X \quad (9)$$

$$\nabla^2 L_2(\omega) = \frac{1}{n} X^\top D X + \nabla^2 r(\omega), \quad \nabla^2 r(\omega) = \text{diag} \left(\lambda \frac{2\alpha(1 - 3\alpha\omega_j^2)}{(1 + \alpha\omega_j^2)^3} \right) \quad (10)$$

36 where the diagonal matrix D has entries

$$D_{ii} = \hat{y}_i(1 - \hat{y}_i) = \sigma(-y_i z_i^\top \omega)(1 - \sigma(-y_i z_i^\top \omega)), \quad (11)$$

37 **Observation:**

38 Since $\log(\hat{y}_i), \log(1 - \hat{y}_i)$ are concave on $(0, \infty)$ it follows that $-\log(\hat{y}_i), -\log(1 - \hat{y}_i)$ are convex
 39 and thus L_1 is a linear combination of convex functions (which is again convex). Meanwhile L_2 is
 40 not guaranteed to be convex due to the non-convex regularization term $r(\omega)$.

41 **2.2 Classic Newton's Method**

42 The classical origin of Newton's method is as an algorithm for finding the roots of functions. In
 43 this paper it is used to find the roots x^* of $\nabla(f(x))$ s.t. $\nabla(f(x^*)) = 0$ and x^* a local minimum of f .
 44 Newton's method uses the update rule [REFERENCE]:

$$\mathbf{x}_{k+1} = \mathbf{x}_k - \eta(\nabla^2 f(\mathbf{x}_k))^{-1} \nabla f(\mathbf{x}_k) \quad (12)$$

45 The inverse Hessian can be interpreted as transforming the gradient landscape to be more isotropic,
 46 thereby improving the conditioning of the problem.

47 2.3 Cubic Newton

48 The cubic Newton method was one of the first to achieve a good complexity guarantee globally
49 [REFERENCE TO DO: What convergence rate exactly?]. It is based on cubic regularization and uses
50 the update rule:

$$\mathbf{x}_{k+1} = \mathbf{x}_k - (\nabla^2 f(\mathbf{x}_k) + H \|\mathbf{x}_{k+1} - \mathbf{x}_k\| \mathbf{I})^{-1} \nabla f(\mathbf{x}_k) \quad (13)$$

51 2.4 Levenberg and Marquardt method

52 The Levenberg-Marquardt's algorithm [REFERENCE] is an early form of regularized Newton's
53 method that modifies the Hessian. For ill conditioned (or singular) H this can increase convergence (or
54 make the problem solvable as $H + \lambda I$ is always invertible for sufficiently large $\text{eig}(H) > -\lambda, \lambda >$
55 0).he update rule is:

$$\mathbf{x}_{k+1} = \mathbf{x}_k - (\nabla^2 f(\mathbf{x}_k) + \lambda_k \mathbf{I})^{-1} \nabla f(\mathbf{x}_k) \quad (14)$$

56 2.5 Regularized Newton

57 In their 2023 article Michenko presents a variation of Newton's method that uses the update rule [2]:

$$\mathbf{x}_{k+1} = \mathbf{x}_k - (\nabla^2 f(\mathbf{x}_k) + \sqrt{H \|\nabla f(\mathbf{x}_k)\|} \mathbf{I})^{-1} \nabla f(\mathbf{x}_k) \quad (15)$$

58 where $H > 0$ is a constant. The convergence rate of this algorithm is $\mathcal{O}(\frac{1}{k^2})$. This method uses an
59 adaptive variant of the Levenberg-Marquardt regularization.

60 2.6 TODO

61 Jorge Nocedal & Stephen J. Wright, Numerical Optimization (2nd ed), Springer (2006). Chapter 7
62 (Newton-CG methods for large-scale optimization) Section 7.2: "Newton-CG Algorithm"

64 2.7 Inexact Newton Method

Given that Newton has cubic complexity we now outline how we aim to reduce the runtime by
extending CG and MINRES methods to the Newton-type methods described in our paper. In order
for the modified algorithms to inherit the convergence guarantees of the algorithms we want to
approximate p s.t.

$$\|Hp + \nabla f\| \leq \epsilon \quad (\text{absolute tolerance}) < \epsilon = 10^{-8}$$

Since $H_{1,2} = \nabla^2 L_{1,2}$ are clearly symmetric (as both $X^\top DX$ and $\nabla^2 r(x)$ are) we can apply the
conjugate gradient method if the H is positive definite or have to fall back on MINRES if it is not pd.
Positive definiteness depends on the data matrix and the regularizer curvature. [TODO: runtime for
MINRES and CG]

Every iteration of Vanilla Newton takes $\mathcal{O}(n^3)$ per iteration because inversion of the Hessian costs
 $\mathcal{O}(n^3)$.

for symmetric applying CG to newton drops the effort for conversion down to

$$\mathcal{O}(k \cdot n^2) = \mathcal{O}(\sqrt{\kappa} \log(1/\epsilon) \cdot n^2)$$

65 where $\kappa(H) = \frac{\lambda_{\max}(H)}{\lambda_{\min}(H)}$

66 Precondition with SSOR to reduce condition number.

67 References

- 68 [1] Slavomír Hanzely, Dmitry Kamzolov, Dmitry Pasechnyuk, Alexander Gasnikov, Peter Richtárik,
69 and Martin Takáč. A damped newton method achieves global $\mathcal{O}(\frac{1}{k^2})$ and local quadratic
70 convergence rate. *Advances in Neural Information Processing Systems*, 35:25320–25334, 2022.
- 71 [2] Konstantin Mishchenko. Regularized newton method with global convergence. *SIAM Journal on*
72 *Optimization*, 33(3):1440–1462, 2023.

73 Checklist

74 The checklist follows the references. Please read the checklist guidelines carefully for information on
75 how to answer these questions. For each question, change the default **[TODO]** to **[Yes]** , **[No]** , or
76 **[N/A]** . You are strongly encouraged to include a **justification to your answer**, either by referencing
77 the appropriate section of your paper or providing a brief inline description. For example:

- 78 • Did you include the license to the code and datasets? **[Yes]** See Section
- 79 • Did you include the license to the code and datasets? **[No]** The code and the data are
80 proprietary.
- 81 • Did you include the license to the code and datasets? **[N/A]**

82 Please do not modify the questions and only use the provided macros for your answers. Note that the
83 Checklist section does not count towards the page limit. In your paper, please delete this instructions
84 block and only keep the Checklist section heading above along with the questions/answers below.

85 1. For all authors...

- 86 (a) Do the main claims made in the abstract and introduction accurately reflect the paper's
87 contributions and scope? **[TODO]**
- 88 (b) Did you describe the limitations of your work? **[TODO]**
- 89 (c) Did you discuss any potential negative societal impacts of your work? **[TODO]**
- 90 (d) Have you read the ethics review guidelines and ensured that your paper conforms to
91 them? **[TODO]**

92 2. If you are including theoretical results...

- 93 (a) Did you state the full set of assumptions of all theoretical results? **[TODO]**
- 94 (b) Did you include complete proofs of all theoretical results? **[TODO]**

95 3. If you ran experiments...

- 96 (a) Did you include the code, data, and instructions needed to reproduce the main experi-
97 mental results (either in the supplemental material or as a URL)? **[TODO]**
- 98 (b) Did you specify all the training details (e.g., data splits, hyperparameters, how they
99 were chosen)? **[TODO]**
- 100 (c) Did you report error bars (e.g., with respect to the random seed after running experi-
101 ments multiple times)? **[TODO]**
- 102 (d) Did you include the total amount of compute and the type of resources used (e.g., type
103 of GPUs, internal cluster, or cloud provider)? **[TODO]**

104 4. If you are using existing assets (e.g., code, data, models) or curating/releasing new assets...

- 105 (a) If your work uses existing assets, did you cite the creators? **[TODO]**
- 106 (b) Did you mention the license of the assets? **[TODO]**
- 107 (c) Did you include any new assets either in the supplemental material or as a URL?
108 **[TODO]**
- 109 (d) Did you discuss whether and how consent was obtained from people whose data you're
110 using/curating? **[TODO]**
- 111 (e) Did you discuss whether the data you are using/curating contains personally identifiable
112 information or offensive content? **[TODO]**

113 5. If you used crowdsourcing or conducted research with human subjects...

- 114 (a) Did you include the full text of instructions given to participants and screenshots, if
115 applicable? **[TODO]**
- 116 (b) Did you describe any potential participant risks, with links to Institutional Review
117 Board (IRB) approvals, if applicable? **[TODO]**
- 118 (c) Did you include the estimated hourly wage paid to participants and the total amount
119 spent on participant compensation? **[TODO]**

120 **A Appendix**

121 Optionally include extra information (complete proofs, additional experiments and plots) in the
122 appendix. This section will often be part of the supplemental material.