

Table 14-3.01  
Primary Endpoint Analysis: ADAS Cog (11) - Change from Baseline to Week 24 - LOCF

|                                    | Placebo<br>(N=79) | Xanomeline<br>Low Dose<br>(N=81) | Xanomeline<br>High Dose<br>(N=74) |
|------------------------------------|-------------------|----------------------------------|-----------------------------------|
| Baseline                           |                   |                                  |                                   |
| n                                  | 79                | 81                               | 74                                |
| Mean (SD)                          | 24.1 (12.19)      | 24.4 (12.92)                     | 21.3 (11.74)                      |
| Median (Range)                     | 21.0 ( 5;61)      | 21.0 ( 5;57)                     | 18.0 ( 3;57)                      |
| Week 24                            |                   |                                  |                                   |
| n                                  | 79                | 81                               | 74                                |
| Mean (SD)                          | 26.7 (13.79)      | 26.4 (13.18)                     | 22.8 (12.48)                      |
| Median (Range)                     | 24.0 ( 5;62)      | 25.0 ( 6;62)                     | 20.0 ( 3;62)                      |
| Change from Baseline               |                   |                                  |                                   |
| n                                  | 79                | 81                               | 74                                |
| Mean (SD)                          | 2.5 ( 5.80)       | 2.0 ( 5.55)                      | 1.5 ( 4.26)                       |
| Median (Range)                     | 2.0 (-11;16)      | 2.0 (-11;17)                     | 1.0 ( -7;13)                      |
| p-value(Dose Response) [1][2]      |                   |                                  | 0.245                             |
| p-value(Xan - Placebo) [1][3]      |                   | 0.569                            | 0.233                             |
| Diff of LS Means (SE)              |                   | -0.5 (0.82)                      | -1.0 (0.84)                       |
| 95% CI                             |                   | (-2.1;1.1)                       | (-2.7;0.7)                        |
| p-value(Xan High - Xan Low) [1][3] |                   |                                  | 0.520                             |
| Diff of LS Means (SE)              |                   |                                  | -0.5 (0.84)                       |
| 95% CI                             |                   |                                  | (-2.2;1.1)                        |

[1] Based on Analysis of covariance (ANCOVA) model with treatment and site group as factors and baseline value as a covariate.

[2] Test for a non-zero coefficient for treatment (dose) as a continuous variable

[3] Pairwise comparison with treatment as a categorical variable: p-values without adjustment for multiple comparisons.

Source: <run interactively>

22:28 Sunday, June 28, 2026