

Table 14-3.06
CIBIC+ - Summary at Week 16 - LOCF

	Placebo (N=79)	Xanomeline Low Dose (N=81)	Xanomeline High Dose (N=74)
Week 16			
n	79	81	74
Mean (SD)	4.2 (0.70)	4.0 (0.77)	4.0 (0.75)
Median (Range)	4.0 (3;6)	4.0 (2;6)	4.0 (2;5)
p-value(Dose Response) [1][2]			0.214
p-value(Xan - Placebo) [1][3]		0.219	0.272
Diff of LS Means (SE)		-0.1 (0.12)	-0.1 (0.12)
95% CI		(-0.4;0.1)	(-0.4;0.1)
p-value(Xan High - Xan Low) [1][3]			0.916
Diff of LS Means (SE)			0.0 (0.12)
95% CI			(-0.2;0.2)

[1] Based on Analysis of covariance (ANCOVA) model with treatment and site group as factors.

[2] Test for a non-zero coefficient for treatment (dose) as a continuous variable.

[3] Pairwise comparison with treatment as a categorical variable: p-values without adjustment for multiple comparisons.

Source: C:\cdisc_pilot\PROGRAMS\DRAFT\TFLs\rtf_eff1.sas

21:05 Monday, June 26, 2006