# **RECCAP2-ocean**

# Glossary or "How to ...?"

Latest update: 2022-06-30

## **Content**

Recommended core analysis		
Priority 1	1	
Priority 2	3	
Calculation of flux components from model simulations	3	
Adjustment of river fluxes	6	
Calculation of regional averages and integrals	7	
Recommended units and terminology	7	
Overview of data sets and recommendations for use	7	
Others	10	

## Recommended core analysis

The following core analysis are recommended with priority 1 and 2, for each regional and the global ocean chapter. The instructions for computation distinguish between the two product classes (models and surface flux products), if necessary.

### **Priority 1**

#### Spatially integrated air-sea flux in terms of Fnet at global, regional and biome level

Instructions for computation

- General instructions
  - o use 2D fgco2 fields with monthly resolution
  - spatially integrate fgco2 using the provided area files and the <u>RECCAP2-ocean</u> region mask
  - o compute annual mean values
- Model specific instructions
  - o use 2D fgco2 fields from simA to derive flux time series
  - add <u>river flux adjustment provided through RECCAP2-ocean</u>, spatially integrated at global, regional and biome level<sup>1</sup>
  - account for potential model drift
    - calculate a linear trend through time-series of fgco2 in simB
    - subtract this trend (=drift) from the time-series in simA.
    - Note: This only works for models that use a climatology or single repeated year forcing and not if looping over a full cycle of interannual varying forcing.
- Surface-flux product specific instructions
  - o use 2D fgco2 fields as provided, without any adjustments

#### Reporting of results

- Long-term mean flux for 1985 2018, as ensemble mean and standard deviation for each product class
- Time series of annual mean values
  - o Ensemble mean for each product class
  - Indicator of spread of ensemble members, e.g. the standard deviation or individual lines for each product
- Linear regression trends for the full time period vs pre/post 2000, reported as mean trend ± Std across all individual products of each product class

#### Surface maps of the ensemble mean air-sea CO2 flux in terms of Fnet

Instructions for computation

General instructions

<sup>&</sup>lt;sup>1</sup> if chapters decide to report model results as they are (i.e. without river flux adjustment), it is recommended that differences between the models and surface flux products are discussed in the light of the provided river flux data

- o use 2D fgco2 fields
- o calculate ensemble-mean fgco2 for the period 1985 2018 per product class
- Model specific instructions
  - o use 2D fgco2 fields from simA without adjustments
- Surface-flux product specific instructions
  - o use fgco2 as provided, without adjustments

## Reporting of results

- plot map of ensemble-mean fgco2 for each product class
- show region and biome boundaries, to achieve a connection to the integrated flux time series

## Maps of column inventory changes of Cant (=Fant,ss+ns) for 1994-2007 (models only)

Instructions for computation

- Model specific instructions
  - o use 3D dissic fields for years 1994 and 2007
  - o subtract dissic simD from dissic simA to compute total Cant in each year
  - o subtract total Cant 1994 from total Cant 2007 to compute delta Cant
  - o multiply delta Cant with provided grid-cell volume
  - o integrate delta Cant across the full water column
  - o divide by provided surface area to get column inventory in mol / m<sup>2</sup>

## Reporting of results

- plot delta Cant column inventory maps for
  - o model ensemble-mean
  - observation based estimates by Gruber et al (2019)<sup>2</sup> and OCIM-v2021

## **Priority 2**

#### Separating CO2 and climate effects on mean, spatially integrated air-sea flux (models only)

Instructions for computation

- Model specific instructions
  - o use 2D fgco2 fields
  - calculate regionally integrated, long-term mean fgco2 for the period 1985 -2018 for each model in simA, simB and simC
  - o calculate the CO2 and climate effects for each model
    - CO2 effect (Fant,ss): simC minus simB
    - climate effect (Fnat,ns + Fant,ns): simA minus simC
    - Note: the separation of <u>flux components</u> is also illustrated in the following chapter of this glossary

<sup>&</sup>lt;sup>2</sup> Cant inventory changes reported by Gruber et al. (2019) are only integrated to 3000m water depth. Differences between their estimates and the models should also be discussed taking into account the modelled Cant accumulation below 3000m water depth

## Reporting of results

- ensemble mean trend ± Std separately for CO2 effect and climate effects
  - $\circ \quad$  could be a table, bar plot or just reported within the text

## Calculation of flux components from model simulations

As a reminder, in the protocol model simulations are defined as:

	Simulation A	Simulation B 'ctrl'	Simulation C	Simulation D
Request level	Mandatory, Tier 1	Mandatory for global models, Tier 1	Tier 2	Tier 2
Atmospheric CO <sub>2</sub>	Increasing	Constant, preindustrial	Increasing	Constant preindustrial
Atmospheric forcing (wind, fluxes of heat and freshwater)	Variable, reanalysis	climatological	climatological	Variable, reanalysis
Considered components <sup>(1)</sup>	$\begin{aligned} & Flux_{ant}^{ ss} + Flux_{ant}^{ ns} + \\ & Flux_{nat}^{ ss} + Flux_{nat}^{ ns} \end{aligned}$	Flux <sub>nat</sub> ss	Flux <sub>ant</sub> ss + Flux <sub>nat</sub> ss	Flux <sub>nat</sub> ss + Flux <sub>nat</sub> ns

<sup>(1)</sup> Total Flux = Flux<sub>ant</sub> ss + Flux<sub>nat</sub> is the air-sea flux of anthropogenic CO<sub>2</sub>, and Flux<sub>nat</sub> that of natural CO<sub>2</sub>. "ss" refers to steady-state and "ns" to non-steady state, i.e., the variability component imposed on natural and anthropogenic CO<sub>2</sub> by all non-seasonal variability. The same differentiation applies also to the change in ocean interior DIC, i.e.,  $\Delta$ DIC =  $\Delta$ DIC<sub>ant</sub> ss +  $\Delta$ DIC ant ss +  $\Delta$ DIC nat should be change in ocean interior DIC, i.e.,  $\Delta$ DIC =  $\Delta$ DIC ant ss +  $\Delta$ DIC nat should be change in ocean interior DIC, i.e.,  $\Delta$ DIC =  $\Delta$ DIC ant ss +  $\Delta$ DIC nat should be change in ocean interior DIC, i.e.,  $\Delta$ DIC =  $\Delta$ DIC nat should be change in ocean interior DIC, i.e.,  $\Delta$ DIC =  $\Delta$ DIC nat should be change in ocean interior DIC, i.e.,  $\Delta$ DIC =  $\Delta$ DIC nat should be change in ocean interior DIC, i.e.,  $\Delta$ DIC =  $\Delta$ DIC nat should be change in ocean interior DIC, i.e.,  $\Delta$ DIC =  $\Delta$ DIC nat should be change in ocean interior DIC, i.e.,  $\Delta$ DIC =  $\Delta$ DIC nat should be change in ocean interior DIC, i.e.,  $\Delta$ DIC =  $\Delta$ DIC nat should be change in ocean interior DIC, i.e.,  $\Delta$ DIC nat should be change in ocean interior DIC, i.e.,  $\Delta$ DIC nat should be change in ocean interior DIC, i.e.,  $\Delta$ DIC nat should be change in ocean interior DIC, i.e.,  $\Delta$ DIC nat should be change in ocean interior DIC, i.e.,  $\Delta$ DIC nat should be change in ocean interior DIC, i.e.,  $\Delta$ DIC nat should be change in ocean interior DIC, i.e.,  $\Delta$ DIC nat should be change in ocean interior DIC, i.e.,  $\Delta$ DIC nat should be change in ocean interior DIC, i.e.,  $\Delta$ DIC nat should be change in ocean interior DIC, i.e.,  $\Delta$ DIC nat should be change in ocean interior DIC nat should be change in oc

# According to the definition of the model simulations above, different carbon flux components should be calculated as follows:

#### Fant,ss = SimC - SimB

This is equivalent to the effect of rising CO<sub>2</sub> alone on the ocean carbon sink.

#### Fnat,ns + Fant,ns = SimA - Sim-C

This describes the effect of climate change and variability on the ocean carbon sink

#### S ocean = SimA - SimB


This corresponds to S\_ocean reported in the Global Carbon Budget and corrects for model drift (=trend in SimB) and biases (=Offset from zero in SimB). Note: This subtraction only works on a global scale, as the assumption that CO2 flux (without rivers) was zero does not hold on a regional scale.

#### Fnat,ns = SimD - SimB

This describes the climate change effect on the natural carbon flux.

#### Fant,ss + Fant,ns = SimA - SimD

This describes the total anthropogenic flux (comparable to Gruber et al. 2019)



### Overview on C-flux components, their calculation and representation by product types. Footnotes indicate:

- 1. Please note that most RECCAP2-ocean models are run without riverine carbon input.
- 2. When outgassing of riverine C input is subtracted, the flux into the ocean increases.
- 3. Not applicable to OCIM models because there is no SimD.
- 4. S\_ocean determined as SimA SimB is only applicable to global models, and should not be interpreted based on regional models.

# **Adjustment of river fluxes**

- Recommended global river flux adjustment:
  - o 0.65 PgC/yr (Regnier et al., 2022)
- Recommended regional river flux adjustments
  - Spatial distribution is the gridded field from <u>Lacroix et al. (2020</u>). CAVEAT: The gridded field requires an upscaling from 0.2 to 0.65 PgC/yr because of too low carbon input to the ocean (P. Regnier, pers. comm.)
  - Upscaled river flux fields are available here: <u>https://reccap2-ocean.github.io/river/</u>
  - Please use our region and biome mask to integrate river fluxes spatially: <a href="https://reccap2-ocean.github.io/regions/">https://reccap2-ocean.github.io/regions/</a>
- Other useful river flux estimates include: <u>Jacobson et al. (2007)</u>, <u>Resplandy et al 2018</u>, Table S1 below from the <u>auxiliary material</u> in <u>Gruber et al. (2009)</u>, <u>Aumont et al. (2001)</u>, and GCB papers.

## Calculation of regional averages and integrals

Assignment of basins and subregions

- Link to basin mask file: <a href="https://reccap2-ocean.github.io/regions/">https://reccap2-ocean.github.io/regions/</a>
- Use of RECCAP2-ocean mask is mandatory on basin scale (i.e. for the definition of chapter boundaries)
- Use of the provided mask on the subregion scale is voluntary

#### Calculation of regionally averaged surface pCO2

- weight gridded spco2 by area of grid cell (from area file provided by data provider)
- take information on the fraction of cells filled with water into account if available
- Note: for the calculation of regionally averaged air-sea disequilibria, Kw (and thus wind speed) are usually taken into account. In this case, the difference between atmospheric pCO2 and regionally averaged spco2 as defined here, would not be identical to regionally averaged disequilibria.

#### Calculation of regionally integrated air-sea CO2 fluxes

- multiply fgco2 with area and sum up over region of interest
- take information on the fraction of cells filled with water into account if available
- convert from molC/s to PgC/yr
  - from per seconds to per year (\* 365.25 \* 24 \* 60 \* 60)
  - o convert from moles C to grams C (\* 12.011 g/mol)

#### Calculation of regional ocean interior integrals

- multiply concentration with volume grid provided by data provider
- take information on the fraction of cells filled with water into account if available

Use of observation-based surface pCO2 products and models (primarily data-assimilated models) that don't cover the full period

- for averages and integrals, exclude data sets from ensemble that do not cover full period
  - full period observations: 1985 to 2018
  - o full period models: 1980 to 2018

# **Recommended units and terminology**

- Carbon inventory: PgC
- Carbon fluxes: PgC yr<sup>-1</sup> (or TgC yr<sup>-1</sup> if appropriate)
- Partial pressure of CO<sub>2</sub>: μatm
- Carbon flux density: mol m-2 yr-1 or mmol m-2 d-1 (outgassing positive, Note: this is opposite to the protocol)

# Overview of data sets and recommendations for use

Note: This information will be moved to a meta data table that is under construction.

model_id	Main features and recommendations for use			
(+ label for publication)				
Group: Global hindcast models				
General remarks: Simulation A of hindcast models covers $F_{co2}$ = Fnat,ss + Fnat,ns + Fant,ss +				
Fant,ns . *Some models also include riverine fluxes (Friv), but often of small magnitude.				
CESM-ETHZ				
CCSM-WHOI	Model output taken from coastal chapter. No specific formatting according to global/regional chapter protocol. Not qc'ed. Ends in 2017.			
CNRM-ESM2-1*				
EC-Earth3				
FESOM_REcoM_LR (FESOM-1.4-REcoM2-LR)	If a chapter would like to use only one FESOM-REcoM simulation, use LR, it has a longer spin-up and is more			
	suitable for analysis of anthropogenic CO2 time-series			
FESOM_RECOM_HR (FESOM-1.4-RECOM2-HR)	High-resolution, shorter spin-up, could be interesting in addition to LR for processes			
MOM6-COBALT2-Princeton (MOM6-Princeton)				
MPIOM-HAMOCC				
MRI-ESM2-0				
NorESM-OC1.2				
ORCA025-GEOMAR	The model simulations have a large positive bias in pCO2 in the upwelling areas off South America and Africa. Consider interpreting the results accordingly, neglecting these areas from your analysis, or excluding this model completely.			
ORCA1-LIM3-PISCES	, , , ,			
(IPSL-NEMO-PISCES)				
Planktom12				
Group: Global data-assimilated models General remarks: Suggest to not average them, but show individually as it is a small				
ensemble. They cover different years, and have quite different results.				
	s cover F <sub>CO2</sub> = Fnat,ss + Fnat,ns + Fant,ss + Fant,ns.			
ECCO-Darwin				
OCIM-v2014-CTL	The OCIM models include simulations A, B, and C and can be treated in the same way as the other models. The point to be aware of is that OCIM is an abiotic model, so the natural fluxes are only the abiotic ones. Also, both OCIM versions have constant circulation. Useful for anthropogenic carbon (Fant,ss) and abiotic carbon. No river fluxes.			

OCIM-v2021	Like OCIM-v2014, but using updated spatially-varying	
	atmospheric pCO2 and time-varying SST and gas transfer	
	velocity. Useful for anthropogenic carbon (Fant,ss) as well	
	as the temperature-forced variability of natural carbon	
	(Fnat,ns forced by temperature). No river fluxes.	
	[ (That, its forced by temperature). No river fluxes.	
Group: Regional hindcast mod	lels	
	dels should only be used in the corresponding regional	
chapters.	acis should only be used in the corresponding regional	
B-SOSE	Southern Ocean (use only in SO chapter), group with	
	data-assimilated models.	
GOA-COBALT	Use only in Pacific chapter	
INCOIS-BIO-ROMS	Use only in Indian Ocean chapter	
ROMS-Southern Ocean-ETHZ	Had to be withdrawn due to errors in the initialization. Will	
Nows southern occur Emile	be removed from the FTP server. Please do not use it!	
ROMS-Pacific-ETHZ	Use only in Pacific chapter	
ROMS-Atlantic-ETHZ	Use only in Atlantic chapter	
ROMS-NYUAD	Use only in Indian Ocean chapter	
NOWIS-NTOAD	Ose only in indian ocean chapter	
Group: Surface CO <sub>2</sub> observation	on based products	
•	based flux products measure the contemporary CO2 flux Fnet	
= Fnat,ss + Fnat,ns + Fant,ss + I		
AOML EXTRAT	Starts in ~1997	
AOIVIL_EXTRAI	Suggestion: exclude from ensembles that average over the	
	full time-period	
CMEMS-LSCE-FFNN	Tull tille-period	
CIVILIVIS-LSCL-I I IVIV		
CSIRML6		
JENA-MLS		
(CarboScope)		
JMAMLR		
NIES-nn		
OceanSODAETHZ		
SOMFFN	Contains smoothened (labelled as spco2/fgco2) fields,	
	which should be used for general analysis. Yet, raw output	
	is also provided (labelled as spco2_raw/fgco2_raw).	
spco2 LDEO HPD		
LDEO 2021 clim RECCAP2	Contains climatological fields only. Updated of Takahashi's	
	pCO <sub>2</sub> climatology.	
UOEX WAT20	Applies adjustments to fCO2 data, hence a different type of	
<del>-</del>	data-product, don't average with the other data-products	
	Uses CCMP wind data only available from 1988 on-wards.	
	For 1985-87 it uses a wind climatology: don't use these	
	years.	
	,	
Group: Ocean interior observa	ation-based products	
General remarks:		

dissic_MOBO-DIC-MPIM	Near-global ocean interior monthly climatology of DIC			
dissicNNGv2LDEO				
ETHZ_Cant_eMLR	Global ocean interior change in Cant storage 1994-2007.			
UW_NOAA_Cant_Pacific				
Group: Atmospheric inversions				
General remarks:				

# **Others**

- Recommended properties for plots
  - o Color scales and ranges
    - outgassing red, uptake blue