


DPP - 1

SOLUTION

Link to View Video Solution:  [Click Here](#)

1. $\frac{r_1}{r_2} = \frac{6}{5}$

$$\frac{m_1}{m_2} = \frac{9}{4}, \quad \frac{q_1}{q_2} = ?$$

$$r = \frac{mv}{qB} = \frac{m}{qB} \sqrt{2k}$$

$$r = \frac{1}{qB} \sqrt{2mk}$$

$$\frac{r_1}{r_2} = \sqrt{\frac{m_1}{m_2}} \times \frac{q_2}{q_1} \Rightarrow \frac{6}{5} = \sqrt{\frac{9}{4}} \times \frac{q_2}{q_1}$$

$$\Rightarrow \frac{q_1}{q_2} = \frac{5}{4}$$

$$q_1 + q_2 = p + q = 9$$

2. $\vec{B} = (2\hat{i} + 3\hat{j})T$

$$\vec{a} = (\alpha\hat{i} - 4\hat{j})m/s^2$$

$$\vec{F} \perp \vec{B} \Rightarrow \vec{a} \perp \vec{B}$$

$$\vec{a} \cdot \vec{B} = 0$$

$$\Rightarrow 2\alpha - 12 = 0$$

$$\alpha = 6$$

3. Proton, deuteron, α -particle

$$m_p \quad 2m_p \quad \approx 4m_p$$

$$q = e \quad q = e \quad q^1 = 2e$$

$$r_p = \frac{\sqrt{2m_p k}}{eB} \quad r_d = \frac{2 \times 2m_p k}{eB}$$

$$r_\alpha = \frac{\sqrt{2 \times 4m_p k}}{2eB}$$

$$r_p : r_d : r_\alpha = 1 : \sqrt{2} : 1 = k : \frac{p}{\sqrt{2}} : 1$$

$$k = 1 \quad p = 2$$

$$k + p = 3$$

Link to View Video Solution: [Click Here](#)

4. (d) : Given : $v = 4 \times 10^5$ m/s, $\theta = 60^\circ$, $m = 1.67 \times 10^{-27}$ kg,


$$q = 1.6 \times 10^{-19} \text{ C}, B = 0.3 \text{ T}$$

$$\text{Time of one revolution, } T = \frac{2\pi m}{qB}$$

$$l = (v \cos 60^\circ) \times (10T) = \frac{10\pi m v}{qB}$$

$$\Rightarrow l = \frac{10 \times 3.14 \times (1.67 \times 10^{-27}) \times (4 \times 10^5)}{(1.6 \times 10^{-19})(0.3)}$$

$$\text{Here, } l = 0.4369 \text{ m} \approx 0.444 \text{ m}$$



5. (b) : Given : $q = 1 \mu\text{C}$; $\vec{v} = 2\hat{i} + 3\hat{j} + 4\hat{k}$ m/s

$$\vec{B} = (5\hat{i} + 3\hat{j} - 6\hat{k}) \times 10^{-3} \text{ T}; \vec{F}_M = \vec{F} \times 10^{-9} \text{ N}$$

$$\text{Magnetic force, } \vec{F}_M = q(\vec{v} \times \vec{B}); \vec{v} \times \vec{B} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 2 & 3 & 4 \\ 5 & 3 & -6 \end{vmatrix} \times 10^{-3}$$

$$= [\hat{i}(-18 - 12) - \hat{j}(-12 - 20) + \hat{k}(6 - 15)] \times 10^{-3}$$

$$= (-30\hat{i} + 32\hat{j} - 9\hat{k}) \times 10^{-3}$$

$$\therefore \vec{F}_M = 1 \times 10^{-6} \times 10^{-3}(-30\hat{i} + 32\hat{j} - 9\hat{k})$$

$$\vec{F}_M = (-30\hat{i} + 32\hat{j} - 9\hat{k}) \times 10^{-9} \text{ N}$$

$$\text{So, } \vec{F} = -30\hat{i} + 32\hat{j} - 9\hat{k}$$

6. $R_1 < R_2$


$$\frac{m_1 v}{q_1 B} < \frac{m_2 v}{q_2 B} \Rightarrow \frac{m_1}{q_1} < \frac{m_2}{q_2}$$

7. $R = \frac{mv}{qB} = \frac{p}{qB}$

$$R = \frac{p}{qB}$$

$$\sin \theta = \frac{d}{R} = \frac{dQB}{p}$$

$$\sin \theta = \frac{dQB}{p} \quad (\text{A})$$




8. \vec{B} outward


$$R = \frac{mv}{qB}$$

$$R \downarrow \text{ ses} \quad v \downarrow \text{ ses}$$

9. $R = \frac{mv}{qB}$

$$T = \frac{2\pi m}{qB}$$



Link to View Video Solution:  [Click Here](#)

$$f = \frac{1}{T} = \frac{qB}{2\pi m}$$

10. $R = \frac{mv}{qB}$

$$\sin\theta = \frac{d}{R} = \frac{dqB}{mv}$$

$$\frac{q}{m} = \frac{v \sin\theta}{dB} \quad (D)$$

