

HOMework-1

1. If  $y = \frac{a + bx^{\frac{3}{5}}}{x^4}$  &  $\frac{dy}{dx}$  vanishes when  $x = 5$  then  $\frac{a}{b} =$   
 (A)  $\sqrt{3}$  (B) 2 (C)  $\sqrt{5}$  (D) 3
2. If  $\frac{d}{dx} \left( \frac{1+x^2+x^4}{1+x+x^2} \right) = ax + b$  then the value of a and b are respectively  
 (A) 2 and 1 (B) -2 and 1 (C) 2 and -1 (D) 3 and 1
3. If  $y = x - x^2$ , then the derivative of  $y^2$  w.r.t.  $x^2$  is  
 (A)  $2x^2 + 3x - 1$  (B)  $2x^2 - 3x + 1$  (C)  $2x^2 + 3x + 1$  (D)  $2x^2 + 5x + 1$
4. The differential coefficient of  $a^{\sin^{-1}x}$  w.r.t.  $\sin^{-1}x$  is -  
 (A)  $a^{\sin^{-1}x} \log_e a$  (B)  $a^{\sin^{-1}x}$  (C)  $\frac{a^{\sin^{-1}x}}{\sqrt{(1-x^2)}}$  (D)  $a^{\sin^{-1}x} \sqrt{(1-x^2)}$
5. The value of derivative of  $\tan^{-1} \left( \frac{2x\sqrt{1-x^2}}{1-2x^2} \right)$  w.r.t  $\sec^{-1} \left( \frac{1}{2x^2-1} \right)$  at  $x = 1/2$  equals-  
 (A) 1 (B) -1 (C) 0 (D) 2
6. If  $y = \cos^{-1}(\cos x)$  then  $\frac{dy}{dx}$  at  $x = \frac{5\pi}{4}$  is equal to  
 (A) 1 (B) -1 (C)  $\frac{1}{\sqrt{2}}$  (D)  $-\frac{1}{\sqrt{2}}$
7. If  $8f(x) + 6f\left(\frac{1}{x}\right) = x + 5$  and  $y = x^2f(x)$ , then  $\frac{dy}{dx}$  at  $x = -1$  is equal to  
 (A) 0 (B)  $\frac{1}{14}$  (C)  $-\frac{1}{14}$  (D) 14
8. If  $f(x) = x^n$ , then the value of  $f(1) - \frac{f'(1)}{1!} + \frac{f''(1)}{2!} - \frac{f'''(1)}{3!} + \dots + \frac{(-1)^n f^{(n)}(1)}{n!}$  is  
 (A)  $2^n$  (B)  $2^{n-1}$  (C) 0 (D) 1
9. If f is differentiable in  $(0, 6)$  &  $f'(4) = 5$  then  $\lim_{x \rightarrow 2} \frac{f(4) - f(x^2)}{2-x} =$   
 (A) 5 (B)  $\frac{5}{4}$  (C) 10 (D) 20
10. If  $u = ax + b$  then  $\frac{d^n}{dx^n} (f(ax + b))$  is equal to  
 (A)  $\frac{d^n}{du^n} (f(u))$  (B)  $a \frac{d^n}{du^n} (f(u))$  (C)  $a^n \frac{d^n}{du^n} (f(u))$  (D)  $a^{-n} \frac{d^n}{du^n} (f(u))$
11. Let  $f(x)$  be a polynomial in x. Then the second derivative of  $f(e^x)$ , is  
 (A)  $f''(e^x) \cdot e^x + f'(e^x)$  (B)  $f''(e^x) \cdot e^{2x} + f'(e^x) \cdot e^{2x}$   
 (C)  $f'(e^x) e^{2x}$  (D)  $f''(e^x) \cdot e^{2x} + f'(e^x) \cdot e^x$
12. If  $y = f(x)$  is an odd differentiable function defined on  $(-\infty, \infty)$  such that  $f'(3) = -2$ , then  $f'(-3)$  equals  
 (A) 4 (B) 2 (C) -2 (D) 0

(MATHEMATICS)

METHOD OF DIFFERENTIATION

13. If  $g$  is inverse of  $f$  and  $f'(x) = \frac{1}{1+x^n}$ , then  $g'(x)$  equals -  
 (A)  $1+x^n$  (B)  $1+(f(x))^n$  (C)  $1+(g(x))^n$  (D)  $1-x^n$
14. Derivative of  $\log_e (\log_e |\sin x|)$  with respect to  $x$  at  $x = \frac{\pi}{6}$  is  
 (A)  $-\frac{\sqrt{3}}{\log_e 2}$  (B)  $\frac{\sqrt{3}}{\log_e 2}$  (C)  $-\frac{\sqrt{3}}{2\log_e 2}$  (D) does not exist
15. If  $f(x) = f'(x) + f''(x) + f'''(x) + f''''(x) + \dots + \infty$  also  $f(0) = 1$  and  $f(x)$  is a differentiable function indefinitely then  $f(x)$  has the value  
 (A)  $e^x$  (C)  $e^{x/2}$  (B)  $e^{2x}$  (D)  $e^{4x}$
16. If  $f(x) = |(x-4)(x-5)|$ , then  $f'(x)$  is  
 (A)  $-2x+9$ , for all  $x \in \mathbb{R}$  (B)  $2x-9$  if  $x > 5$   
 (C)  $-2x+9$  if  $4 < x < 5$  (D) not defined for  $x = 4, 5$
17. If  $f$  is twice differentiable such that  $f''(x) = -f(x)$  and  $f'(x) = g(x)$ . If  $h(x)$  is twice differentiable function such that  $h'(x) = [f(x)]^2 + [g(x)]^2$ . If  $h(0) = 2$ ,  $h(1) = 4$ , then the equation  $y = h(x)$  represents  
 (A) a curve of degree 2 (B) a curve passing through the origin  
 (C) a straight line with slope 2 (D) a straight line with  $y$  intercept equal to 2.
18. Two functions  $f$  &  $g$  have first & second derivatives at  $x = 0$  satisfy the relations,  
 $f(0) = \frac{2}{g(0)}$ ,  $f'(0) = 2$ ,  $g'(0) = 4$ ,  $g''(0) = 5$ ,  $f''(0) = g(0) = 3$  then  
 (A) if  $h(x) = \frac{f(x)}{g(x)}$  then  $h'(0) = \frac{32}{9}$  (B) if  $k(x) = f(x) \cdot g(x) \sin x$  then  $k'(0) = 2$   
 (C)  $\lim_{x \rightarrow 0} \frac{g'(x)}{f'(x)} = \frac{1}{2}$  (D)  $f'(x) = g'(x)$
19. Differentiate the following functions with respect to  $x$ .  
 (i)  $x^{2/3} + 7e - \frac{5}{x} + 7 \tan x$  (ii)  $\ln \tan \left( \frac{\pi}{4} + \frac{x}{2} \right)$   
 (iii)  $\frac{\sin x - x \cos x}{x \sin x + \cos x}$  (iv)  $\tan \left( \tan^{-1} \sqrt{\frac{1 - \cos x}{1 + \cos x}} \right)$
20. Differentiate  $x^2 \cdot \ln x \cdot e^x$  with respect to  $x$ .
21. If  $\cos \frac{x}{2} \cdot \cos \frac{x}{2^2} \cdot \cos \frac{x}{2^3} \dots \infty = \frac{\sin x}{x}$  then find the value of  $\frac{1}{2^2} \sec^2 \frac{x}{2} + \frac{1}{2^4} \sec^2 \frac{x}{2^2} + \frac{1}{2^6} \sec^2 \frac{x}{2^3} + \dots \infty$ .
22. Let  $f, g$  and  $h$  are differentiable functions. If  $f(0) = 1$ ;  $g(0) = 2$ ;  $h(0) = 3$  and the derivatives of their pair wise products at  $x = 0$  are  $(fg)'(0) = 6$ ;  $(gh)'(0) = 4$  and  $(hf)'(0) = 5$  then compute the value of  $(fgh)'(0)$ .
23. If  $f: \mathbb{R} \rightarrow \mathbb{R}$  is a function such that  $f(x) = x^3 + x^2 f'(1) + x f''(2) + f'''(3)$  for all  $x \in \mathbb{R}$ , then prove that  $f(2) = f(1) - f(0)$ .

Paragraph for Question Nos. 24 to 26

$f(x)$  is a polynomial function  $f : \mathbb{R} \rightarrow \mathbb{R}$  such that  $f(2x) = f'(x)f''(x)$ .

24. The value of  $f(3)$  is  
 (A) 4 (B) 12 (C) 15 (D) 11
25.  $f(x)$  is :  
 (A) one-one and onto (B) one-one and into  
 (C) many-one and onto (D) many-one and into
26. Equation  $f(x) = x$  has:  
 (A) three real and positive roots (B) three real and negative roots  
 (C) one real root (D) three real such that sum of roots is zero
27. Let  $f : (-1, 1) \rightarrow \mathbb{R}$  be a differentiable function with  $f(0) = -1$  and  $f'(0) = 1$ .  
 Let  $g(x) = [f(2f(x) + 2)]^2$ , then  $g'(0) =$  [AIEEE 2010]  
 (A) 4 (C) 0 (B) -4 (D) -2
28. If  $f(x) = \frac{2x-4}{x^2-1}$  and  $f'(x) = \frac{p}{(x^2-1)^2}$ , then  $p$  equals-  
 (A)  $x^2 - 8x - 2$  (B)  $-2x^2 + 8x + 2$   
 (C)  $4x + 2$  (D)  $-2x^2 + 8x - 2$
29. If  $y = \sqrt{\frac{1-\cos x}{1+\cos x}}$ ,  $x \in (0, \pi)$  then  $\frac{dy}{dx}$  equals-  
 (A)  $\frac{1}{2} \sec^2 x/2$  (B)  $\frac{1}{2} \operatorname{cosec}^2 x/2$   
 (C)  $\sec^2 x/2$  (D)  $\operatorname{cosec}^2 x/2$
30.  $\frac{d}{d\theta} \left\{ \tan^{-1} \left( \frac{1-\cos \theta}{\sin \theta} \right) \right\}$  equals-  
 (A)  $1/2$  (B) 1 (C)  $\sec \theta$  (D)  $\operatorname{cosec} \theta$
31.  $d/dx(\sec x^\circ)$  equals -  
 (A)  $\sec x \tan x$  (B)  $\sec x^\circ \tan x^\circ$   
 (C)  $\left(\frac{\pi}{180}\right) \sec x^\circ \tan x^\circ$  (D)  $\left(\frac{\pi}{180}\right) \sec x \tan x$
32. If  $\operatorname{acos}^2(x+y) = b$ , then  $dy/dx$  equals-  
 (A) 2 (B) -2 (C) 1 (D) -1
33. If  $y = \log \sqrt{\frac{1-\sin x}{1+\sin x}}$ , then  $dy/dx$  equals-  
 (A)  $\sec x$  (B)  $-\sec x$   
 (C)  $\operatorname{cosec} x$  (D)  $\sec x \tan x$

(MATHEMATICS)

METHOD OF DIFFERENTIATION

34. If  $y = \log_{10} (\sin x)$ , then  $dy/dx$  equals-
- (A)  $\sin x \log_{10} e$  (B)  $\cos x \log_{10} e$   
 (C)  $\cot x \log_{10} e$  (D)  $\cot x$
35. The derivative of  $x|x|$  is-
- (A)  $2x$  (B)  $-2x$  (C)  $2|x|$  (D) Does not exist

ANSWER KEY

1. (C) 2. (C) 3. (B) 4. (A) 5. (B) 6. (B) 7. (C)  
 8. (C) 9. (D) 10. (C) 11. (D) 12. (C) 13. (C) 14. (D)  
 15. (B) 16. (BCD) 17. (CD) 18. (ABC)
19. (i)  $\frac{2}{3}x^{-\frac{1}{3}} + \frac{5}{x^2} + 7\sec^2 x$  (ii)  $\sec x$   
 (iii)  $\frac{x^2}{(x \sin x + \cos x)^2}$  (iv)  $\frac{1}{2}\sec^2\left(\frac{x}{2}\right)$
20.  $x^2 \ln x \cdot e^x + xe^x + 2xe^x \ln x$  21.  $\operatorname{cosec}^2 x - \frac{1}{x^2}$
22. 16 24. (B) 25. (A) 26. (D) 27. (B) 28. (D) 29. (A)  
 30. (A) 31. (C) 32. (D) 33. (B) 34. (C) 35. (C)