

EXERCISE # (O-I)


Electrode Potential, Galvanic Cell

1. In a galvanic cell
 - (A) chemical reaction produces electrical energy
 - (B) electrical energy produces chemical reaction
 - (C) reduction occurs at anode
 - (D) oxidation occurs at cathode
2. Which of the following statement is wrong about galvanic cell ?
 - (A) Cathode is positive charged
 - (B) Anode is negatively charged
 - (C) Reduction takes place at the anode
 - (D) Electrons move from anode to cathode
3. Which of the following has been universally accepted as a reference electrode at all temperatures and has been assigned a value of zero volt?
 - (A) Standard platinum electrode
 - (B) Standard copper electrode
 - (C) Standard graphite electrode
 - (D) Standard hydrogen electrode
4. A standard hydrogen electrode has zero electrode potential because
 - (A) hydrogen is easier to oxidise
 - (B) electrode potential is assumed to be zero
 - (C) hydrogen atom has only one electron
 - (D) hydrogen is the lightest element.
5. Which is not true for a standard hydrogen electrode ?
 - (A) The hydrogen ion concentration must be 1 M
 - (B) Temperature must be 25°C
 - (C) Pressure of hydrogen must be 1 bar
 - (D) Temperature can vary
6. The equation representing the process by which standard reduction potential of zinc can be defined is :
 - (A) $\text{Zn}^{2+}(\text{s}) + 2\text{e}^- \longrightarrow \text{Zn}(\text{s})$
 - (B) $\text{Zn}(\text{g}) \longrightarrow \text{Zn}^{2+}(\text{g}) + 2\text{e}^-$
 - (C) $\text{Zn}^{2+}(\text{g}) + 2\text{e}^- \longrightarrow \text{Zn}(\text{s})$
 - (D) $\text{Zn}^{2+}(\text{aq.}) + 2\text{e}^- \longrightarrow \text{Zn}(\text{s})$

 $\Delta G = -nFE$ theoretical

7. E° for $\text{F}_2 + 2\text{e}^- = 2\text{F}^-$ is 2.8 V, E° for $\frac{1}{2}\text{F}_2 + \text{e}^- = \text{F}^-$ is ?
 - (A) 2.8 V
 - (B) 1.4 V
 - (C) -2.8 V
 - (D) -1.4 V
8. Doubling all the coefficients in the equation for the cell reaction -
 - (A) Doubles both E° and ΔG°
 - (B) Doubles E° , but does not change ΔG°
 - (C) Doubles ΔG° , but does not change E°
 - (D) Does not change E° or ΔG°

9. $E^\circ = 0.6 \text{ V}$ for the following reaction :





What is ΔG° for this reaction at 25°C -

- (A) -115.8 kJ (B) -57.9 kJ (C) 57.9 kJ (D) 115.8 kJ

$$E_1 + E_2 = E_3$$

10. The standard electrode potentials for the reactions



at 25°C are 0.80 volt and -0.14 volt , respectively. The standard emf of the cell formed by these half-cells is

- (A) 0.66 volt (B) 0.80 volt (C) 1.08 volt (D) 0.94 volt





11. $E^\circ(\text{Ni}^{2+}/\text{Ni}) = -0.25 \text{ volt}$, $E^\circ(\text{Au}^{3+}/\text{Au}) = 1.50 \text{ volt}$. The standard emf of the voltaic cell.

- (A) 1.25 volt (B) -1.75 volt (C) 1.75 volt (D) 4.0 volt

12. For the cell reaction $2\text{Ce}^{4+} + \text{Co} \longrightarrow 2\text{Ce}^{3+} + \text{Co}^{2+}$ $E^\circ_{\text{cell}} = 1.89 \text{ V}$.

If $E^\circ_{\text{Co}^{2+}/\text{Co}}$ is -0.28 V , what is the value of $E^\circ_{\text{Ce}^{4+}/\text{Ce}^{3+}}$?

13. From the following E° values of half cells,




What combination of two half cells would result in a cell with the largest potential ?

- (A) (ii) and (iii) (B) (ii) and (iv) (C) (i) and (iii) (D) (i) and (iv)

14. The standard reduction potentials for two half-cell reactions are given below,





The standard free energy change for the reaction




- (A) 115.8 kJ (B) -115.8 kJ (C) -231.6 kJ (D) 231.6 kJ

15. Given electrode potentials :



E°_{cell} for the cell reaction,



- (A) $(2 \times 0.771 - 0.536) = 1.006 \text{ volts}$ (B) $(0.771 - 0.5 \times 0.536) = 0.503 \text{ volts}$
 (C) $0.771 - 0.536 = 0.235 \text{ volts}$ (D) $0.536 - 0.771 = -0.235 \text{ volts}$

Electrode Potential And Equilibrium Constant

16. If $E^\circ_{\text{Au}^+/\text{Au}}$ is 1.69 V & $E^\circ_{\text{Au}^{3+}/\text{Au}}$ is 1.40 V, then $E^\circ_{\text{Au}^{3+}/\text{Au}^+}$ will be
 (A) 0.19 V (B) 2.945 V (C) 1.255 V (D) None
17. What is the standard electrode potential for the electrode $\text{MnO}_4^- / \text{MnO}_2$ in solution
 $E^\circ_{\text{MnO}_4^-/\text{Mn}^{2+}} = 1.51\text{volt}$, $E^\circ_{\text{MnO}_2/\text{Mn}^{2+}} = 1.23\text{volt}$
 (A) 0.7 V (B) 3.33 V (C) 1.70 V (D) 2.5 V
18. Calculate the equilibrium constant for the reaction $\text{Fe} + \text{CuSO}_4 \rightleftharpoons \text{FeSO}_4 + \text{Cu}$ at 25°C .
 Given $E^\circ(\text{Fe}/\text{Fe}^{2+}) = 0.44\text{V}$, $E^\circ(\text{Cu}/\text{Cu}^{2+}) = -0.337\text{V}$.
19. At 25°C the standard emf of cell having reactions involving two electrons change is found to be 0.295V. The equilibrium constant of the reaction is –
 (A) 29.5×10^{-2} (B) 10 (C) 10^{10} (D) 29.5×10^{10}
20. The equilibrium constant for the reaction
 $\text{Sr(s)} + \text{Mg}^{2+}(\text{aq}) \rightleftharpoons \text{Sr}^{2+}(\text{aq}) + \text{Mg(s)}$ is 4×10^{12} at 25°C
 The E° for a cell made up of the Sr/Sr^{2+} and Mg^{2+}/Mg half cells
 ($\log 2 = 0.3$)
 (A) 0.3717 V (B) 0.7434 V (C) 0.1858 V (D) 0.135 V
21. For a reaction - $\text{A(s)} + 2\text{B}^+(\text{aq}) \rightarrow \text{A}^{2+}(\text{aq}) + 2\text{B(s)}$ K_c has been found to be 10^{12} . The E°_{cell} is:
 (A) 0.354 V (B) 0.708 V (C) 0.0098 V (D) 1.36 V
22. Determine the value of E°_{cell} for the following reaction -
 $\text{Cu}^{2+}(\text{aq}) + \text{Sn}^{2+}(\text{aq}) \longrightarrow \text{Cu(s)} + \text{Sn}^{4+}(\text{aq})$ Equilibrium constant is 10^6
 (A) 0.177 (B) 0.0177 (C) 0.215 (D) 1.77

Application Of Electrode Potential, Reduction And Oxidation Power

23. A standard reduction electrode potential of four metals are
 $A = -0.250\text{ V}$, $B = -0.140\text{ V}$, $C = -0.126\text{ V}$, $D = -0.402\text{ V}$
 The metal that displaces A from its aqueous solution is :-
 (A) B (B) C (C) D (D) None of the above
24. The standard reduction potentials at 25°C for the following half reactions are :
 $\text{Zn}^{2+}(\text{aq.}) + 2\text{e}^- \rightleftharpoons \text{Zn(s)}; -0.762$
 $\text{Cr}^{3+}(\text{aq.}) + 3\text{e}^- \rightleftharpoons \text{Cr(s)}; -0.740$
 $2\text{H}^+ + 2\text{e}^- \rightleftharpoons \text{H}_2(\text{g}); 0.00$
 $\text{Fe}^{3+} + \text{e}^- \rightleftharpoons \text{Fe}^{2+}; 0.77$
 Which is the strongest reducing agent ?
 (A) Zn (B) Cr (C) $\text{H}_2(\text{g})$ (D) $\text{Fe}^{2+}(\text{aq})$




25. The reduction potential values are given below:





Which one is the best reducing agent ?

- (A) Al (B) Mg (C) Cu (D) I₂

26. The following facts are available :-







Which of the following statements is correct :-



Cell Representation, Types of Half Cell,

NERNST EQUATION & ITS APPLICATIONS

27. For the cell $\text{Pt}(\text{Cl}_2) | \text{Cl}^-(\text{aq}) || \text{Ag}^+(\text{aq}) | \text{Ag}$, cell reaction is given by



28. For a reaction, $\text{M}^{n+1} + n e^- \longrightarrow \text{M}^{n+}$, the Nernst equation has a form of :

(A) $E = E^\circ + \frac{RT}{nF} \lambda_n \frac{[\text{M}^+]}{[\text{M}^{n+1}]}$

(B) $E = E^\circ - \frac{RT}{nF} \lambda_n \frac{[\text{M}^+]}{[\text{M}^{n+1}]}$

(C) $E = E^\circ - \frac{RT}{nF} \log \frac{[\text{M}^+]}{[\text{M}^{n+1}]}$

(D) $E = E^\circ + \frac{RT}{nF} \log \frac{[\text{M}^+]}{[\text{M}^{n+1}]}$

29. The reduction potential of hydrogen electrode ($P_{\text{H}_2} = 1 \text{ atm}$; $[\text{H}^+] = 0.1 \text{ M}$) at 25°C will be -

(A) 0.00 V

(B) -0.059 V

(C) 0.118 V

(D) 0.059 V

30. Which of the following represents the reduction potential of silver wire dipped into 0.1 M AgNO_3 solution at 25°C ?

(A) E°_{red}

(B) $(E^\circ_{\text{red}} + 0.059)$

(C) $(E^\circ_{\text{oxi}} - 0.059)$

(D) $(E^\circ_{\text{red}} - 0.059)$

31. By how much times will potential of half cell Cu^{+2}/Cu change if, the solution is diluted to 100 times at 298 K :-

(A) Increases by 59 mV

(B) Decrease by 59 mV

(C) Increases by 29.5 mV

(D) Decreases by 29.5 mV

32. How much will the reduction potential of a hydrogen electrode change when its solution initially at pH = 0 is neutralised to pH = 7 ?
 (A) increase by 0.059V (B) decrease by 0.059V
 (C) increase by 0.413V (D) decrease by 0.413V
33. If the pressure of H₂ gas is increased from 1 atm to 100 atm keeping H⁺ concentration constant at 1 M, the change in reduction potential of hydrogen half-cell at 25°C will be
 (A) 0.059 V (B) 0.59 V (C) 0.0295 V (D) 0.118 V

Nernst Equation .Full Reaction

34. Which of the following will increase the voltage of the cell with following cell reaction
 $\text{Sn(s)} + 2\text{Ag}^+(\text{aq}) \rightarrow \text{Sn}^{2+}(\text{aq}) + 2\text{Ag(s)}$
 (A) Decrease in the concentration of Ag⁺ ions
 (B) Increase in the concentration of Sn²⁺ ions
 (C) Increase in the concentration of Ag⁺ ions
 (D) (A) & (B) both
35. The emf of the cell
 $\text{Ti} / \text{Ti}^+ (0.01\text{M}) || \text{Cu}^{2+} (0.1\text{M}) / \text{Cu}$ is 0.83 V
 The emf of this cell will be increased by :
 (A) Increase the concentration of Cu²⁺ ions (B) Decreasing the concentration of Ti⁺
 (C) Increasing the concentration of both (D) (A) & (B) both
36. Consider the cell, $\text{Cu} | \text{Cu}^{2+} || \text{Ag}^+ | \text{Ag}$. If the concentration of Cu²⁺ and Ag⁺ ions becomes ten times then emf of the cell :-
 (A) Becomes 10 times (B) Remains same
 (C) Increase by 0.0295 V (D) Decrease by 0.0295 V
37. The emf of the cell in which the following reaction,
 $\text{Zn(s)} + \text{Ni}^{2+}(\text{aq}) (a = 0.1) \rightarrow \text{Zn}^{2+}(\text{aq}) (a = 1.0) + \text{Ni(s)}$
 occurs, is found to be 0.5105 V at 298 K. The standard e.m.f. of the cell is :-
 (A) -0.5105 V (B) 0.5400 V (C) 0.4810 V (D) 0.5696 V
38. The standard emf for the cell reaction,
 $\text{Zn(s)} + \text{Cu}^{2+}(\text{aq}) \longrightarrow \text{Zn}^{2+}(\text{aq}) + \text{Cu(s)}$ is 1.10 volt at 25 °C. The emf for the cell reaction when 0.1 M Cu²⁺ and 0.1 M Zn²⁺ solution are used at 25°C is :
 (A) 1.10 volt (B) 0.110 volt (C) -1.10 volt (D) -0.110 volt
39. Consider the reaction,
 $\text{Cl}_2(\text{g}) + 2\text{Br}^-(\text{aq}) \longrightarrow 2\text{Cl}^-(\text{aq}) + \text{Br}_2(\text{g})$
 The emf of the cell when [Cl⁻] = [Br⁻] = 0.01M and Cl₂ gas at 1 atm pressure while Br₂(g) at 0.01 atm will be (E° for the above reaction is = 0.29 volt) :
 (A) 0.54 volt (B) 0.35 volt (C) 0.24 volt (D) -0.29 volt

Concentration Cell

40. What is the potential of the cell containing two hydrogen electrodes as represented below
 $\text{Pt} | \text{H}_2(\text{g}) | \text{H}^+_{(\text{aq})}(10^{-8} \text{ M}) || \text{H}^+_{(\text{aq})}(0.001 \text{ M}) | \text{H}_2(\text{g}) | \text{Pt}$
 (A) -0.295 V (B) -0.0591 V (C) 0.295 V (D) 0.0591 V
41. Which represent a concentration cell?
 (A) $\text{Pt} | \text{H}_2 | \text{HCl} || \text{HCl} | \text{PtH}_2$ (B) $\text{Pt} | \text{H}_2 | \text{HCl} || \text{Cl}_2 | \text{Pt}$
 (C) $\text{Zn} | \text{Zn}^{2+} || \text{Cu}^{2+} | \text{Cu}$ (D) $\text{Fe} | \text{Fe}^{2+} || \text{Cu}^{2+} | \text{Cu}$
42. $\text{Zn} | \text{Zn}^{2+} (\text{C}_1) || \text{Zn}^{2+} (\text{C}_2) | \text{Zn}$. for this cell ΔG is negative if -
 (A) $\text{C}_1 = \text{C}_2$ (B) $\text{C}_1 > \text{C}_2$ (C) $\text{C}_2 > \text{C}_1$ (D) None

Metal Sparingly Soluble Salt

Metal, Metal Insoluble Salt- Ion Electrode


43. The chemical reaction
 $2\text{AgCl}(\text{s}) + \text{H}_2(\text{g}) \longrightarrow 2\text{HCl}(\text{aq}) + 2\text{Ag}(\text{s})$
 taking place in a galvanic cell is represented by the notation
 (A) $\text{Pt}(\text{s}) | \text{H}_2(\text{g}), 1 \text{ bar} | 1 \text{ M KCl}(\text{aq}) | \text{AgCl}(\text{s}) | \text{Ag}(\text{s})$
 (B) $\text{Pt}(\text{s}) | \text{H}_2(\text{g}), 1 \text{ bar} | 1 \text{ M HCl}(\text{aq}) | 1 \text{ M Ag}^+(\text{aq}) | \text{Ag}(\text{s})$
 (C) $\text{Pt}(\text{s}) | \text{H}_2(\text{g}), 1 \text{ bar} | 1 \text{ M HCl}(\text{aq}) | \text{AgCl}(\text{s}) | \text{Ag}(\text{s})$
 (D) $\text{Pt}(\text{s}) | \text{H}_2(\text{g}), 1 \text{ bar} | 1 \text{ M HCl}(\text{aq}) | \text{Ag}(\text{s}) | \text{AgCl}(\text{s})$
44. A silver wire dipped in 0.1 M HCl solution saturated with AgCl develops oxidation potential of -0.209 V . If $E^0_{\text{Ag}/\text{Ag}^+} = -0.799 \text{ V}$, the K_{sp} of AgCl in pure water will be
 (A) 3×10^{-11} (B) 10^{-11} (C) 4×10^{-11} (D) 3×10^{-11}
45. The solubility product of silver iodide is 8.3×10^{-17} and the standard reduction potential of Ag , Ag^+ electrode is $+0.8 \text{ volts}$ at 25°C . The standard reduction potential of AgAgI/I^- electrode from these data is
 (A) -0.30 V (B) $+0.15 \text{ V}$ (C) $+0.10 \text{ V}$ (D) -0.15 V

Metal Sparingly Soluble Salt

THERMODYNAMIC FUNCTIONS OF CELL

46. $\Delta G = \Delta H - T\Delta S$ and $\Delta G = \Delta H + T \left[\frac{d(\Delta G)}{dT} \right]_p$ then $\left(\frac{dE_{\text{cell}}}{dT} \right)$ is :
 (A) $\frac{\Delta S}{nF}$ (B) $\frac{nE}{\Delta S}$ (C) $-nFE_{\text{cell}}$ (D) $+nFE_{\text{cell}}$
47. The maximum efficiency of cell is given by-
 (A) $\frac{\Delta H}{\Delta G}$ (B) $\frac{nFE_{\text{cell}}}{\Delta G}$ (C) $-\frac{nFE_{\text{cell}}}{\Delta H}$ (D) Zero

48. The efficiency of a hypothetical cell is about 84% which involves the following reaction :



Then, the standard electrode potential of the cell will be (Assume as $\Delta S = 0$)

- (A) 1.20 (B) 2.40 V (C) 1.10 V (D) 1.24 V
49. The temperature coefficient, of the emf i.e. $\frac{dE}{dT} = -0.00065 \text{ volt. deg}^{-1}$ for the cell $\text{Cd} | \text{CdCl}_2 (1\text{M}) || \text{AgCl}(s) | \text{Ag}$ at 25°C . Calculate the entropy changes $\Delta S_{298\text{K}}$ for the cell reaction, $\text{Cd} + 2\text{AgCl} \rightarrow \text{Cd}^{2+} + 2\text{Cl}^- + 2\text{Ag}$
- (A) -105.5 JK^{-1} (B) -150.2 JK^{-1} (C) -75.7 JK^{-1} (D) -125.5 JK^{-1}

Salt Bridge

50. KCl can be used in salt bridge if electrolyte of a galvanic cell contains:
- (A) Ag^+ - ions (B) Pb^{2+} - ions (C) Hg_2^{2+} -ions (D) Au^{3+} - ions
51. KCl can be used in salt bridge as electrolyte in which of the following cells?
- (A) $\text{Zn} | \text{ZnCl}_2 || \text{AgNO}_3 | \text{Ag}$ (B) $\text{Pb} | \text{Pb}(\text{NO}_3)_2 || \text{Cu}(\text{NO}_3)_2 | \text{Cu}$
 (C) $\text{Cu} | \text{CuSO}_4 || \text{AuCl}_3 | \text{Au}$ (D) $\text{Fe} | \text{FeSO}_4 || \text{Pb}(\text{NO}_3)_2 | \text{Pb}$

Electrolytic Cell Calculations

52. When an electric current is passed through a cell containing an electrolyte, positive ions move towards the cathode and negative ions towards the anode. What will happen if the cathode is pulled out of the solution?
- (A) The positive ions will start moving towards the anode and negative ions will stop moving.
 (B) The negative ions will continue to move towards the anode and the positive ions will stop moving
 (C) Both positive and negative ions will move towards the anode.
 (D) None of these movements will take place.
53. When one coulomb of electricity is passed through an electrolytic solution the mass deposited on the electrode is equal to :
- (A) equivalent weight (B) molecular weight
 (C) electrochemical equivalent (D) one gram
54. When the same electric current is passed through the solution of different electrolytes in series the amounts of elements deposited on the electrodes are in the ratio of their:
- (A) atomic number (B) atomic masses (C) specific gravities (D) equivalent masses
55. The electric charge for electro deposition of 1 equivalent of a substance is :
- (A) one ampere for one second (B) 1 coulomb
 (C) one ampere for one hour (D) charge on one mole of electrons

56. Electro chemical equivalent of a substance is 0.0006; its equivalent weight is :
(A) 57.9 (B) 28.95 (C) 115.8 (D) cannot be calculated
57. A current of 9.65 amp. flowing for 10 minutes deposits 3.0 g of a metal. The equivalent wt. of the metal is :
(A) 10 (B) 30 (C) 50 (D) 96.5
58. The amount of electricity that can deposit 108 g. of silver from silver nitrate solution is:
(A) 1 ampere (B) 1 coulomb (C) 1 Faraday (D) 2 ampere
59. W g of copper deposited in a copper voltameter when an electric current of 2 ampere is passed for 2 hours. If one ampere of electric current is passed for 4 hours in the same voltameter, copper deposited will be :
(A) W (B) W/2 (C) W/4 (D) 2W
60. A current of 9.65 amp. passing for 16 min. 40 sec. through a molten tin salt deposits 5.95 gm of Tin. The oxidation state of the tin in the salt is : (at. wt of Sn = 119)
(A) +4 (B) +3 (C) +2 (D) +1
61. One mole of electron passes through each of the solution of AgNO_3 , CuSO_4 and AlCl_3 when Ag, Cu and Al are deposited at cathode. The molar ratio of Ag, Cu and Al deposited are
(A) 1 : 1 : 1 (B) 6 : 3 : 2 (C) 6 : 3 : 1 (D) 1 : 3 : 6
62. The ratio of weights of hydrogen and magnesium deposited by the same amount of electricity from aqueous H_2SO_4 and fused MgSO_4 are :
(A) 1 : 8 (B) 1 : 12 (C) 1 : 16 (D) None of these
63. The same amount of electricity was passed through two separate electrolytic cells containing solutions of nickel nitrate $[\text{Ni}(\text{NO}_3)_2]$ and chromium nitrate $[\text{Cr}(\text{NO}_3)_3]$ respectively. If 0.3 g of nickel was deposited in the first cell, the amount of chromium deposited is :
(at. wt. of Ni = 59, at. wt. of Cr = 52)
(A) 0.1 g (B) 0.17 g (C) 0.3 g (D) 0.6 g
64. When an electric current is passed through acid diluted water, 112 ml. of hydrogen gas at 1 atm and 273 K collects at the cathode in 965 second. The current passed, in ampere is :
(A) 1.0 (B) 0.5 (C) 0.1 (D) 2.0
65. How many coulombs of electric charge are required for the oxidation of 1 mole of H_2O to O_2 ?
(A) $9.65 \times 10^4 \text{ C}$ (B) $4.825 \times 10^5 \text{ C}$ (C) $1.93 \times 10^5 \text{ C}$ (D) $1.93 \times 10^4 \text{ C}$
66. An electric current is passed through silver voltameter connected to a water voltameter in series. The cathode of the silver voltameter weighed 0.108g more at the end of the electrolysis. The volume of oxygen evolved at STP is :
(A) 56 cm^3 (B) 550 cm^3 (C) 5.675 cm^3 (D) 11.2 cm^3

67. The charge required for the oxidation of one mole Mn_3O_4 into MnO_4^{2-} in presence of alkaline medium is
(A) $5 \times 96500 \text{ C}$ (B) 96500 C (C) $10 \times 96500 \text{ C}$ (D) $2 \times 96500 \text{ C}$
68. The time required to coat a surface of 80 cm^2 with $5 \times 10^{-3} \text{ cm}$ thick layer of silver (density 1.08 g cm^{-3}) with the passage of 9.65 A current through a silver nitrate solution is :
(A) 10 sec. (B) 40 sec. (C) 30 sec. (D) 20 sec.

Order Of Deposition

69. The products formed when an aqueous solution of NaBr is electrolyzed in a cell having inert electrodes are :
(A) Na and Br_2 (B) Na and O_2 (C) H_2 , Br_2 and NaOH (D) H_2 and O_2
70. Electrolysis of a CuSO_4 produces :-
(A) An increase in pH (B) A decrease in pH
(C) H_2 will produce at anode (D) H_2 will produce at cathode
71. A solution of sodium sulphate in water is electrolysed using inert electrodes. The products at the cathode and anode are respectively.
(A) H_2 , O_2 (B) O_2 , H_2 (C) O_2 , Na (D) none
72. When an aqueous solution of lithium chloride is electrolysed using graphite electrodes
(A) Cl_2 is liberated at the anode.
(B) Li is deposited at the cathode
(C) as the current flows, pH of the solution remains constant
(D) as the current flows, pH of the solution decreases.

Cells and batteries

Commercial Cells & Corrosion

73. A fuel cell uses $\text{CH}_4(\text{g})$ and forms CO_3^{2-} at the anode. It is used to power a car with 80 Amp. for 0.96 hr. How many litres of $\text{CH}_4(\text{g})$ (at 1 atm and 273 K) would be required ?($F = 96500$). Assume 100% efficiency.
(A) 5 (B) 6 (C) 7 (D) 8
74. When lead storage battery is charged :
(A) lead dioxide dissolves
(B) sulphuric acid is regenerated
(C) the lead electrode becomes coated with lead sulphate
(D) the amount of sulphuric acid in solution decreases
75. Which is not correct method for prevention of iron from Rusting -
(A) Galvanisation (B) Connecting to sacrificial electrode of Mg
(C) Making medium alkaline (D) Making medium acidic

Conductance ,Conductivity, Molar Conductance

76. Electrolytic conduction differs from metallic conduction from the fact that in the former
 (A) The resistance increases with increasing temperature
 (B) The resistance decreases with increasing temperature
 (C) The resistance remains constant with increasing temperature
 (D) The resistance is independent of the length of the conductor
77. Which of the following solution of KCl has the lowest value of specific conductance :
 (A) 1 M (B) 0.1 M (C) 0.01 M (D) 0.001 M
78. Which of the following solutions of KCl has the lowest value of molar conductance ?
 (A) 1 M (B) 0.1 M (C) .01 M (D) .001 M
79. The resistance of 0.5 M solution of an electrolyte in a cell was found to be 50 Ω . If the electrodes in the cell are 2.2 cm apart and have an area of 4.4 cm² then the molar conductivity (in S m² mol⁻¹) of the solution is
 (A) 0.2 (B) 0.02 (C) 0.002 (D) None of these
80. If 0.01 M solution of an electrolyte has a resistance of 40 ohms in a cell having a cell constant of 0.4 cm⁻¹ then its molar conductance in ohm⁻¹ cm² mol⁻¹ is :
 (A) 10 (B) 10² (C) 10³ (D) 10⁴
81. The resistance of 0.01 N solution of an electrolyte was found to be 200 ohm at 298 K using a conductivity cell of cell constant 1.5 cm⁻¹. The equivalent conductance of solution is :-
 (A) 750 mho cm² eq⁻¹ (B) 75 mho cm² eq⁻¹
 (C) 750 mho⁻¹ cm² eq⁻¹ (D) 75 mho⁻¹cm² eq⁻¹
82. Equal volumes of 0.015 M CH₃COOH & 0.015 M NaOH are mixed together. What would be molar conductivity of mixture if conductivity of CH₃COONa is 6.3 $\times 10^{-4}$ S cm⁻¹
 (A) 8.4 S cm² mol⁻¹ (B) 84 S cm² mol⁻¹ (C) 4.2 S cm² mol⁻¹ (D) 42 S cm² mol⁻¹

Kohlrausch Law, Kohlrausch And Alpha

83. The molar conductance at infinite dilution of AgNO₃, AgCl and NaCl are 115, 120 and 110 respectively. The molar conductance of NaNO₃ is :-
 (A) 110 (B) 105 (C) 130 (D) 150
84. Equivalent conductance of Ba²⁺ and Cl⁻ ions are 127 & 76 ohm⁻¹ cm² eq⁻¹ respectively. Equivalent conductance of BaCl₂ at infinite dilution is -
 (A) 139.5 (B) 101.5 (C) 203 (D) 279
85. The equivalent conductivity of 0.1 N CH₃COOH at 25 °C is 80 and at infinite dilution 400. The degree of dissociation of CH₃COOH is :
 (A) 1 (B) 0.2 (C) 0.1 (D) 0.5

86. Equivalent conductance of 0.1 M HA(weak acid) solution is $10 \text{ Scm}^2\text{equivalent}^{-1}$ and that at infinite dilution is $200 \text{ S cm}^2 \text{eq}^{-1}$ Hence pH of HA solution is
(A) 1.3 (B) 1.7 (C) 2.3 (D) 3.7
87. The dissociation constant of n-butyric acid is $1.6 \times 10^{-5} \text{ M}$ and the molar conductivity at infinite dilution is $380 \times 10^{-4} \text{ S m}^2 \text{mol}^{-1}$. The specific conductance of the 0.01 M acid solution is
(A) $1.52 \times 10^{-5} \text{ S m}^{-1}$ (B) $1.52 \times 10^{-2} \text{ S m}^{-1}$
(C) $1.52 \times 10^{-3} \text{ S m}^{-1}$ (D) None

K_{sp} And Kohlrausch

88. The conductivity of a saturated solution of BaSO_4 is $3.06 \times 10^{-6} \text{ ohm}^{-1} \text{ cm}^{-1}$ and its molar conductance is $1.53 \text{ ohm}^{-1} \text{ cm}^2 \text{mol}^{-1}$. The K_{sp} of BaSO_4 will be :
(A) 4×10^{-12} (B) 2.5×10^{-9} (C) 2.5×10^{-13} (D) 4×10^{-6}
89. The conductivity of a saturated solution of Ag_3PO_4 is $9 \times 10^{-6} \text{ S m}^{-1}$ and its equivalent conductivity is $1.50 \times 10^{-4} \text{ S m}^2 \text{equivalent}^{-1}$. The K_{sp} of Ag_3PO_4 is
(A) 4.32×10^{-18} (B) 1.8×10^{-9} (C) 8.64×10^{-13} (D) None of these

EXERCISE # S-I

Electrode Potential, Galvanic Cell

Galvanic Cell, Its Representation & Salt Bridge

1. In the galvanic cell $\text{Cu} | \text{Cu}^{2+} || \text{Ag}^+ | \text{Ag}$, the electrons flow from Cu-electrode to Ag-electrode. Answer the following questions regarding this cell :
- Which is the anode ?
 - Which is the cathode ?
 - What happens at anode-reduction or oxidation ?
 - What happens at cathode-oxidation or reduction ?
 - Which electrode loses mass ?
 - Which electrode gains mass ?
 - Write the electrode reactions.
 - Write the cell reaction
 - Which metal has greater tendency to loss electron-Cu or Ag ?
 - Which is the more reactive metal-Cu or Ag ?
 - What is the function of salt bridge represented by the symbol $||$?


$$E_1 + E_2 = E_3$$

STANDARD CELL POTENTIAL AND ELECTROCHEMICAL SERIES

2. What is E°_{Cell} if :





$$E^\circ = -0.74 \text{ V}$$



$$E^\circ = 0.94 \text{ V}$$

3. Determine the standard reduction potential for the half reaction :



$$E^\circ_{\text{Cell}} = -0.15 \text{ V}$$




$$E^\circ = 1.20 \text{ V}$$


 ΔG And Electrode Potential

4. If $E^\circ_{\text{Fe}^{2+}|\text{Fe}} = -0.44 \text{ V}$, $E^\circ_{\text{Fe}^{3+}|\text{Fe}^{2+}} = 0.77 \text{ V}$. Calculate $E^\circ_{\text{Fe}^{3+}|\text{Fe}}$.

5. If for the half cell reactions




$$E^\circ = 0.15 \text{ V}$$




$$E^\circ = 0.34 \text{ V}$$

Calculate E° of the half-cell reaction




6. Consider the standard reduction potentials (in volts) as shown in figure. Find E° .




7. If for the half cell reactions
- $$\text{Cu}^{2+} + e^{-} \longrightarrow \text{Cu}^{+} \quad E^{\circ} = 0.15 \text{ V}$$
- $$\text{Cu}^{+} + e^{-} \longrightarrow \text{Cu} \quad E^{\circ} = 0.53 \text{ V}$$

predict whether Cu^{+} undergoes disproportionation or not.

8. The standard oxidation potentials for Mn^{2+} ion acid solution are



Is the reaction



spontaneous under conditions of unit activity? What is the change in free energy?

Electrode Potential And Equilibrium Constant

9. Calculate the equilibrium constant for the reaction
 $\text{Fe}^{2+} + \text{Ce}^{4+} \rightleftharpoons \text{Fe}^{3+} + \text{Ce}^{3+}$,
 [given : $E^{\circ}_{\text{Ce}^{4+}/\text{Ce}^{3+}} = 1.44\text{V}$; $E^{\circ}_{\text{Fe}^{3+}/\text{Fe}^{2+}} = 0.68\text{V}$]
10. For a cell $\text{Mg(s)} \mid \text{Mg}^{2+}(\text{aq}) \parallel \text{Ag}^{+}(\text{aq}) \mid \text{Ag}$,
 (i) Calculate the equilibrium constant at 25°C .
 (ii) Also find the maximum work per mole Mg that can be obtained by operating the cell.
 $E^{\circ}(\text{Mg}^{2+}/\text{Mg}) = -2.37\text{V}$, $E^{\circ}(\text{Ag}^{+}/\text{Ag}) = 0.8 \text{ V}$.
11. The standard oxidation potential of Zn referred to SHE is 0.76V and that of Cu is -0.34V at 25°C .
 When excess of Zn is added to CuSO_4 , Zn displaces Cu^{2+} till equilibrium is reached. What is the approximate value of $\log \frac{[\text{Zn}^{2+}]}{[\text{Cu}^{2+}]}$ at equilibrium?
12. The standard reduction potential at 25°C for the reduction of water
 $2\text{H}_2\text{O} + 2e^{-} \rightleftharpoons \text{H}_2 + 2\text{OH}^{-}$ is -0.84 volt .
 Calculate the equilibrium constant for the reaction
 $\text{H}_2\text{O} \rightleftharpoons \text{H}^{+} + \text{OH}^{-}$ at 25°C .
 (Take $\frac{2.303RT}{F} = 0.06$)
13. Calculate the equilibrium constant for the reaction:
 $3\text{Sn(s)} + 2\text{Cr}_2\text{O}_7^{2-} + 28\text{H}^{+} \longrightarrow 3\text{Sn}^{4+} + 4\text{Cr}^{3+} + 14\text{H}_2\text{O}$
 E° for $\text{Sn}/\text{Sn}^{2+} = 0.136 \text{ V}$
 E° for $\text{Sn}^{2+}/\text{Sn}^{4+} = -0.154 \text{ V}$
 E° for $\text{Cr}_2\text{O}_7^{2-}/\text{Cr}^{3+} = 1.33 \text{ V}$
14. For the reaction,
 $4\text{Al(s)} + 3\text{O}_2(\text{g}) + 6\text{H}_2\text{O} + 4\text{OH}^{-} \rightleftharpoons 4[\text{Al}(\text{OH})_4^{-}]$; $E^{\circ}_{\text{cell}} = 2.73 \text{ V}$.
 If $\Delta G^{\circ}_f(\text{OH}^{-}) = -157 \text{ kJ mol}^{-1}$ and $\Delta G^{\circ}_f(\text{H}_2\text{O}) = -237.2 \text{ kJ mol}^{-1}$,
 Determine $\Delta G^{\circ}_f[\text{Al}(\text{OH})_4^{-}]$.

Application Of Electrode Potential, Reduction And Oxidation Power

15. The reduction potential values are given below
 $\text{Al}^{3+}/\text{Al} = -1.67 \text{ volt}$, $\text{Mg}^{2+}/\text{Mg} = -2.34 \text{ volt}$, $\text{Cu}^{2+}/\text{Cu} = +0.34 \text{ volt}$
 $\text{I}_2 / \text{I}^- = +0.53 \text{ volt}$. Which one is the best reducing agent ?
16. The standard reduction potential value of the three metallic cations X, Y and Z are 0.52, - 3.03 and - 1.18 V respectively. Write the decreasing order of reducing power of the corresponding metals :
17. The value of E° for electrode reactions,
 $\text{Fe} \longrightarrow \text{Fe}^{2+} + 2\text{e}^- \quad E^\circ = 0.44 \text{ volt}$
 $\text{Cu} \longrightarrow \text{Cu}^{2+} + 2\text{e}^- \quad E^\circ = -0.33 \text{ volt}$
 $\text{Zn} \longrightarrow \text{Zn}^{2+} + 2\text{e}^- \quad E^\circ = 0.76 \text{ volt}$
 State which of these metals can replace the other two from the solution of their salts ?
18. Is 1.0 M H^+ solution under H_2SO_4 at 1.0 atm capable of oxidising silver metal in the presence of 1.0 M Ag^+ ion?

$$E^\circ_{\text{Ag}^+/\text{Ag}} = 0.80 \text{ V},$$

$$E^\circ_{\text{H}^+/\text{H}_2(\text{Pt})} = 0.0 \text{ V}$$

Cell Representation, Types Of Half Cell


19. Write cell reaction of the following cells :
 (a) $\text{Ag} | \text{Ag}^+ (\text{aq}) || \text{Cu}^{2+} (\text{aq}) | \text{Cu}$ (b) $\text{Pt} | \text{Fe}^{2+}, \text{Fe}^{3+} || \text{MnO}_4^-, \text{Mn}^{2+}, \text{H}^+ | \text{Pt}$
 (c) $\text{Pt}, \text{Cl}_2 | \text{Cl}^- (\text{aq}) || \text{Ag}^+ (\text{aq}) | \text{Ag}$ (d) $\text{Pt}, \text{H}_2 | \text{H}^+ (\text{aq}) || \text{Cd}^{2+} (\text{aq}) | \text{Cd}$
20. Write cell representation for following cells.
 (a) $\text{Cd}^{2+} (\text{aq}) + \text{Zn} (\text{s}) \longrightarrow \text{Zn}^{2+} (\text{aq}) + \text{Cd} (\text{s})$
 (b) $2\text{Ag}^+ (\text{aq}) + \text{H}_2 (\text{g}) \longrightarrow 2\text{H}^+ (\text{aq}) + 2\text{Ag} (\text{s})$
 (c) $\text{Cr}_2\text{O}_7^{2-} (\text{aq}) + 14\text{H}^+ (\text{aq}) + 6\text{Fe}^{2+} (\text{aq}) \longrightarrow 6\text{Fe}^{3+} (\text{aq}) + 2\text{Cr}^{3+} (\text{aq}) + 7\text{H}_2\text{O} (\text{l})$

NERNST EQUATION

Nernst equation, Half reaction

21. Calculate the oxidation potential of a hydrogen electrode at $\text{pH} = 1$ ($T = 298 \text{ K}$).
22. At what concentration of Cu^{2+} in a solution of CuSO_4 will the electrode potential be zero at 25°C ? Given : $E^\circ (\text{Cu} | \text{Cu}^{2+}) = -0.34 \text{ V}$.
23. A zinc electrode is placed in a 0.1M solution at 25°C . Assuming that the salt is 20% dissociated at this dilution calculate the electrode reduction potential. $E^\circ (\text{Zn}^{2+} | \text{Zn}) = -0.76 \text{ V}$.
24. The standard reduction potential of $\text{Cu}^{2+} / \text{Cu}$ couple is 0.34 V at 25°C . Calculate the reduction potential at $\text{pH} = 14$ for this couple.
 (Given : $K_{\text{sp}}, \text{Cu} (\text{OH})_2 = 1.0 \times 10^{-19}$).

25. Calculate the EMF of a Daniel cell when the concentration of ZnSO_4 and CuSO_4 are 0.001 M and 0.1M respectively. The standard potential of the cell is 1.1V.
26. Calculate E^0 and E for the cell $\text{Sn} | \text{Sn}^{2+} (1\text{M}) || \text{Pb}^{2+} (10^{-3}\text{M}) | \text{Pb}$,
 $E^0 (\text{Sn}^{2+} | \text{Sn}) = -0.14\text{V}$, $E^0 (\text{Pb}^{2+} | \text{Pb}) = -0.13\text{V}$. Is cell representation is correct?
27. Consider the following electrochemical cell :



- Write a balanced net ionic equation for the spontaneous reaction that take place in the cell.
- Calculate the standard cell potential E° for the cell reaction.
- If the cell emf is 1.6 V, what is the concentration of Zn^{2+} ?
- How will the cell potential be affected if KI is added to Ag^+ half-cell ?

28. Calculate the EMF of the following cell

$$\text{Zn} \mid \text{Zn}^{2+} (0.01\text{M}) \parallel \text{Zn}^{2+} (0.1 \text{ M}) \mid \text{Zn}$$

at 298 K.

- 29.** Calculate pH using the following cell :

$$\text{Pt (H}_2\text{)} \mid \text{H}^+ (\text{x M}) \parallel \text{H}^+ (1 \text{ M}) \mid \text{Pt (H}_2\text{)} \quad \text{if } E_{\text{cell}} = 0.2364 \text{ V.}$$


$$1 \text{ atm} \qquad \qquad \qquad 1 \text{ atm}$$

- 30.** Equinormal Solutions of two weak acids, HA ($pK_a = 3$) and HB ($pK_a = 5$) are each placed in contact with equal pressure of hydrogen electrode at 25°C to form two hydrogen half-cell. Find the emf of a cell constructed by interconnecting them.
- 31.** In two vessels each containing 500ml water, 0.5 mmol of aniline ($K_b = 10^{-9}$) and 50 mmol of HCl are added separately. Two hydrogen electrodes are constructed using these solutions. Calculate the emf of cell made by connecting them appropriately.

Metal Sparingly Soluble Salt


Metal- Metal Insoluble Salt- Ion Electrode

32. Write cell reaction from given cell diagrams :



33. For the galvanic cell : $\text{Ag} | \text{AgBr}(\text{s}) | \text{KBr} (0.001\text{M}) || \text{KCl} (0.2\text{M}) | \text{AgCl}(\text{s}) | \text{Ag}$,

Calculate the EMF generated? (Take $\frac{2.303RT}{F} = 0.06$)




34. The emf of the cell $\text{Ag} | \text{AgI}(\text{s}) | \text{KI}(0.05\text{M}) || \text{AgNO}_3(0.05\text{M}) | \text{Ag}$ is 0.788V. Calculate the solubility product of AgI.

35. Consider the cell $\text{Ag} | \text{AgBr}(\text{s}) | \text{Br}^- || \text{Cl}^- | \text{AgCl}(\text{s}) | \text{Ag}$ at 25°C . The solubility product constants of AgBr & AgCl are respectively 5×10^{-13} & 1×10^{-10} . For what ratio of the concentrations of Br^- & Cl^- ions would the emf of the cell be zero ?

36. Given, $E^\circ = -0.27 \text{ V}$ for the $\text{Cl}^- | \text{PbCl}_2 | \text{Pb}$ couple and -0.12 V for the $\text{Pb}^{2+} | \text{Pb}$ couple, determine


K_{sp} for PbCl_2 at 25°C ? (Take $\frac{2.303RT}{F} = 0.06$)

37. The pK_{sp} of AgI is 16. if the E° value for $\text{Ag}^+ | \text{Ag}$ is 0.8 V. Find the E° for the half cell reaction





Electrolytic Cell Calculations

38. Calculate the no. of electrons lost or gained during electrolysis of



39. How many faradays of electricity are required to convert



40. 0.5 mole of electron is passed through two electrolytic cells in series. One contains silver ions, and the other zinc ions. Assume that only cathode reaction in each cell is the reduction of the ion to the metal. How many gm of each metal will be deposited.

41. 3A current was passed through an aqueous solution of an unknown salt of Pd for 1Hr. 2.977g of Pd^{+n} was deposited at cathode. Find n. (Given Atomic mass of Pd = 106.4)

42. If 0.224 litre of H_2 gas is formed at the cathode, how much O_2 gas is formed at the anode under identical conditions?
43. Assume 96500 C as one unit of electricity. If cost of electricity of producing x gm Al is Rs x, what is the cost of electricity of producing x gm Mg?
44. Chromium metal can be plated out from an acidic solution containing CrO_3 according to following equation:
 $CrO_3(aq) + 6H^+(aq) + 6e^- \rightarrow Cr(s) + 3H_2O$
 Calculate :
 (i) How many grams of chromium will be plated out by 24125 coulombs and
 (ii) How long will it take to plate out 1.5 gm of chromium by using 12.5 ampere current
45. A current of 3 amp was passed for 2 hours through a solution of $CuSO_4$. 3 g of Cu^{2+} ions were deposited as Cu at cathode. Calculate percentage current efficiency of the process.
46. The electrosynthesis of MnO_2 is carried out from a solution of $MnSO_4$ in H_2SO_4 (aq). If a current of 25.5 ampere is used with a current efficiency of 85%, how long would it take to produce 1 kg of MnO_2 ?
47. Electrolysis of a solution of HSO_4^- ions produce $S_2O_8^{2-}$. Assuming 75% current efficiency, what current should be employed to achieve a production rate of 1 mole of $S_2O_8^{2-}$ per hour?

Order Of Deposition

48.

	ELECTROLYTE	Product at ANODE	Product at CATHODE
1	NaCl (Molten) with Pt electrode		
2	NaCl (aq) with Pt electrode		
3	Na_2SO_4 (aq) with Pt electrode		
4	$NaNO_3$ (aq) with Pt electrode		
5	$AgNO_3$ (aq) with Pt electrode		
6	$CuSO_4$ (aq) with Inert electrode		
7	$CuSO_4$ (aq) with Copper electrode		

49. A constant current of 30 A is passed through an aqueous solution of NaCl for a time of 1.0 hr. How many moles of NaOH are produced? What is volume of Cl_2 gas at 1atm, 273K produced (in litre) ?
50. After electrolysis of 600 ml NaCl solution with inert electrodes for a certain period of time, was found to be 1N in NaOH. During the same time, 31.75 g of Cu was deposited in the copper voltameter in series with the electrolytic cell. Calculate the percentage yield of NaOH obtained considering 100% yield in copper voltameter.
51. A solution of $Ni(NO_3)_2$ is electrolysed between platinum electrodes using a current of 5 ampere for 20 minutes. What mass of Ni is deposited at the cathode?
52. A current of 3.7A is passed for 6hrs. between Ni electrodes in 0.5L of 2M solution of $Ni(NO_3)_2$. What will be the molarity of solution at the end of electrolysis?

Molar Conductance

53. When a solution of specific conductance $1.342 \text{ ohm}^{-1} \text{ m}^{-1}$ was placed in a conductivity cell with parallel electrodes, the resistance was found to be 170.5 ohm . Area of electrodes is $1.86 \times 10^{-4} \text{ m}^2$. Calculate separation of electrodes.
54. The resistance of a conductivity cell filled with 0.01 M solution of NaCl is 200 ohm at 18°C . Calculate the molar conductivity of the solution. The cell constant of the conductivity cell is 0.88 cm^{-1} .
55. A solution containing 2.08 g of anhydrous barium chloride is 500 CC of water has a specific conductance $0.005 \text{ ohm}^{-1} \text{ cm}^{-1}$. What are molar and equivalent conductivities of this solution.
56. For 0.01 N KCl , the resistivity 800 ohm cm . Calculate the conductivity and equivalent conductance.
57. The molar conductivity of $0.1 \text{ M CH}_3\text{COOH}$ solution is $4 \text{ S cm}^2 \text{ mole}^{-1}$. What is the conductivity and resistivity of the solution ?
58. The conductivity of pure water in a conductivity cell with electrodes of cross sectional area 4 cm^2 and 2 cm apart is $8 \times 10^{-7} \text{ S cm}^{-1}$.
(i) What is resistance of conductivity cell ?
(ii) What current would flow through the cell under an applied potential difference of 1 volt ?

Kohlrausch Law, Kohlrausch and Alpha

59. For the strong electrolytes NaOH , NaCl and BaCl_2 the molar ionic conductivities at infinite dilution are 240×10^{-4} , 125×10^{-4} and $280.0 \times 10^{-4} \text{ mho cm}^2 \text{ mol}^{-1}$ respectively. Calculate the molar conductivity of Ba(OH)_2 at infinite dilution.
60. The value of Λ_m^∞ for HCl , NaCl and $\text{CH}_3\text{CO}_2\text{Na}$ are 425 , 125 and $100 \text{ S cm}^2 \text{ mol}^{-1}$ respectively. Calculate the value of Λ_m^∞ for acetic acid. If the molar conductivity of a given acetic acid solution is $48 \text{ S cm}^2 \text{ mol}^{-1}$ at 25°C , calculate its degree of dissociation.
61. Equivalent conductance of $0.01 \text{ N Na}_2\text{SO}_4$ solution is $120 \text{ ohm}^{-1} \text{ cm}^2 \text{ eq}^{-1}$. The equivalent conductance at infinite dilution is $150 \text{ ohm}^{-1} \text{ cm}^2 \text{ eq}^{-1}$. What is the degree of dissociation in $0.01 \text{ N Na}_2\text{SO}_4$ solution?
62. Hydrofluoric acid is weak acid. At 25°C , the molar conductivity of 0.002 M HF is $200 \text{ ohm}^{-1} \text{ cm}^2 \text{ mole}^{-1}$. If its $\Lambda_m^\infty = 400 \text{ ohm}^{-1} \text{ cm}^2 \text{ mole}^{-1}$, calculate its degree of dissociation and equilibrium constant at the given concentration.

Ksp and Kohlrausch


63. Saturated solution of AgCl at 25°C has specific conductance of $1.12 \times 10^{-6} \text{ ohm}^{-1} \text{ cm}^{-1}$. The λ_∞ Ag^+ and $\lambda_\infty \text{Cl}^-$ are 54 and $58 \text{ ohm}^{-1} \text{ cm}^2 \text{ eq}^{-1}$. respectively. Calculate the solubility product of AgCl at 25°C .
64. Specific conductance of a saturated solution of AgBr is $8.75 \times 10^{-7} \text{ ohm}^{-1} \text{ cm}^{-1}$ at 25°C . Specific conductance of pure water at 25°C is $0.75 \times 10^{-7} \text{ ohm}^{-1} \text{ cm}^{-2}$. Λ_m^∞ for KBr , AgNO_3 and KNO_3 are 140 , 130 , $110 \text{ (S cm}^2 \text{ mol}^{-1})$ respectively. Calculate the solubility of AgBr in gm/litre .
65. At 25°C , $\lambda_\infty(\text{H}^+) = 3.5 \times 10^{-2} \text{ S m}^2 \text{ mol}^{-1}$ and $\lambda_\infty(\text{OH}^-) = 2 \times 10^{-2} \text{ S m}^2 \text{ mol}^{-1}$.
Given: Sp. conductance = $5.5 \times 10^{-6} \text{ S m}^{-1}$ for H_2O , determine pH and K_w .

EXERCISE # (O-II)

Single correct

1. The standard emf of the cell, $\text{Cd(s)} \mid \text{CdCl}_2(\text{aq}) (0.1 \text{ M}) \parallel \text{AgCl(s)} \mid \text{Ag(s)}$ in which the cell reaction is,
 $\text{Cd(s)} + 2\text{AgCl(s)} \longrightarrow 2\text{Ag(s)} + \text{Cd}^{2+}(\text{aq}) + 2\text{Cl}^{-}(\text{aq})$ is 0.6915 V at 0°C and 0.6753 V at 25°C .
 The ΔH° of the reaction at 25°C is :
 (A) - 176 kJ (B) - 234.7 kJ (C) + 123.5 kJ (D) - 167.6 kJ





2.



	Content	Resistance	Volume of gas produced (at same T & P)
Cell-1	$\text{CuSO}_4(\text{aq})$	10Ω	at anode is V_1
Cell-2	$\text{NaCl}(\text{aq})$	2Ω	at anode is V_2
Cell-3	$\text{ZnSO}_4(\text{aq})$	6Ω	at cathode is V_3




The ratio $V_1 : V_2 : V_3$ is

- (A) 2 : 3 : 1 (B) 3 : 2 : 1 (C) 3 : 1 : 2 (D) 2 : 1 : 3
3. For saturated solution of CaF_2 , choose the correct relationship-
- (A) $\Lambda_{\text{mCaF}_2}^\infty = \lambda_{\text{eqCa}^{2+}}^\infty + 2\lambda_{\text{eqF}^\ominus}^\infty$ (B) $\Lambda_{\text{mCaF}_2}^\infty = 2(\lambda_{\text{eqCa}^{2+}}^\infty + \lambda_{\text{eqF}^\ominus}^\infty)$
 (C) $\Lambda_{\text{mCaF}_2}^\infty = 2(\lambda_{\text{eqCa}^{2+}}^\infty + \lambda_{\text{eqF}^\ominus}^\infty)$ (D) $\Lambda_{\text{mCaF}_2}^\infty = \frac{1}{2}(\lambda_{\text{eqCa}^{2+}}^\infty + \lambda_{\text{eqF}^\ominus}^\infty)$
4. Which of the following is the most appropriate graph for conductometric titration when $\text{AgNO}_3(\text{aq.})$ solution is titrated against $\text{KCl}(\text{aq.})$ solution. (Assuming that these ions do not differ much in ionic conductance)



Multiple correct:





5. The value of E° for elements are as follows :



Based on the above data correct statements are -


- (A) Strongest reducing agent is Cr^{3+} (B) Strongest oxidising agent is Fe^{3+}
 (C) Sn^{4+} can oxidise I^- (D) Fe^{3+} can oxidise I^-

6. Pick out the correct statements among the following from inspection of standard reduction potentials (Assume standard state conditions).



- (A) Cl_2 can oxidise SO_4^{2-} from solution
 (B) Cl_2 can oxidise Br^- and I^- from aqueous solution
 (C) I_2 can oxidise Cl^- and Br^- from aqueous solution
 (D) $\text{S}_2\text{O}_8^{2-}$ can oxidise Cl^- , Br^- and I^- from aqueous solution

7. Using the standard electrode potential values given below, decide which of the statements, I, II, III and IV are correct. Choose the right answer from (A), (B), (C) and (D).



- I. Copper can displace iron from FeSO_4 solution.
 II. Iron can displace copper from CuSO_4 solution.
 III. Silver can displace copper from CuSO_4 solution.
 IV. Iron can displace silver from AgNO_3 solution.

- (A) I and II (B) II and III (C) II and IV (D) I and IV

8. The EMF of the following cell is 0.22 volt.
 $\text{Ag(s)} \mid \text{AgCl(s)} \mid \text{KCl (1M)} \mid \text{H}^+ \text{(1M)} \mid \text{H}_2 \text{(g) (1atm)} \mid \text{Pt(s)}$.
 Which of the following will decrease the EMF of cell.
 (A) increasing pressure of $\text{H}_2 \text{(g)}$ from 1 atm to 2 atm
 (B) increasing Cl^- concentration in Anodic compartment
 (C) increasing H^+ concentration in cathodic compartment
 (D) decreasing KCl concentration in Anodic compartment.
9. Select correct statements for the cell
 $\text{Ag} \mid \text{AgCl} \mid \text{HCl} \parallel \text{AgNO}_3 \mid \text{Ag}$; $(\Delta S)_{\text{reaction}} = -ve$
 (A) $E_{\text{Cell}}^\circ > 0$
 (B) $\Delta G^\circ < 0$
 (C) E_{cell} decrease by increasing temperature
 (D) E_{cell} decrease by diluting anodic solution
10. Which of the following arrangement will produce oxygen at anode during electrolysis ?
 (A) Dilute H_2SO_4 solution with Cu electrodes.
 (B) Dilute H_2SO_4 solution with inert electrodes.
 (C) Fused NaOH with inert electrodes.
 (D) Dilute NaCl solution with inert electrodes.
11. If 270.0 g of water is electrolysed during an experiment performed by miss Abhilasha with 75% current efficiency then
 (A) 168 L of $\text{O}_2 \text{(g)}$ will be evolved at anode at 1 atm & 273 K
 (B) Total 504 L gases will be produced at 1 atm & 273 K.
 (C) 336 L of $\text{H}_2 \text{(g)}$ will be evolved at anode at 1 atm & 273 K
 (D) 45 F electricity will be consumed
12. During discharging of lead storage battery, which of the following is/are true ?
 (A) H_2SO_4 is produced (B) H_2O is consumed
 (C) PbSO_4 is formed at both electrodes (D) Density of electrolytic solution decreases
13. Select the INCORRECT options (s) for electrolytic solution -
 (A) Molar conductivity and equivalent conductivity both increases on increasing dilution for strong electrolyte as well as weak electrolyte solution
 (B) Conductance and conductivity both increases on increasing dilution for strong electrolyte as well as weak electrolyte solution
 (C) On increasing temperature conductivity & conductance increases.
 (D) If conductivity of solution decreases then molar conductivity also decreases.

14. Select correct statement at infinite dilution for an electrolyte -
- (A) Λ_m is maximum (B) Resistance is zero
- (C) Λ_m° is constant for any temperature (D) No interaction remain among the ions

Paragraph for question nos. 6 to 8

A half cell is prepared by $K_2Cr_2O_7$ in a buffer solution of pH = 1. Concentration of $K_2Cr_2O_7$ is 1 M. To 3 litre of this solution 570 gm of $SnCl_2$ is added which is oxidised completely to $SnCl_4$.

Given : $E^\circ_{Cr_2O_7^{2-}/Cr^{+3}, H^+} = 1.33 \text{ V}$; $\frac{2.303}{F} RT = 0.06$;

Atomic mass of Sn = 119 ; $E^\circ_{Sn^{+4}/Sn^{+2}} = 0.15$

15. Number of moles of Cr^{+3} formed are
- (A) 2 (B) 6 (C) 4 (D) 3
16. Half cell potential $E_{Cr_2O_7^{2-}/Cr^{+3}}$ after the reaction of $SnCl_2$ is
- (A) 1.187 (B) 1.191 (C) 1.285 (D) None of these
17. Emf of the cell $Pt | Sn^{+2}, Sn^{+4}, H^+ || Cr_2O_7^{2-}, Cr^{+3}, H^+ | Pt$
- (0.1M) (0.2 M) (1M), (0.2M) (1M) (1M)
- (A) -1.18 V (B) 1.164 V (C) 1.18 V (D) None of these

Paragraph for Q.18 & Q.19



The cell $Pt (H_2) (1 \text{ atm}) | H^+(pH = ?) || I^- (a = 1) | AgI(s), Ag(s) | Pt$ has emf, $E_{298K} = 0$. The standard electrode potential for the reaction $AgI + e^- \rightarrow Ag + I^\ominus$ is - 0.151 volt.

18. Calculate the pH value.
- (A) 3.37 (B) 5.26 (C) 2.56 (D) 4.62
19. Calculate the solubility product of AgI in water at $25^\circ C$ [$E^\circ_{(Ag^+, Ag)} = + 0.799 \text{ volt}$]
- (A) 1.97×10^{-17} (B) 8.43×10^{-17} (C) 1.79×10^{-17} (D) 9.17×10^{-17}


Paragraph for Question 20 to 22

H_2O_2 can be produced by the Ammonium hydrogen sulphate.

Reactions occurring in electrolytic cell.



Hydrolysis of Ammonium persulphate



20. What is the current efficiency when 100 Amp current is passed for 965 sec. in order to produce 17 gm of H_2O_2 .
 (A) 20 % (B) 40 % (C) 80 % (D) 100 %
21. What volume of hydrogen gas at 1 atm & 273 K will be produced at cathode in above question.
 (A) 22.4 L (B) 11.2 L (C) 5.6 L (D) 2.8 L
22. How many moles of electrons are to be passed in order to produce enough H_2O_2 which when reacted with excess of KI then liberated iodine required 100 ml of centimolar Hypo solution.
 (A) 10^{-1} (B) 10^{-2} (C) 10^{-3} (D) 5×10^{-4}


Paragraph for Q.23 & Q.24

For the titration of a solution of HCl against NaOH solution, hydrogen electrode is dipped into HCl solution. Then this electrode is connected to normal calomel electrode (NCE).

In such experiment 25 ml of HCl (0.1M) is titrated against a standard 0.1 M NaOH solution

$$E_{\text{NCE(reduction)}} = 0.24\text{V}; \frac{2.303RT}{F} = 0.06$$

23. Select the correct graph for titration



24. E_{cell} after adding 24.95 ml of NaOH
 (A) 0.3 (B) 0 V (C) 0.48 V (D) 0.64 V

Assertion & Reasoning type questions :

25. Statement -1 : The voltage of mercury cell remains constant for long period of time.
 Statement -2 : It is because net cell reaction does not involve active species.
 (A) Statement-1 is true, statement-2 is true and statement-2 is correct explanation for statement-1.
 (B) Statement-1 is true, statement-2 is true and statement-2 is NOT the correct explanation for statement-1.
 (C) Statement-1 is true, statement-2 is false.
 (D) Statement-1 is false, statement-2 is true.

26. Statement-1 : The SRP of three metallic ions A^+, B^{2+}, C^{3+} are $-0.3, -0.5, 0.8$ volt respectively, so oxidising power of ions is $C^{3+} > A^+ > B^{2+}$.

Statement-2 : Higher the SRP, higher the oxidising power.

(A) Statement-1 is true, statement-2 is true and statement-2 is correct explanation for statement-1.

(B) Statement-1 is true, statement-2 is true and statement-2 is NOT the correct explanation for statement-1.

(C) Statement-1 is true, statement-2 is false.

(D) Statement-1 is false, statement-2 is true.

Match the column:

27. **Column I**

- (A) Dilute solution of HCl
- (B) Dilute solution of NaCl
- (C) Concentrated solution of NaCl
- (D) $AgNO_3$ solution

Column II

(Electrolysis product using inert electrode)

- (P) O_2 evolved at anode
- (Q) H_2 evolved at cathode
- (R) Cl_2 evolved at anode
- (S) Ag deposition at cathode

28. **List-I**

- (P) Conductivity (k) increases
- (Q) Molar conductance (λ_m) increases
- (R) Conductance (G)
- (S) Molar conductance λ_m^∞ at infinite

List-II


- (1) Depends on cell constant as well as concentration
- (2) Independent of concentration as well as cell constant
- (3) With decreases in concentration but independent of cell constant
- (4) With increase in concentration but independent dilution of cell constant

Code:

	P	Q	R	S
(A)	4	1	3	2
(B)	4	3	2	1
(C)	3	4	1	2
(D)	4	3	1	2

EXERCISE # (S-II)

1. From the standard potentials shown in the following diagram, calculate the potentials E_1^0 and E_2^0 .



2. The EMF of the cell $M | M^{n+} (0.02M) || H^+ (1M) | H_2(g) (1 \text{ atm}), Pt$ at $25^\circ C$ is $0.81V$. Calculate the valency of the metal if the standard oxidation of the metal is $0.76V$
3. Calculate the EMF of the cell,
 $Zn - Hg(c_1M) | Zn^{2+} (aq) | Hg - Zn(c_2M)$
 at $25^\circ C$, if the concentrations of the zinc amalgam are: $c_1 = 10g$ per $100g$ of mercury and $c_2 = 1g$ per $100g$ of mercury.
4. Calculate the emf of the cell
 $Pt, H_2(1.0 \text{ atm}) | CH_3COOH (0.1M) || NH_3(aq, 0.01M) | H_2 (1.0 \text{ atm}), Pt$
 $K_a(CH_3COOH) = 1.8 \times 10^{-5}, K_b(NH_3) = 1.8 \times 10^{-5}$.
5. The Edison storage cell is represented as $Fe(s) | FeO(s) | KOH(aq) | Ni_2O_3(s) | Ni(s)$ The half-cell reaction are
 $Ni_2O_3(s) + H_2O(l) + 2e^- \rightleftharpoons 2NiO(s) + 2OH^-, E^0 = +0.40V$
 $FeO(s) + H_2O(l) + 2e^- \rightleftharpoons Fe(s) + 2OH^-, E^0 = -0.87V$
 (i) What is the cell reaction?
 (ii) What is the cell emf? How does it depend on the concentration of KOH ?
 (iii) What is the maximum amount of electrical energy that can be obtained from one mole of Ni_2O_3 ?
6. For the galvanic cell : $Ag | AgCl(s) | KCl (0.2M) || KBr (0.001 M) | AgBr(s) | Ag$,
 Calculate the EMF generated and assign correct polarity to each electrode for a spontaneous process after taking into account the cell reaction at $25^\circ C$.
 $[K_{sp}(AgCl) = 2.8 \times 10^{-19}; K_{sp}(AgBr) = 3.3 \times 10^{-13}]$
7. Calculate the quantity of electricity that would be required to reduce $12.3g$ of nitrobenzene to aniline, if the current efficiency for the process is 50% . If the potential drop across the cell is 3.0 volts, how much energy will be consumed?
8. How long a current of $2A$ has to be passed through a solution of $AgNO_3$ to coat a metal surface of $80cm^2$ with 5μ thick layer? Density of silver = $10.8g/cm^3$.

9. 10g solution of CuSO_4 is electrolyzed using 0.01F of electricity. Calculate:
(a) The weight of resulting solution (b) Equivalents of acid or alkali in the solution.
10. Cadmium amalgam is prepared by electrolysis of a solution of CdCl_2 using a mercury cathode. How long should a current of 5A be passed in order to prepare 12% Cd-Hg amalgam on a cathode of 2gm Hg ($\text{Cd} = 112.4$)
11. One of the methods of preparation of per disulphuric acid, $\text{H}_2\text{S}_2\text{O}_8$, involve electrolytic oxidation of H_2SO_4 at anode ($2\text{H}_2\text{SO}_4 \rightarrow \text{H}_2\text{S}_2\text{O}_8 + 2\text{H}^+ + 2\text{e}^-$) with oxygen and hydrogen as by-products. In such an electrolysis, 9.722 L of H_2 and 2.35 L of O_2 were generated at STP. What is the weight of $\text{H}_2\text{S}_2\text{O}_8$ formed?
12. Dal lake has water 8.2×10^{12} litre approximately. A power reactor produces electricity at the rate of 1.5×10^6 coulomb per second at an appropriate voltage. How many years would it take to electrolyse the lake?
13. The equivalent conductance of 0.10 N solution of MgCl_2 is $97.1 \text{ mho cm}^2 \text{ eq}^{-1}$ at 25°C . a cell with electrode that are 1.5 cm^2 in surface area and 0.5 cm apart is filled with 0.1 N MgCl_2 solution. How much current will flow when potential difference between the electrodes is 5 volt.
14. The specific conductance at 25°C of a saturated solution of SrSO_4 is $1.482 \times 10^{-4} \text{ ohm}^{-1} \text{ cm}^{-1}$ while that of water used is $1.5 \times 10^{-6} \text{ mho cm}^{-1}$. Determine at 25°C the solubility in gm per litre of SrSO_4 in water. Molar ionic conductance of Sr^{2+} and SO_4^{2-} ions at infinite dilution are 59.46 and $79.8 \text{ ohm}^{-1} \text{ cm}^2 \text{ mole}^{-1}$ respectively. [$\text{Sr} = 87.6$, $\text{S} = 32$, $\text{O} = 16$]

EXERCISE # (JEE-MAINS)

1. The molar conductivity of a conductivity cell filled with 10 mole of 20 mL NaCl solution is Λ_{m1} and that of 20 moles another identical cell having 80 mL NaCl solution is Λ_{m2} , The conductivities exhibited by these two cells are same.

The relationship between Λ_{m2} and Λ_{m1} is

- (A) $\Lambda_{m2} = 2\Lambda_{m1}$ (B) $\Lambda_{m2} = \Lambda_{m1} / 2$ (C) $\Lambda_{m2} = 2\Lambda_{m1}$ (D) $\Lambda_{m2} = 4\Lambda_{m1}$

[JEE Main, July 2022]

2. The limiting molar conductivities of NaI, NaNO₃ and AgNO₃ are 12.7, 12.0 and 13.3 mS m² mol⁻¹, respectively (all at 25°C). The limiting molar conductivity of AgI at this temperature is ____ mS m² mol⁻¹

[JEE Main, June 2022]

3. A dilute solution of Sulphuric acid is electrolysed using a current of 0.10 A for 2 hours to produce hydrogen and oxygen gas. The total volume of gases produced at STP is ____ cm³. (Nearest integer) [Given : Faraday constant $F = 96500 \text{ C mol}^{-1}$ at STP, molar volume of an ideal gas is 22.7 L mol⁻¹]

[JEE Main, June 2022]

4. Given below are two statements:

Statement I: The limiting molar conductivity of KCl (strong electrolyte) is higher compared to that of CH₃COOH (weak electrolyte).


Statement II: Molar conductivity decreases with decrease in concentration of electrolyte.

In the light of the above statements, choose the most appropriate answer from the options given below:

[JEE Main, August 2021]

- (A) Statement I is true but Statement II is false.
 (B) Statement I is false but Statement II is true.
 (C) Both Statement I and Statement II are true.
 (D) Both Statement I and Statement II are false.

5. For the reaction



the magnitude of the standard molar free energy change, $\Delta_r G_m^\circ = -$ ____ kJ.

(Round off to the Nearest Integer)


$$\left[\begin{array}{l} E_{\text{Fe}^{2+}/\text{Fe}(\text{s})}^0 = -0.440\text{V}; \quad E_{\text{Fe}^{3+}/\text{Fe}(\text{s})}^0 = -0.036\text{V} \\ E_{\text{I}_2/2\text{I}^{-}}^0 = 0.539\text{V}; \quad F = 96500\text{C} \end{array} \right]$$

(Physical Chemistry)

ELECTROCHEMISTRY

6. Consider the cell at 25°C [JEE Main, March 2021]
 $\text{Zn} | \text{Zn}^{2+} (\text{aq}), (1\text{M}) || \text{Fe}^{3+} (\text{aq}), \text{Fe}^{2+} (\text{aq}) | \text{Pt} (\text{s})$
 The fraction of total iron present as Fe^{3+} ion at the cell potential of 1.500 V is $x \times 10^{-2}$. The value of x is _____. (Nearest integer)
 (Given: $E^\ominus_{\text{Fe}^{3+}/\text{Fe}^{2+}} = 0.77\text{V}$, $E^\ominus_{\text{Zn}^{2+}/\text{Zn}} = -0.76\text{V}$) [JEE Main, July 2021]
7. The conductivity of a weak acid HA of concentration 0.001 mol L^{-1} is $2.0 \times 10^{-5} \text{ S cm}^{-1}$. If $\Lambda^\ominus_m(\text{HA}) = 190 \text{ S cm}^2 \text{ mol}^{-1}$, the ionization constant (K_a) of HA is equal to $___ \times 10^{-6}$.
 (Round off to the nearest integer) [JEE Main, July 2021]
8. The resistance of a conductivity cell with cell constant 1.14 cm^{-1} , containing 0.001 M KCl at 298 K is 1500Ω . The molar conductivity of 0.001 M KCl solution at 298 K in $\text{S cm}^2 \text{ mol}^{-1}$ is _____.
 (Integer answer) [JEE Main, August 2021]
9. 108g of silver (molar mass 108 g mol^{-1}) is deposited at cathode from $\text{AgNO}_3(\text{aq})$ solution by a certain quantity of electricity. The volume (in L) of oxygen gas produced at 273 K and 1 bar pressure from water by the same quantity of electricity is _____. [Jee Main, 2020]
10. An acidic solution of dichromate is electrolyzed for 8 minutes using 2A current. As per the following equation
 $\text{Cr}_2\text{O}_7^{2-} + 14\text{H}^+ + 6\text{e}^- \rightarrow 2\text{Cr}^{3+} + 7\text{H}_2\text{O}$
 The amount of Cr^{3+} obtained was 0.104 g. The efficiency of the process (in %) is
 (Take : $F = 96000 \text{ C}$, At. mass of chromium = 52) _____.
 Given 60.31 [Jee Main, 2020]
11. Given that the standard potentials (E^\ominus) of Cu^{2+}/Cu and Cu^+/Cu are 0.34 V and 0.522 V respectively, the E^\ominus of $\text{Cu}^{2+}/\text{Cu}^+$ is : [Jee Main, 2020]
 (A) -0.182 V (B) -0.158 V (C) 0.182 V (D) +0.158 V
12. For the given cell ;
 $\text{Cu}(\text{s}) | \text{Cu}^{2+} (\text{C}_1\text{M}) || \text{Cu}^{2+} (\text{C}_2\text{M}) | \text{Cu}(\text{s})$ change in Gibbs energy (ΔG) is negative, if : [Jee Main, 2020]
 (A) $\text{C}_1 = 2\text{C}_2$ (B) $\text{C}_2 = \frac{\text{C}_1}{\sqrt{2}}$ (C) $\text{C}_1 = \text{C}_2$ (D) $\text{C}_2 = \sqrt{2}\text{C}_1$
13. Given that $E^\ominus_{\text{O}_2/\text{H}_2\text{O}} = 1.23 \text{ V}$; [JEE Mains 2019 (Apr.)]
 $E^\ominus_{\text{S}_2\text{O}_8^{2-}/\text{SO}_4^{2-}} = 2.05 \text{ V}$
 $E^\ominus_{\text{Br}_2/\text{Br}^-} = +1.09 \text{ V}$
 $E^\ominus_{\text{Au}^{3+}/\text{Au}} = +1.4 \text{ V}$
 The strongest oxidizing agent is :
 (A) O_2 (B) $\text{S}_2\text{O}_8^{2-}$ (C) Br_2 (D) Au^{3+}

14. Calculate the standard cell potential (in V) of the cell in which following reaction takes place :



[JEE Mains 2019 (Apr.)]





Given that

$$E^\circ_{\text{Ag}^+/\text{Ag}} = x \text{ V}$$





$$E^\circ_{\text{Fe}^{2+}/\text{Fe}} = y \text{ V}$$

$$E^\circ_{\text{Fe}^{3+}/\text{Fe}} = z \text{ V}$$

- (A) $x - z$ (B) $x + 2y - 3z$ (C) $x - y$ (D) $x + y - z$
15. The standard Gibbs energy for the given cell reaction in kJ mol^{-1} at 298 K is :
 $\text{Zn}(\text{s}) + \text{Cu}^{2+}(\text{aq}) \rightarrow \text{Zn}^{2+}(\text{aq}) + \text{Cu}(\text{s})$, $E^\circ = 2 \text{ V}$ at 298 K
 (Faraday's constant, $F = 96000 \text{ C mol}^{-1}$) [JEE Mains 2019 (Apr.)]
 (A) -384 (B) 384 (C) -192 (D) 192
16. A solution of $\text{Ni}(\text{NO}_3)_2$ is electrolysed between platinum electrodes using 0.1 Faraday electricity. How many mole of Ni will be deposited at the cathode? [JEE Mains 2019 (Apr.)]
 (A) 0.20 (B) 0.15 (C) 0.10 (D) 0.05
17. Consider the statements S1 and S2 :
 S1: Conductivity always increases with decrease in the concentration of electrolyte.
 S2: Molar conductivity always increases with decrease in the concentration of electrolyte.
 The correct option among the following is: [JEE Mains 2019 (Apr.)]
 (A) Both S1 and S2 are wrong (B) Both S1 and S2 are correct
 (C) S1 is wrong and S2 is correct (D) S1 is correct and S2 is wrong
18. Which one of the following graphs between molar conductivity (Λ_m) versus \sqrt{C} is correct ? [JEE Mains 2019 (Apr.)]



19. Given: [JEE Mains 2019 (Apr.)]



Oxidizing power of the species will increase in the order:

- (A) $\text{Ce}^{4+} < \text{Pb}^{4+} < \text{Bi}^{3+} < \text{Co}^{3+}$ (B) $\text{Co}^{3+} < \text{Pb}^{4+} < \text{Ce}^{4+} < \text{Bi}^{3+}$
 (C) $\text{Bi}^{3+} < \text{Ce}^{4+} < \text{Pb}^{4+} < \text{Co}^{3+}$ (D) $\text{Co}^{3+} < \text{Ce}^{4+} < \text{Bi}^{3+} < \text{Pb}^{4+}$

(Physical Chemistry)

ELECTROCHEMISTRY

20. The decreasing order of electrical conductivity of the following aqueous solutions is:
 0.1 M Formic acid (A),
 0.1 M Acetic acid (B),
 0.1 M Benzoic acid (C).
 (A) $A > B > C$ (B) $C > A > B$ (C) $A > C > B$ (D) $C > B > A$ [JEE Mains 2019 (Apr.)]
21. The anodic half-cell of lead-acid battery is recharged using electricity of 0.05 Faraday. The amount of PbSO_4 electrolyzed in g during the process is : (Molar mass of $\text{PbSO}_4 = 303 \text{ g mol}^{-1}$)
 (A) 22.8 (B) 15.2 (C) 7.6 (D) 11.4 [JEE Mains 2019 (Jan.)]
22. If the standard electrode potential for a cell is 2 V at 300 K, the equilibrium constant (K) for the reaction
 $\text{Zn(s)} + \text{Cu}^{2+}(\text{aq}) \rightleftharpoons \text{Zn}^{2+}(\text{aq}) + \text{Cu(s)}$ at 300 K is approximately.
 (R = $8 \text{ J K}^{-1} \text{ mol}^{-1}$, F = 96000 C mol^{-1})
 (A) e^{-80} (B) e^{-160} (C) e^{160} (D) e^{320} [JEE Mains 2019 (Jan.)]
23. Consider the following reduction processes :
 $\text{Zn}^{2+} + 2e^- \rightarrow \text{Zn(s)}$; $E^\circ = -0.76 \text{ V}$
 $\text{Ca}^{2+} + 2e^- \rightarrow \text{Ca(s)}$; $E^\circ = -2.87 \text{ V}$
 $\text{Mg}^{2+} + 2e^- \rightarrow \text{Mg(s)}$; $E^\circ = -2.36 \text{ V}$
 $\text{Ni}^{2+} + 2e^- \rightarrow \text{Ni(s)}$; $E^\circ = -0.25 \text{ V}$
 The reducing power of the metals increases in the order :
 (A) $\text{Ca} < \text{Mg} < \text{Zn} < \text{Ni}$ (B) $\text{Zn} < \text{Mg} < \text{Ni} < \text{Ca}$
 (C) $\text{Ni} < \text{Zn} < \text{Mg} < \text{Ca}$ (D) $\text{Ca} < \text{Zn} < \text{Mg} < \text{Ni}$ [JEE Mains 2019 (Jan.)]
24. In the cell $\text{Pt(s)} | \text{H}_2(\text{g}, 1\text{bar}) | \text{HCl(aq)} | \text{AgCl(s)} | \text{Ag(s)} | \text{Pt(s)}$ the cell potential is 0.92 V when a 10^{-6} molal HCl solution is used. The standard electrode potential of (AgCl/Ag, Cl^-) electrode is :
 $\left\{ \text{Given, } \frac{2.303RT}{F} = 0.06 \text{ V at } 298 \text{ K} \right\}$
 (A) 0.94 V (B) 0.40 V (C) 0.20 V (D) 0.76 V [JEE Mains 2019 (Jan.)]
25. For the cell $\text{Zn(s)} | \text{Zn}^{2+}(\text{aq}) || \text{M}^{x+}(\text{aq}) | \text{M(s)}$, different half cells and their standard electrode potentials are given below : [JEE Mains 2019 (Jan.)]
- | | | | | |
|--|--|---|---|--|
| $\text{M}^{x+}(\text{aq}) / \text{M(s)}$ | $\text{Au}^{3+}(\text{aq}) / \text{Au(s)}$ | $\text{Ag}^+(\text{aq}) / \text{Ag(s)}$ | $\text{Fe}^{3+}(\text{aq}) / \text{Fe}^{2+}(\text{aq})$ | $\text{Fe}^{2+}(\text{aq}) / \text{Fe(s)}$ |
| $E^\circ_{\text{M}^{x+}/\text{M}^{(v)}}$ | 1.40 | 0.80 | 0.77 | -0.44 |
- If $E^\circ_{\text{Zn}^{2+}/\text{Zn}} = -0.76 \text{ V}$, which cathode will give a maximum value of E°_{cell} per electron transferred?
 (A) $\text{Fe}^{2+} / \text{Fe}$ (B) $\text{Ag}^{3+} / \text{Au}$ (C) $\text{Fe}^{3+} / \text{Fe}^{2+}$ (D) Ag^+ / Ag

(Physical Chemistry)

ELECTROCHEMISTRY

26. Given the equilibrium constant : [JEE Mains 2019 (Jan.)]
 K_c of the reaction :
 $\text{Cu(s)} + 2\text{Ag}^+(\text{aq}) \rightarrow \text{Cu}^{2+}(\text{aq}) + 2\text{Ag(s)}$ is
 10×10^{15} , calculate the E_{cell}^0 of this reaction at 298 K
 $[2.303 \frac{RT}{F} \text{ at } 298\text{K is } 0.059\text{V}]$
 (A) 0.4736 V (B) 0.04736 mV (C) 0.4736 mV (D) 0.04736 V
27. The standard electrode potential E^\ominus and its temperature coefficient $\left(\frac{dE^\ominus}{dT}\right)$ for a cell are 2V and $-5 \times 10^{-4} \text{ VK}^{-1}$ at 300 K respectively. The cell reaction is $\text{Zn(s)} + \text{Cu}^{2+}(\text{aq}) \rightarrow \text{Zn}^{2+}(\text{aq}) + \text{Cu(s)}$ The standard reaction enthalpy $(\Delta_r H^\ominus)$ at 300 K in kJ mol^{-1} is,
 [Use $R = 8 \text{ JK}^{-1} \text{ mol}^{-1}$ and $F = 96,000 \text{ C mol}^{-1}$] [JEE Mains 2019 (Jan.)]
 (A) 192.0 (B) -384.0 (C) 206.4 (D) -412.8
28. \wedge_m° for NaCl, HCl and NaA are 126.4, 425.9 and $100.5 \text{ S cm}^2 \text{ mol}^{-1}$, respectively. If the conductivity of 0.001 M HA is $5 \times 10^{-5} \text{ S cm}^{-1}$, degree of dissociation of HA is:
[JEE Mains 2019 (Jan.)]
 (A) 0.50 (B) 0.125 (C) 0.25 (D) 0.75
29. How long (approximate) should water be electrolysed by passing through 100 amperes current so that the oxygen released can completely burn 27.66 g of diborane ?
 (Atomic weight of B = 10.8 u) [JEE Mains 2018]
 (A) 1.6 hours (B) 6.4 hours (C) 0.8 hours (D) 3.2 hours
30. Given
 $E_{\text{Cr}^{2+}/\text{Cr}^{3+}}^0 = 1.36 \text{ V}$, $E_{\text{Cr}^{3+}/\text{Cr}}^0 = -0.74 \text{ V}$
 $E_{\text{Cl}_2/\text{Cl}^-}^0 = 1.33 \text{ V}$, $E_{\text{MnO}_4^-/\text{Mn}^{2+}}^0 = 1.51 \text{ V}$
 Among the following, the strongest reducing agent is [JEE Mains 2017]
 (A) Cr (B) Mn^{2+} (C) Cr^{3+} (D) Cl^-
31. Oxidation of succinate ion produces ethylene and carbon dioxide gases. On passing 0.2 Faraday electricity through on aqueous solution of potassium succinate, the total volume of gases (at both cathode and anode) at STP (1 atm and 273 K) is : [JEE-MAINS (ONLINE) 2016]
 (A) 8.96 L (B) 2.24 L (C) 4.48 L (D) 6.72 L

32. What will occur if a block of copper metal is dropped into a beaker containing a solution of 1M ZnSO_4 [JEE-MAINS (ONLINE) 2016]
 (A) The copper metal will dissolve and zinc metal will be deposited
 (B) No reaction will occur
 (C) The copper metal will dissolve with evolution of oxygen gas
 (D) The copper metal will dissolve with evolution of hydrogen gas
33. Two Faraday of electricity is passed through a solution of CuSO_4 . The mass of copper deposited at the cathode is : (at. mass of Cu = 63.5 amu) [JEE-MAINS 2015]
 (A) 2g (B) 127 g (C) 0 g (D) 63.5 g
34. A variable, opposite external potential (E_{ext}) is applied to the cell $\text{Zn}|\text{Zn}^{2+} (1 \text{ M}) || \text{Cu}^{2+} (1 \text{ M}) | \text{Cu}$, of potential 1.1 V. When $E_{\text{ext}} < 1.1 \text{ V}$ and $E_{\text{ext}} > 1.1 \text{ V}$, respectively electrons flow from : [JEE-MAINS (ONLINE) 2015]
 (A) anode to cathode in both cases (B) anode to cathode and cathode to anode
 (C) cathode to anode in both cases (D) cathode to anode and anode to cathode
35. At 298 K, the standard reduction potentials are 1.51 V for $\text{MnO}_4^-|\text{Mn}^{2+}$, 1.36 V for $\text{Cl}_2|\text{Cl}^-$, 1.07 V for $\text{Br}_2|\text{Br}^-$, and 0.54 V for $\text{I}_2|\text{I}^-$. At pH = 3, permanganate is expected to oxidize $\left(\frac{RT}{F} = 0.059 \text{ V}\right)$: [JEE-MAINS (ONLINE) 2015]
 (A) Cl^- and Br^- (B) Cl^- , Br^- and I^- (C) Br^- and I^- (D) I^- only
36. Resistance of 0.2 M solution of an electrolyte is 50 Ω . The specific conductance of the solution is 1.4 S m^{-1} . The resistance of 0.5 M solution of the same electrolyte is 280 Ω . The molar conductivity of 0.5 M solution of the electrolyte in $\text{S m}^2 \text{ mol}^{-1}$ is : [JEE-MAINS 2014]
 (A) 5×10^3 (B) 5×10^2 (C) 5×10^{-4} (D) 5×10^{-3}
37. The equivalent conductance of NaCl at concentration C and at infinite dilution are λ_c and λ_∞ , respectively. The correct relationship between λ_c and λ_∞ is given as : [JEE-MAINS 2014]
 (where the constant B is positive)
 (A) $\lambda_c = \lambda_\infty - (B) \sqrt{C}$ (B) $\lambda_c = \lambda_\infty + (B) \sqrt{C}$
 (C) $\lambda_c = \lambda_\infty + (B) C$ (D) $\lambda_c = \lambda_\infty - (B) C$
38. Given :
 $E_{\text{Cr}^{3+}/\text{Cr}}^0 = 0.74 \text{ V}$; $E_{\text{MnO}_4^-/\text{Mn}^{2+}}^0 = 1.51 \text{ V}$
 $E_{\text{Cr}_2\text{O}_7^{2-}/\text{Cr}^{3+}}^0 = 1.33 \text{ V}$; $E_{\text{Cl}_2/\text{Cl}^-}^0 = 1.36 \text{ V}$
 Based on the data given above, strongest oxidising agent will be :
 (A) Cl^- (B) Cr^{3+} (C) Mn^{2+} (D) MnO_4^-

(Physical Chemistry)

ELECTROCHEMISTRY

39. The standard reduction potentials for $\text{Zn}^{2+} / \text{Zn}$, $\text{Ni}^{2+} / \text{Ni}$ and $\text{Fe}^{2+} / \text{Fe}$ are -0.76 , -0.23 and -0.44 V respectively. The reaction $\text{X} + \text{Y}^{2+} \rightarrow \text{X}^{2+} + \text{Y}$ will be spontaneous when [AIEEE 2012]
 (A) $\text{X} = \text{Zn}$, $\text{Y} = \text{Ni}$ (B) $\text{X} = \text{Ni}$, $\text{Y} = \text{Fe}$ (C) $\text{X} = \text{Ni}$, $\text{Y} = \text{Zn}$ (D) $\text{X} = \text{Fe}$, $\text{Y} = \text{Zn}$
40. The reduction potential of hydrogen half-cell will be negative if :- [AIEEE 2011]
 (A) $p(\text{H}_2) = 2$ atm $[\text{H}^+] = 1.0$ M (B) $p(\text{H}_2) = 2$ atm and $[\text{H}^+] = 2.0$ M
 (C) $p(\text{H}_2) = 1$ atm and $[\text{H}^+] = 2.0$ M (D) $p(\text{H}_2) = 1$ atm and $[\text{H}^+] = 1.0$ M
41. Resistance of 0.2 M solution of an electrolyte is 50Ω . The specific conductance of the solution is 1.3 S m^{-1} . If resistance of the 0.4 M solution of the same electrolyte is 260Ω , its molar conductivity is :- [AIEEE 2011]
 (A) $6250 \text{ S m}^2 \text{ mol}^{-1}$ (B) $6.25 \times 10^{-4} \text{ S m}^2 \text{ mol}^{-1}$
 (C) $625 \times 10^{-4} \text{ S m}^2 \text{ mol}^{-1}$ (D) $62.5 \text{ S m}^2 \text{ mol}^{-1}$
42. The Gibbs energy for the decomposition of Al_2O_3 at 500°C is as follows : [AIEEE 2010]

$$\frac{2}{3} \text{Al}_2\text{O}_3 \longrightarrow \frac{4}{3} \text{Al} + \text{O}_2, \Delta_r G = +966 \text{ kJ mol}^{-1}$$

 The potential difference needed for electrolytic reduction of Al_2O_3 at 500°C is at least :-
 (A) 5.0 V (B) 4.5 V (C) 3.0 V (D) 2.5 V
43. Given : [AIEEE-2009]
 $E_{\text{Fe}^{3+}/\text{Fe}}^0 = -0.036\text{V}$, $E_{\text{Fe}^{2+}/\text{Fe}}^0 = -0.439\text{V}$. The value of standard electrode potential for the change.
 $\text{Fe}^{3+}(\text{aq}) + \text{e}^- \longrightarrow \text{Fe}^{2+}(\text{aq})$ will be :-
 (A) 0.770 V (B) -0.27 V (C) -0.072 V (D) 0.385 V
44. Given : $E_{\text{Cr}^{3+}/\text{Cr}}^0 = -0.72$ V, $E_{\text{Fe}^{2+}/\text{Fe}}^0 = -0.42$ V. The potential for the cell, [AIEEE 2008]
 $\text{Cr}(\text{s}) | \text{Cr}^{3+}(\text{aq}) (0.1 \text{ M}) || \text{Fe}^{2+}(\text{aq}) (0.01 \text{ M}) | \text{Fe}(\text{s})$ is
 (A) 0.26 V (B) 0.339 V (C) -0.339 V (D) -0.26 V
45. The cell $\text{Zn} | \text{Zn}^{2+}(\text{aq}) (1\text{M}) || \text{Cu}^{2+}(\text{aq}) (1\text{M}) | \text{Cu}$ ($E^\circ_{\text{cell}} = 1.10$ V) was allowed to be completely discharged at 298 K. The relative concentration of Zn^{2+} to Cu^{2+} , $\left\{ \frac{[\text{Zn}^{2+}]}{[\text{Cu}^{2+}]} \right\}$ is : [AIEEE 2007]
 (A) 9.65×10^4 (B) Antilog (24.08) (C) 37.3 (D) 1037.3
46. Given the data at 25°C ,
 $\text{Ag}(\text{s}) + \text{I}^-(\text{aq}) \rightarrow \text{AgI}(\text{s}) + \text{e}^-$, $E^\circ = 0.152\text{V}$
 $\text{Ag}(\text{s}) \rightarrow \text{Ag}^+(\text{aq}) + \text{e}^-$, $E^\circ = -0.800$ V
 What is the value of $\log K_{\text{sp}}$ for AgI ?
 (Where K_{sp} = solubility product) [AIEEE 2006]

$$\left(2.303 \frac{RT}{F} = 0.59\text{V} \right)$$

 (A) -8.12 (B) $+8.612$ (C) -37.83 (D) -16.13

47. Resistance of a conductivity cell filled with a solution of an electrolyte of concentration 0.1M is 100Ω. The conductivity of this solution is 1.29 Sm⁻¹. Resistance of the same cell when filled with 0.02M of the same solution is 520Ω. The molar conductivity of 0.02M solution of the electrolyte will be. [AIEEE 2006]
- (A) $124 \times 10^{-4} \text{ Sm}^2 \text{ mol}^{-1}$ (B) $1240 \times 10^{-4} \text{ Sm}^2 \text{ mol}^{-1}$
 (C) $1.24 \times 10^4 \text{ Sm}^2 \text{ mol}^{-1}$ (D) $12.4 \times 10^{-4} \text{ Sm}^2 \text{ mol}^{-1}$
48. The molar conductivities, Λ_{NaOAc}^0 and Λ_{HCl}^0 at infinite dilution in water at 25°C are 91.0 and 426.2 S cm²/mol respectively. To calculate Λ_{HOAc}^0 the additional value required is : [AIEEE 2006]
- (A) KCl (B) NaOH (C) NaCl (D) H₂O
49. For a spontaneous reaction the ΔG, equilibrium constant (K) and E_{Cell}^0 will be respectively [AIEEE 2005]
- (A) -ve, < 1, -ve (B) -ve, > 1, -ve (C) -ve, > 1, +ve (D) +ve, > 1, -ve
50. Aluminium oxide may be electrolysed at 1000°C to furnish aluminium metal (1 Faraday = 96500 Coulombs). The cathode reaction is [AIEEE 2005]
- $\text{Al}^{3+} + 3e^- \longrightarrow \text{Al}$
- To prepare 5.12 kg of aluminium metal by this method would require,
- (A) $5.49 \times 10^4 \text{ C}$ of electric charge (B) $5.49 \times 10^1 \text{ C}$ of electric charge
 (C) $5.49 \times 10^7 \text{ C}$ of electric charge (D) $1.83 \times 10^7 \text{ C}$ of electric charge
51. The highest electrical conductivity of the following aqueous solution is of [AIEEE 2005]
- (A) 0.1 M fluoroacetic acid (B) 0.1 M difluoroacetic acid
 (C) 0.1 M acetic acid (D) 0.1 M chloroacetic acid
52. Electrolyte $\Lambda^\infty (\text{S cm}^2 \text{ mol}^{-1})$
- | | |
|------------------|-------|
| KCl | 149.9 |
| KNO ₃ | 145.0 |
| HCl | 426.2 |
| NaOAc | 91.0 |
| NaCl | 126.5 |
- Calculate $\Lambda_{\text{HOAc}}^\infty$ using appropriate molar conductances of the electrolytes listed above at infinite dilution in H₂O at 25°C [AIEEE 2005]
- (A) 390.7 (B) 217.5 (C) 517.2 (D) 552.7
53. Consider the following E^0 values, [AIEEE 2004]
- $E_{\text{Fe}^{3+}/\text{Fe}^{2+}}^0 = +0.77\text{V}$ $E_{\text{Sn}^{2+}/\text{Sn}}^0 = -0.14\text{V}$
- Under standard conditions the potential for the reaction,
 $\text{Sn(s)} + 2\text{Fe}^{3+}(\text{aq}) \longrightarrow 2\text{Fe}^{2+}(\text{aq}) + \text{Sn}^{2+}(\text{aq})$ is:
- (A) 0.91V (B) 1.40V (C) 1.68V (D) 0.63V

54. The standard e.m.f. of a cell, involving one electron change is found to be 0.591 V at 25°C. The equilibrium constant of the reaction is ($F = 96,500 \text{ C mol}^{-1}$; $R = 8.314 \text{ J K}^{-1} \text{ mol}^{-1}$) [AIEEE 2004]
- (A) 1.0×10^{10} (B) 1.0×10^5 (C) 1.0×10^1 (D) 1.0×10^{30}
55. The limiting molar conductivities Λ^0 for NaCl, KBr and KCl are 126, 152 and $150 \text{ S cm}^2 \text{ mol}^{-1}$ respectively. The Λ_m^0 for NaBr is : [AIEEE 2004]
- (A) $278 \text{ S cm}^2 \text{ mol}^{-1}$ (B) $176 \text{ S cm}^2 \text{ mol}^{-1}$ (C) $128 \text{ S cm}^2 \text{ mol}^{-1}$ (D) $302 \text{ S cm}^2 \text{ mol}^{-1}$
56. In a cell that utilises the reaction, [AIEEE 2004]
- $$\text{Zn(s)} + 2\text{H}^+(\text{aq}) \longrightarrow \text{Zn}^{2+}(\text{aq}) + \text{H}_2(\text{g})$$
- addition of H_2SO_4 to cathode compartment, will :
- (A) increase the E_{cell} and shift equilibrium to the right
 (B) lower the E_{cell} and shift equilibrium to the right
 (C) lower the E_{cell} and shift equilibrium to the left
 (D) increase the E_{cell} and shift equilibrium to the left
47. The $E_{\text{M}^{3+}/\text{M}^{2+}}^0$ values for Cr, Mn, Fe and Co are -0.41, +1.57, +0.77 and +1.97V respectively. For which one of these metals the change in oxidation state from +2 to +3 is easiest? [AIEEE 2004]
- (A) Fe (B) Mn (C) Cr (D) Co
58. For a cell reaction involving a two-electron change, the standard e.m.f. of the cell is found to be 0.295V at 25°C. The equilibrium constant of the reaction at 25°C will be : [AIEEE 2003]
- (A) 10 (B) 1×10^{10} (C) 1×10^{-10} (D) 29.5×10^{-2}
59. Standard reduction electrode potentials of three metals A, B and C are respectively +0.5V, -3.0V and -1.2 V. The reducing powers of these metals are : [AIEEE 2003]
- (A) $\text{C} > \text{B} > \text{A}$ (B) $\text{A} > \text{C} > \text{B}$ (C) $\text{B} > \text{C} > \text{A}$ (D) $\text{A} > \text{B} > \text{C}$
60. For the redox reaction :
- $$\text{Zn(s)} + \text{Cu}^{2+} (0.1\text{M}) \rightarrow \text{Zn}^{2+} (1\text{M}) + \text{Cu(s)}$$
- taking place in a cell, E°_{Cell} is 1.10 volt. E_{Cell} for the cell will be $\left(2.303 \frac{RT}{F} = 0.0591 \right)$ [AIEEE 2003]
- (A) 1.07 volt (B) 0.82 volt (C) 2.14 volt (D) 1.80 volt
61. When, during electrolysis of a solution of AgNO_3 9650 coulombs of charge pass through the electroplating bath, the mass of silver deposited on the cathode will be : [AIEEE 2003]
- (A) 21.6g (B) 108g (C) 1.08g (D) 10.8g

EXERCISE # (JEE-ADVANCED)

1. Plotting $1/\Lambda_m$ against $c\Lambda_m$ for aqueous solutions of a monobasic weak acid (HX) resulted in a straight line with y-axis intercept of P and slope of S. The ratio P/S is [JEE Adv.2023]

$[\Lambda_m = \text{molar conductivity}]$

$\Lambda_m^\circ = \text{limiting molar conductivity}$

$c = \text{molar concentration}$

$K_a = \text{dissociation constant of HX}$

- (A) $K_a\Lambda_m^\circ$ (B) $K_a\Lambda_m^\circ/2$ (C) $2 K_a\Lambda_m^\circ$ (D) $1/(K_a\Lambda_m^\circ)$

2. The reduction potential (E° , in V) of $\text{MnO}_4^- (\text{aq})/\text{Mn}(\text{s})$ is

[Given: $E^\circ_{(\text{MnO}_4^- (\text{aq})/\text{MnO}_2 (\text{s}))} = 1.68 \text{ V}$; $E^\circ_{(\text{MnO}_2 (\text{s})/\text{Mn}^{2+} (\text{aq}))} = 1.21 \text{ V}$; $E^\circ_{(\text{Mn}^{2+} (\text{aq})/\text{Mn}(\text{s}))} = -1.03 \text{ V}$]

[JEE Adv.2022]

3. Consider the strong electrolytes Z_mX_n , U_mY_p and V_mX_n . Limiting molar conductivity (Λ°) of U_mY_p and V_mX_n are 250 and $440 \text{ S cm}^2 \text{ mol}^{-1}$, respectively. The value of $(m + n + p)$ is


Given:

Ion	Z^{n+}	U^{p+}	V^{n+}	X^{m-}	Y^{m-}
$\lambda^\circ (\text{S cm}^2 \text{ mol}^{-1})$	50.0	25.0	100.0	80.0	100.0


λ° is the limiting molar conductivity of ions

The plot of molar conductivity (Λ) of Z_mX_n vs $c^{1/2}$ is given below.

[JEE Adv.2022]





4. Consider a 70% efficient hydrogen-oxygen fuel cell working under standard conditions at 1 bar and 298 K. Its cell reaction is [JEE Adv.2020]



The work derived from the cell on the consumption of 1.0×10^{-3} mol of $\text{H}_2(\text{g})$ is used to compress 1.00 mol of a monoatomic ideal gas in a thermally insulated container. What is the change in the temperature (in K) of the ideal gas?

The standard reduction potentials for the two half-cells are given below.



Use $F = 96500 \text{ C mol}^{-1}$, $R = 8.314 \text{ J mol}^{-1} \text{ K}^{-1}$.

Passage 5 and 6

At 298 K, the limiting molar conductivity of a weak monobasic acid is $4 \times 10^2 \text{ S cm}^2 \text{ mol}^{-1}$. At 298 K, for an aqueous solution of the acid, the degree of dissociation is α and the molar conductivity is $y \times 10^2 \text{ S cm}^2 \text{ mol}^{-1}$. At 298 K, upon 20 times dilution with water, the molar conductivity of the solution becomes $3y \times 10^2 \text{ S cm}^2 \text{ mol}^{-1}$ [JEE Adv.2020]

5. The value of α is ____.

6. The value of y is ____.

7. For the electrochemical cell,



the standard emf of the cell is 2.70 V at 300 K. When the concentration of Mg^{2+} is changed to $x \text{ M}$, the cell potential changes to 2.67 V at 300 K. The value of x is ____.

(given, $\frac{F}{R} = 11500 \text{ KV}^{-1}$), where F is the Faraday constant and R is the gas constant,

$$\ln(10) = 2.30$$

[JEE Adv. 2018]

8. Consider an electrochemical cell: $\text{A}(\text{s}) | \text{A}^{n+}(\text{aq}, 2 \text{ M}) || \text{B}^{2n+}(\text{aq}, 1 \text{ M}) | \text{B}(\text{s})$. The value of ΔH^0 for the cell reaction is twice that of ΔG^0 at 300 K. If the emf of the cell is zero, the ΔS^0 (in $\text{J K}^{-1} \text{ mol}^{-1}$) of the cell reaction per mole of B formed at 300 K is ____.

(Given: $\ln(2) = 0.7$, R (universal gas constant) = $8.3 \text{ J K}^{-1} \text{ mol}^{-1}$. H , S and G are enthalpy, entropy and Gibbs energy, respectively.)

[JEE Adv. 2018]

9. The conductance of a 0.0015 M aqueous solution of a weak monobasic acid was determined by using a conductivity cell consisting of platinized Pt electrodes. The distance between the electrodes is 120 cm with an area of cross section of 1 cm^2 . The conductance of this solution was found to be $5 \times 10^{-7} \text{ S}$. The pH of the solution is 4. The value of limiting molar conductivity (Λ_m^0) of this weak monobasic acid in aqueous solution is $Z \times 10^2 \text{ S cm}^{-1} \text{ mol}^{-1}$. The value of Z is

[JEE Adv. 2017]

10. For the following cell,




when the concentration of Zn^{2+} is 10 times the concentration of Cu^{2+} , then expression for ΔG (in J mol^{-1}) is [F is faraday constant; R is gas constant; T is temperature; $E^\circ(\text{cell}) = 1.1\text{V}$]

[JEE Adv. 2017]

- (A) $2.303RT - 2.2 F$ (B) $- 2.2 F$
(C) $1.1F$ (D) $2.303RT + 1.1 F$

11. For the following electrochemical cell at 298K,

[JEE-Adv. 2016]



$$E_{\text{cell}} = 0.092 \text{ V when } \frac{[\text{M}^{2+}(\text{aq})]}{[\text{M}^{4+}(\text{aq})]} = 10^x$$

$$\text{Given : } E_{\text{M}^{4+}/\text{M}^{2+}}^\circ = 0.151 \text{ V ; } 2.303 \frac{RT}{F} = 0.059 \text{ V}$$

The value of x is -

- (A) -2 (B) -1 (C) 1 (D) 2
12. All the energy released from the reaction $\text{X} \rightarrow \text{Y}$, $\Delta_r G^\circ = -193 \text{ kJ mol}^{-1}$ is used for the oxidizing M^+ and $\text{M}^+ \rightarrow \text{M}^{3+} + 2\text{e}^-$, $E^\circ = -0.25 \text{ V}$.

Under standard conditions, the number of moles of M^+ oxidized when one mole of X is converted to Y is [F = 96500 C mol^{-1}]

[JEE-Adv. 2015]

13. The molar conductivity of a solution of a weak acid HX (0.01 M) is 10 times smaller than the molar conductivity of a solution of a weak acid HY (0.1 M). If $\lambda_{\text{X}^-}^0 \approx \lambda_{\text{Y}^-}^0$, the difference in their pK_a values, $\text{pK}_a(\text{HX}) - \text{pK}_a(\text{HY})$, is (consider degree of ionization of both acids to be $\ll 1$).

[JEE-Adv. 2015]

14. In a galvanic cell, the salt bridge -

[JEE-Adv. 2014]

- (A) Does not participate chemically in the cell reaction
(B) Stops the diffusion of ions from one electrode to another
(C) Is necessary for the occurrence of the cell reaction
(D) Ensures mixing of the two electrolytic solutions

15. An aqueous solution of X is added slowly to an aqueous solution of Y as shown in List-I. The variation in conductivity of these reactions is given in List-II. Match List-I with List-II and select the correct answer using the code given below the lists : [JEE-Adv. 2013]

List-I		List-II	
(P)	$(\text{C}_2\text{H}_5)_3\text{N} + \text{CH}_3\text{COOH}$ X Y	(1)	Conductivity decreases and then increases
(Q)	$\text{KI}(0.1\text{M}) + \text{AgNO}_3(0.01\text{M})$ X Y	(2)	Conductivity decreases and then does not change much
(R)	$\text{CH}_3\text{COOH} + \text{KOH}$ X Y	(3)	Conductivity increases and then does not change much
(S)	$\text{NaOH} + \text{HI}$ X Y	(4)	Conductivity does not change much and then increases

Codes :

	P	Q	R	S		P	Q	R	S
(A)	3	4	2	1	(B)	4	3	2	1
(C)	2	3	4	1	(D)	1	4	3	2

16. The standard reduction potential data at 25°C is given below [JEE-Adv. 2013]

$$E^\circ (\text{Fe}^{3+}, \text{Fe}^{2+}) = +0.77 \text{ V} ;$$

$$E^\circ (\text{Fe}^{2+}, \text{Fe}) = -0.44 \text{ V} ;$$

$$E^\circ (\text{Cu}^{2+}, \text{Cu}) = +0.34 \text{ V} ;$$

$$E^\circ (\text{Cu}^+, \text{Cu}) = +0.52 \text{ V} ;$$

$$E^\circ [\text{O}_2(\text{g}) + 4\text{H}^+ + 4\text{e}^- \rightarrow 2\text{H}_2\text{O}] = +1.23 \text{ V} ;$$

$$E^\circ [(\text{O}_2(\text{g}) + 2\text{H}_2\text{O} + 4\text{e}^- \rightarrow 4\text{OH}^-)] = +0.40 \text{ V} ;$$

$$E^\circ (\text{Cr}^{3+}, \text{Cr}) = -0.74 \text{ V} ;$$

$$E^\circ (\text{Cr}^{2+}, \text{Cr}) = -0.91 \text{ V} ;$$

Match E° of the redox pair in List-I with the values given in List-II and select the correct answer using the code given below the lists :

List-I		List-II	
(P)	$E^\circ(\text{Fe}^{3+}, \text{Fe})$	(1)	- 0.18 V
(Q)	$E^\circ(4\text{H}_2\text{O} \rightleftharpoons 4\text{H}^+ + 4\text{OH}^-)$	(2)	- 0.4 V
(R)	$E^\circ(\text{Cu}^{2+} + \text{Cu} \rightarrow 2\text{Cu}^+)$	(3)	- 0.04 V
(S)	$E^\circ(\text{Cr}^{3+}, \text{Cr}^{2+})$	(4)	- 0.83 V

Codes :

	P	Q	R	S		P	Q	R	S
(A)	4	1	2	3	(B)	2	3	4	1
(C)	1	2	3	4	(D)	3	4	1	2

Paragraph for Question 17 and 18

The electrochemical cell shown below is a concentration cell.


$M | M^{2+}$ (saturated solution of a sparingly soluble salt, MX_2) $|| M^{2+}$ ($0.001 \text{ mol dm}^{-3}$) $| M$ The emf of the cell depends on the difference in concentrations of M^{2+} ions at the two electrodes.

The emf of the cell at 298 K is 0.059V.

[JEE 2012]

17. The value of ΔG (kJ mol^{-1}) for the given cell is (take $F = 96500 \text{ C mol}^{-1}$)
 (A) -5.7 (B) 5.7 (C) 11.4 (D) -11.4.
18. The solubility product (K_{sp} ; $\text{mol}^3 \text{ dm}^{-9}$) of MX_2 at 298 K based on the information available for the given concentration cell is (take $2.303 \times R \times 298/F = 0.059 \text{ V}$)
 (A) 1×10^{-15} (B) 4×10^{-15} (C) 1×10^{-12} (D) 1×10^{-12}
19. AgNO_3 (a) was added to an aqueous KCl solution gradually and the conductivity of the solution was measured. the plot of conductance (Λ) versus the volume of AgNO_3 is

[JEE 2011]



- (A) (P) (B) (Q) (C) (R) (D) (S)
20. Consider the following cell reaction : [JEE 2011]
 $2\text{Fe(s)} + \text{O}_{2(\text{g})} + 4\text{H}^+_{(\text{aq})} \rightarrow 2\text{Fe}^{2+}_{(\text{aq})} + 2\text{H}_2\text{O(l)} \quad E^\circ = 1.67 \text{ V}$
 At $[\text{Fe}^{2+}] = 10^{-3} \text{ M}$, $P(\text{O}_2) = 0.1 \text{ atm}$ and $\text{pH} = 3$, the cell potential at 25°C is –
 (A) 1.47 V (B) 1.77 V (C) 1.87 V (D) 1.57 V

Paragraph for Questions 21 to 22

The concentration of potassium ions inside a biological cell is at least twenty times higher than the outside. The resulting potential difference across the cell is important in several processes such as transmission of nerve impulses and maintaining the ion balance. A simple model for such a concentration cell involving a metal M is :

[JEE 2010]

$M(\text{s}) | M^+(\text{aq}; 0.05 \text{ molar}) || M^+(\text{aq}; 1 \text{ molar}) | M(\text{s})$

For the above electrolytic cell the magnitude of the cell potential $|E_{\text{cell}}| = 70 \text{ mV}$.

21. For the above cell :–
 (A) $E_{\text{cell}} < 0$; $\Delta G > 0$ (B) $E_{\text{cell}} > 0$; $\Delta G < 0$ (C) $E_{\text{cell}} < 0$; $\Delta G > 0$ (D) $E_{\text{cell}} > 0$; $\Delta G < 0$
22. If the 0.05 molar solution of M^+ is replaced by a 0.0025 molar M^+ solution, then the magnitude of the cell potential would be :–
 (A) 35 mV (B) 70 mV (C) 140 mV (D) 700 mV

23. For the reaction of NO_3^- ion in an aqueous solution, E° is +0.96 V. Values of E° for some metal ions are given below



[JEE 2009]

The pair(s) of metal that is(are) oxidised by NO_3^- in aqueous solution is(are)

- (A) V and Hg (B) Hg and Fe (C) Fe and Au (D) Fe and V
24. Electrolysis of dilute aqueous NaCl solution was carried out by passing 10 milli ampere current. The time required to liberate 0.01 mol of H_2 gas at the cathode is (1 Faraday = 96500 C mol^{-1})

[JEE 2008]

- (A) $9.65 \times 10^4 \text{ sec}$ (B) $19.3 \times 10^4 \text{ sec}$ (C) $28.95 \times 10^4 \text{ sec}$ (D) $38.6 \times 10^4 \text{ sec}$

Paragraph for Question Nos. 25 & 26 (2 questions)

Redox reactions play a pivoted role in chemistry and biology. The values of standard redox potential (E°) of two half-cell reactions decide which way the reaction is expected to proceed. A simple example is a Daniel cell in which zinc goes into solution and copper gets deposited. Given below are a set of half-cell reactions (acidic medium) along with their E° (V with respect to normal hydrogen electrode) values. Using this data obtain the correct explanations to Questions 14-16.



[JEE 2007]

25. Among the following, identify the correct statement.
- (A) Chloride ion is oxidised by O_2 (B) Fe^{2+} is oxidised by iodine
(C) Iodine ion is oxidised by chlorine (D) Mn^{2+} is oxidised by chlorine
26. While Fe^{3+} is stable, Mn^{3+} is not stable in acid solution because
- (A) O_2 oxidises Mn^{2+} to Mn^{3+}
(B) O_2 oxidises both Mn^{2+} to Mn^{3+} and Fe^{2+} to Fe^{3+}
(C) Fe^{3+} oxidises H_2O to O_2
(D) Mn^{3+} oxidises H_2O to O_2


Paragraph for Question Nos. 27 to 29 (3 questions)

Chemical reactions involve interaction of atoms and molecules. A large number of atoms/molecules (approximately 6.023×10^{23}) are present in a few grams of any chemical compound varying with their atomic/molecular masses. To handle such large numbers conveniently, the mole concept was introduced. This concept has implications in diverse areas such as analytical chemistry, biochemistry, electrochemistry and radiochemistry. The following example illustrates a typical case, involving chemical/ electrochemical reaction, which requires a clear understanding of the mole concept. A 4.0 molar aqueous solution of NaCl is prepared and 500 mL of this solution is electrolysed. This leads to the evolution of chlorine gas at one of the electrodes (atomic mass : Na = 23, Hg = 200; 1 Faraday = 96500 coulombs) [JEE 2007]

27. The total number of moles of chlorine gas evolved is
(A) 0.5 (B) 1.0 (C) 2.0 (D) 3.0
28. If the cathode is a Hg electrode, the maximum weight (g) of amalgam formed from this solution is
(A) 200 (B) 225 (C) 400 (D) 446
29. The total charge (coulombs) required for complete electrolysis is
(A) 24125 (B) 48250 (C) 96500 (D) 193000

Question No. 30 to 31 (3 questions)

Tollen's reagent is used for the detection of aldehyde when a solution of AgNO_3 is added to glucose with NH_4OH then gluconic acid is formed



$$\left[\text{Use } 2.303 \times \frac{RT}{F} = 0.0592 \text{ and } \frac{F}{RT} = 38.92 \text{ at } 298 \text{ K} \right]$$


[JEE 2006]

30. $2\text{Ag}^+ + \text{C}_6\text{H}_{12}\text{O}_6 + \text{H}_2\text{O} \longrightarrow 2\text{Ag(s)} + \text{C}_6\text{H}_{12}\text{O}_7 + 2\text{H}^+$
Find $\ln K$ of this reaction
(A) 66.13 (B) 58.38 (C) 28.30 (D) 46.29
31. When ammonia is added to the solution, pH is raised to 11. Which half-cell reaction is affected by pH and by how much?
(A) E_{oxd} will increase by a factor of 0.65 from E^0_{oxd}
(B) E_{oxd} will decrease by a factor of 0.65 from E^0_{oxd}
(C) E_{red} will increase by a factor of 0.65 from E^0_{red}
(D) E_{red} will decrease by a factor of 0.65 from E^0_{red}

32. Ammonia is always added in this reaction. Which of the following must be incorrect?
 (A) NH_3 combines with Ag^+ to form a complex.
 (B) $\text{Ag}(\text{NH}_3)_2^+$ is a weaker oxidising reagent than Ag^+ .
 (C) In absence of NH_3 silver salt of gluconic acid is formed.
 (D) NH_3 has affected the standard reduction potential of glucose/gluconic acid electrode.
33. We have taken a saturated solution of $\text{AgBr.K}_{\text{sp}}$ of AgBr is 12×10^{-14} . If 10^{-7} mole of AgNO_3 are added to 1 litre of this solution find conductivity (specific conductance) of this solution in terms of $10^{-7} \text{ S m}^{-1} \text{ mol}^{-1}$. [JEE 2006]

Given : $\lambda_{(\text{Ag}^+)}^0 = 6 \times 10^{-3} \text{ S m}^2 \text{ mol}^{-1}$; $\lambda_{(\text{Br}^-)}^0 = 8 \times 10^{-3} \text{ S m}^2 \text{ mol}^{-1}$; $\lambda_{(\text{NO}_3^-)}^0 = 7 \times 10^{-3} \text{ S m}^2 \text{ mol}^{-1}$


34. The half cell reactions for rusting of iron are: [JEE 2005]



ΔG^0 (in kJ) for the reaction is :

- (A) -76 (B) -322 (C) -122 (D) -176

35. (a) Calculate ΔG_f^0 of the following reaction [JEE 2005]



Given : $\Delta G_f^0 (\text{AgCl}) = -109 \text{ kJ/mole}$, $\Delta G_f^0 (\text{Cl}^-) = -129 \text{ kJ/mole}$, $\Delta G_f^0 (\text{Ag}^+) = 77 \text{ kJ/mole}$

Represent the above reaction in form of a cell

Calculate E^0 of the cell. Find $\log_{10} K_{\text{SP}}$ of AgCl

(B) (b) $6.539 \times 10^{-2} \text{ g}$ of metallic Zn ($\text{amu} = 65.39$) was added to 100 ml of saturated

solution of AgCl . Calculate $\frac{[\text{Zn}^{2+}]}{[\text{Ag}^+]^2} \log_{10}$, given that




Also find how many moles of Ag will be formed ?

36. $\text{Zn} | \text{Zn}^{2+} (a = 0.1\text{M}) || \text{Fe}^{2+} (a = 0.01\text{M}) | \text{Fe}$. The emf of the above cell is 0.2905 V. Equilibrium constant for the cell reaction is [JEE 2004]

- (A) $10^{0.32/0.0591}$ (B) $10^{0.32/0.0295}$ (C) $10^{0.26/0.0295}$ (D) $e^{0.32/0.295}$

37. Find the equilibrium constant at 298 K for the reaction,



Given that $E_{\text{Cu}^{2+}|\text{Cu}^+}^0 = 0.15 \text{ V}$, $E_{\text{In}^{3+}|\text{In}^{2+}}^0 = -0.42 \text{ V}$, $E_{\text{In}^{2+}|\text{In}^+}^0 = -0.40 \text{ V}$

[JEE 2004]

38. In the electrolytic cell, flow of electrons is from:

- (A) Cathode to anode in solution (B) Cathode to anode through external supply
 (C) Cathode to anode through internal supply (D) Anode to cathode through internal supply.

39. Two students use same stock solution of ZnSO_4 and a solution of CuSO_4 . The e.m.f of one cell is 0.03 V higher than the other. The conc. of CuSO_4 in the cell with higher e.m.f value is 0.5 M. Find out the conc. of CuSO_4 in the other cell $\left(\frac{2.303 RT}{F} = 0.06 \right)$. [JEE 2003]
40. Standard electrode potential data are useful for understanding the suitability of an oxidant in a redox titration. Some half cell reactions and their standard potentials are given below:
 $\text{MnO}_4^- (\text{aq}) + 8\text{H}^+ (\text{aq}) + 5\text{e}^- \longrightarrow \text{Mn}^{2+} (\text{aq}) + 4\text{H}_2\text{O} (\text{l}); E^\circ = 1.51 \text{ V}$ [JEE 2002]
 $\text{Cr}_2\text{O}_7^{2-} (\text{aq}) + 14\text{H}^+ (\text{aq}) + 6\text{e}^- \longrightarrow 2\text{Cr}^{3+} (\text{aq}) + 7\text{H}_2\text{O} (\text{l}); E^\circ = 1.38 \text{ V}$
 $\text{Fe}^{3+} (\text{aq}) + \text{e}^- \longrightarrow \text{Fe}^{2+} (\text{aq}); E^\circ = 0.77 \text{ V}$
 $\text{Cl}_2 (\text{g}) + 2\text{e}^- \longrightarrow 2\text{Cl}^- (\text{aq}); E^\circ = 1.40 \text{ V}$
 Identify the only incorrect statement regarding quantitative estimation of aqueous $\text{Fe}(\text{NO}_3)_2$
 (A) MnO_4^- can be used in aqueous HCl (B) $\text{Cr}_2\text{O}_7^{2-}$ can be used in aqueous HCl
 (C) MnO_4^- can be used in aqueous H_2SO_4 (D) $\text{Cr}_2\text{O}_7^{2-}$ can be used in aqueous H_2SO_4
41. The reaction, [JEE 2001]
 $3\text{ClO}^- (\text{aq}) \longrightarrow \text{ClO}_3^- (\text{aq}) + 2\text{Cl}^- (\text{aq})$
 is an example of
 (A) Oxidation reaction (B) Reduction reaction
 (C) Disproportionation reaction (D) Decomposition reaction
42. The correct order of equivalent conductance at infinite dilution of LiCl , NaCl and KCl is
 (A) $\text{LiCl} > \text{NaCl} > \text{KCl}$ (B) $\text{KCl} > \text{NaCl} > \text{LiCl}$ [JEE 2001]
 (C) $\text{NaCl} > \text{KCl} > \text{LiCl}$ (D) $\text{LiCl} > \text{KCl} > \text{NaCl}$
43. Saturated solution of KNO_3 is used to make salt bridge because
 (A) velocity of K^+ is greater than that of NO_3^-
 (B) velocity of NO_3^- is greater than that of K^+
 (C) velocities of both K^+ and NO_3^- are nearly the same
 (D) KNO_3 is highly soluble in water [JEE 2001]
44. The standard potential of the following cell is 0.23 V at 15°C & 0.21 V at 35°C
 $\text{Pt} | \text{H}_2 (\text{g}) | \text{HCl} (\text{aq}) | \text{AgCl} (\text{s}) | \text{Ag} (\text{s})$
 (i) Write the cell reaction.
 (ii) Calculate ΔH° , ΔS° for the cell reaction by assuming that these quantities remain unchanged in the range 15°C to 35°C .
 (iii) Calculate the solubility of AgCl in water at 25°C . Given standard reduction potential of the Ag^+/Ag couple is 0.80 V at 25°C . [JEE 2001]

45. For the electrochemical cell, $M | M^+ || X^- | X$, $E^\circ (M^+/M) = 0.44 \text{ V}$ and $E^\circ (X/X^-) = 0.33 \text{ V}$. From this data, one can deduce that [JEE 2000]
- (A) $M + X \longrightarrow M^+ + X^-$ is the spontaneous reaction
 (B) $M^+ + X^- \longrightarrow M + X$ is the spontaneous reaction
 (C) $E_{\text{cell}} = 0.77 \text{ V}$
 (D) $E_{\text{cell}} = -0.77 \text{ V}$
46. Copper sulphate solution (250 mL) was electrolysed using a platinum anode and a copper cathode. A constant current of 2 mA was passed for 16 min. It was found that after electrolysis, the absorbance (concentration) of the solution was reduced to 50% of its original value. Calculate the concentration of copper sulphate in the solution to begin with. [JEE 2000]
47. The following electrochemical cell has been set up
 $\text{Pt(I)} | \text{Fe}^{3+}, \text{Fe}^{2+} (a=1) || \text{Ce}^{4+}, \text{Ce}^{3+} (a=1) | \text{Pt(II)}$
 $E_{\text{Fe}^{3+}/\text{Fe}^{2+}}^0 = 0.77 \text{ V}$ and $E_{\text{Ce}^{4+}/\text{Ce}^{3+}}^0 = 1.61 \text{ V}$
 If an ammeter is connected between the two platinum electrodes, predict the direction of flow of current. Will the current increase or decrease with time? [JEE 2000]
48. A gas X at 1 atm is bubbled through a solution containing a mixture of 1 M Y and 1 M Z at 25°C. If the reduction potential of $Z > Y > X$, then [JEE 1999]
- (A) Y will oxidise X and not Z
 (B) Y will oxidise Z and X
 (C) Y will oxidise both X and Z
 (D) Y will reduce both X and Z.
49. Calculate the equilibrium constant for the reaction, $2\text{Fe}^{3+} + 3\text{I}^- \rightleftharpoons 2\text{Fe}^{2+} + \text{I}_3^-$. The standard reduction potentials in acidic conditions are 0.77 and 0.54 V respectively for $\text{Fe}^{3+} / \text{Fe}^{2+}$ and $\text{I}_3^- / \text{I}^-$ couples. [JEE 1998]
50. Find the solubility product of a saturated solution of Ag_2CrO_4 in water at 298 K if the emf of the cell $\text{Ag} | \text{Ag}^+ (\text{satd. Ag}_2\text{CrO}_4 \text{ soln.}) || \text{Ag}^+ (0.1 \text{ M}) | \text{Ag}$ is 0.164 V at 298 K. [JEE 1998]

ANSWER KEY

EXERCISE # (O-I)

- | | | | | | | |
|---------|---------|---------|---------|-------------------------------------|---------|---------|
| 1. (A) | 2. (C) | 3. (D) | 4. (B) | 5. (B) | 6. (D) | 7. (A) |
| 8. (C) | 9. (A) | 10. (D) | 11. (C) | 12. (1.61 V) | | 13. (A) |
| 14. (C) | 15. (C) | 16. (C) | 17. (C) | 18. ($K_c = 2.18 \times 10^{26}$) | | 19. (C) |
| 20. (A) | 21. (A) | 22. (A) | 23. (C) | 24. (A) | 25. (B) | 26. (B) |
| 27. (A) | 28. (B) | 29. (B) | 30. (D) | 31. (B) | 32. (D) | 33. (A) |
| 34. (C) | 35. (D) | 36. (C) | 37. (B) | 38. (A) | 39. (B) | 40. (C) |
| 41. (A) | 42. (C) | 43. (C) | 44. (B) | 45. (D) | 46. (A) | 47. (C) |
| 48. (D) | 49. (D) | 50. (D) | 51. (C) | 52. (D) | 53. (C) | 54. (D) |
| 55. (D) | 56. (A) | 57. (C) | 58. (C) | 59. (A) | 60. (C) | 61. (B) |
| 62. (B) | 63. (B) | 64. (A) | 65. (C) | 66. (C) | 67. (C) | 68. (B) |
| 69. (C) | 70. (B) | 71. (A) | 72. (A) | 73. (D) | 74. (B) | 75. (D) |
| 76. (B) | 77. (D) | 78. (A) | 79. (C) | 80. (C) | 81. (A) | 82. (B) |
| 83. (B) | 84. (C) | 85. (B) | 86. (C) | 87. (B) | 88. (D) | 89. (A) |

EXERCISE S-I

- (a) Cu (b) Ag

(c) oxidation (d) reduction

(e) Cu (f) Ag

(g) anode-Cu \longrightarrow $\text{Cu}^{2+} + 2\text{e}^-$; cathode- $\text{Ag}^+ + \text{e}^- \longrightarrow \text{Ag}$

(h) $\text{Cu} + 2\text{Ag}^+ \longrightarrow \text{Cu}^{2+} + 2\text{Ag}$ (i) Cu

(j) Cu (k) to maintain electrical neutrality in solution
- (1.68 V)
- (1.35 V)
- (- 0.0367 V)
- (0.53 V)
- (-0.756 V)
- (yes)
- (Spontaneous, - 48250 J)
- ($K_c = 7.6 \times 10^{12}$)
- ($K_c = 2.868 \times 10^{107}$, $\Delta G^0 = - 611.8 \text{ kJ}$)
- ($\log [\text{Zn}^{2+}] / [\text{Cu}^{2+}] = 37.22$)
- ($K_w \gg 10^{-14}$)
- ($K = 10^{268}$)
- (-1.30 $\times 10^3 \text{ kJ mol}^{-1}$)
- (Mg)
- (Y > Z > X)
- Zn
- (- 0.80 V, NO)
- (a) $2\text{Ag} + \text{Cu}^{2+} \longrightarrow 2\text{Ag}^+ + \text{Cu}$, (b) $\text{MnO}_4^- + 5\text{Fe}^{2+} + 8\text{H}^+ \longrightarrow \text{Mn}^{2+} + 5\text{Fe}^{3+} + 4\text{H}_2\text{O}$

(c) $2\text{Cl}^- + 2\text{Ag}^+ \longrightarrow 2\text{Ag} + \text{Cl}_2$, (d) $\text{H}_2 + \text{Cd}^{2+} \longrightarrow \text{Cd} + 2\text{H}^+$
- (a) $\text{Zn} | \text{Zn}^{2+} || \text{Cd}^{2+} | \text{Cd}$,

(b) $\text{Pt}, \text{H}_2 | \text{H}^+ || \text{Ag}^+ | \text{Ag}$,

(c) $\text{Pt} | \text{Fe}^{2+}, \text{Fe}^{3+} || \text{Cr}_2\text{O}_7^{2-}, \text{H}^+, \text{Cr}^{3+} | \text{Pt}$

(Physical Chemistry)

ELECTROCHEMISTRY

21. (0.059 volt) 22. $[\text{Cu}^{2+}] = 2.97 \times 10^{-12} \text{M}$ for $E = 0$ 23. ($E = -0.81 \text{ V}$)
24. (-0.22 V) 25. ($E = 1.159 \text{ V}$)
26. ($E^0_{\text{cell}} = +0.01 \text{ V}$, $E_{\text{cell}} = -0.0785 \text{ V}$,
No, correct representation is $\text{Pb}|\text{Pb}^{2+} (10^{-3} \text{ M})||\text{Sn}^{2+} (1 \text{ M})|\text{Sn}$)
27. (a) The spontaneous cell reaction : $\text{Zn} + 2\text{Ag}^+ (\text{aq}) \rightarrow \text{Zn}^{2+} (\text{aq}) + 2\text{Ag} (\text{s})$
(b) 1.56 V (c) $[\text{Zn}^{2+}] = 4 \times 10^{-4} \text{ M}$
(d) As we add KI to cathode chamber, some Ag^+ will precipitate out as :
$$\text{Ag}^+ + \text{I}^- \rightarrow \text{AgI}$$

The above reaction reducing $[\text{Ag}^+]$ from cathode chamber. This will reduce E_{cell} according to Nernst's equation.
28. (0.0295 V) 29. ($\text{pH} = 4$) 30. ($E = 0.059$)
31. ($E = 0.413 \text{ V}$)
32. (A) $\text{Hg}_2\text{Cl}_2 (\text{s}) + \text{Cu} (\text{s}) \rightarrow \text{Cu}^{2+} (\text{aq}) + 2\text{Cl}^- (\text{aq}) + 2\text{Hg} (\text{l})$
(B) $2\text{Ag}_{(\text{s})} + 2\text{IO}_3^- + \text{Zn}^{2+} \longrightarrow 2\text{AgIO}_{3(\text{s})} + \text{Zn}_{(\text{s})}$
(C) $\text{Mn}_{(\text{s})} + 2\text{OH}^- + \text{Cu}^{2+} \longrightarrow \text{Mn}(\text{OH})_{2(\text{s})} + \text{Cu}_{(\text{s})}$
33. (0.42 V)
34. ($K_{\text{sp}} = 1.1 \times 10^{-16}$)
35. ($[\text{Br}^-] : [\text{Cl}^-] = 1 : 200$)
36. ($K_{\text{SP}} = 10^{-5}$)
37. (-0.16 V)
38. (a) 6×10^{22} electrons lost,
(b) 1.2×10^{24} electrons gained,
(c) 1.80×10^{23} electrons gained
39. (a) 0.75 F, (b) 0.69 F, (c) 1.1 F
40. (i) 54 gm, (ii) 16.35 gm
41. ($n = 4$)
42. (0.112 litre)
43. (Rs. 0.75 x)
44. (i) 2.17 gm ; (ii) 1336.15 sec
45. (42.2 %)
46. ($1.023 \times 10^5 \text{ sec}$)

47. (71.5 amp)

48.

	ELECTROLYTE	ANODE Product	CATHODE Product
1	NaCl (Molten) with Pt electrode	Cl ₂ (g)	Na
2	NaCl (aq) with Pt electrode	Cl ₂ (g)	H ₂ (g)
3	Na ₂ SO ₄ (aq) with Pt electrode	O ₂ (g)	H ₂ (g)
4	NaNO ₃ (aq) with Pt electrode	O ₂ (g)	H ₂ (g)
5	AgNO ₃ (aq) with Pt electrode	O ₂ (g)	Ag
6	CuSO ₄ (aq) with Inert electrode	O ₂ (g)	Cu
7	CuSO ₄ (aq) with Copper electrode	Cu dissolve	Cu

49. (1.12 mol, 12.535 litre)

50. (60 %)

51. (1.825g)

52. (2M)

53. (4.25 × 10⁻² m)

54. (400 S cm² mol⁻¹)

55. (i) 250 mho cm² mol⁻¹, (ii) 125 mho cm² equivalent⁻¹

56. (0.0125 mho g eq⁻¹ m², 1.25 × 10⁻³ mho cm⁻¹)

57. (0.00040 S cm⁻¹; 2500 ohm cm)

58. (i) 6.25 × 10⁵ ohm, (ii) 1.6 × 10⁻⁶ amp

59. (510 × 10⁻⁴ mho cm² mol⁻¹)

60. (i) 400 S cm² mol⁻¹ (ii) 12 %

61. (0.8)

62. (a = 0.5, k = 10 × 10⁻⁴)

63. (10⁻¹⁰ mole² /litre²)

64. (9.4 × 10⁻⁴ gm/litre)

65. (i) 7 (ii) 1 × 10⁻¹⁴

EXERCISE # (O-II)

- | | | | | | |
|---|-----------|--------------|--------|-------------|-----------|
| 1. (D) | 2. (A) | 3. (B) | 4. (A) | 5. (B, D) | 6. (B,D) |
| 7. (C) | 8. (A,D) | 9. (A,B,C,D) | | 10. (B,C,D) | 11. (A,B) |
| 12. (C,D) | 13. (B,D) | 14. (A,D) | | 15. (A) | 16. (B) |
| 17. (B) | 18. (C) | 19. (B) | | 20. (D) | 21. (B) |
| 22. (C) | 23. (B) | 24. (C) | | 25. (A) | 26. (A) |
| 27. (A) P, Q (B) P, Q (C) Q, R, (D) P,S | | | | 28. (D) | |

EXERCISE # (S-II)

- | | | | |
|---|----------------------|-----------------------|---------------|
| 1. (0.52 V, 0.61 V) | 2. (n = 2) | 3. (0.0295 V) | 4. (- 0.46 V) |
| 5. (ii). 1.27 V, (iii) 245.1 kJ | | 6. (-0.037 V) | |
| 7. (115800C, 347.4 kJ) | | 8. (t = 193 sec) | |
| 9. (Final weight = 9.6g, 0.01 Eq of acid) | 10. (t = 93.65 sec.) | 11. (43.456g) | |
| 12. (1.9×10^6 year) | 13. (0.1456 ampere) | 14. (0.1934 gm/litre) | |

EXERCISE # (JEE-MAINS)

- | | | | | | |
|---------|----------|-------------------|---------|---------|---------|
| 1. (A) | 2. (14) | 3. (127) | 4. (D) | 5. (46) | 6. (24) |
| 7. (12) | 8. (760) | 9. (5.67 or 5.68) | 10. 60 | 11. (D) | 12. (D) |
| 13. (B) | 14. (B) | 15. (A) | 16. (D) | 17. (C) | 18. (C) |
| 20. (C) | 21. (C) | 22. (C) | 23. (C) | 24. (C) | 25. (B) |
| 27. (D) | 28. (B) | 29. (D) | 30. (A) | 31. (A) | 32. (B) |
| 34. (B) | 35. (C) | 36. (C) | 37. (A) | 38. (D) | 39. (A) |
| 41. (B) | 42. (D) | 43. (A) | 44. (A) | 45. (D) | 46. (D) |
| 48. (C) | 49. (C) | 50. (C) | 51. (B) | 52. (A) | 53. (A) |
| 55. (C) | 56. (A) | 57. (C) | 58. (B) | 59. (C) | 60. (A) |
| | | | | | 61. (D) |

EXERCISE # (JEE-ADVANCED)

- | | | | | | |
|--|---|---|------------|-------------|-----------|
| 1. (A) | 2. (0.77) | 3. (7) | 4. (13.32) | 5. (0.215) | 6. (0.86) |
| 7. (10) | 8. (-11.62) | 9. (6) | 10. (A) | 11. (D) | 12. (4) |
| 13. (3) | 14. (A,B) | 15. (A) | 16. (D) | 17. (D) | 18. (B) |
| 19. (D) | 20. (D) | 21. (B) | 22. (C) | 23. (A,B,D) | 24. (B) |
| 25. (C) | 26. (D) | 27. (B) | 28. (D) | 29. (D) | 30. (A) |
| 31. (A) | 32. (D) | 33. (55 S m^{-1}) | 34. (B) | | |
| 35. (a) $E^0 = 0.59 \text{ V}$, $\log_{10} K_{\text{SP}} = -10$ | (b) $52.8, 10^{-6}$ moles | 36. (B) | | | |
| 37. ($K_{\text{C}} = 10^{10}$) | 38. (C) | 39. (0.05) | 40. (A) | 41. (C) | 42. (B) |
| 43. (C) | 44. ($\Delta H^0 = -49987 \text{ J mol}^{-1}$, $\Delta S^0 = -96.5 \text{ J mol}^{-1} \text{ K}^{-1}$, $s = 1.47 \times 10^{-5} \text{ M}$) | | | | |
| 45. (B) | 46. ($7.95 \times 10^{-5} \text{ M}$) | 47. (decrease with time) | | | |
| 48. (A) | 49. ($K_{\text{C}} = 6.26 \times 10^7$) | 50. ($K_{\text{SP}} = 2.287 \times 10^{-12}$) | | | |