

Projectile Motion

Q. An inclined plane makes an angle $\theta_0 = 30^\circ$ with the horizontal. A particle is projected from this plane with a speed of 5 m s^{-1} at an angle of elevation $\beta = 30^\circ$ with the horizontal as shown in Fig.

a. Find the range of the particle on the plane when it strikes the plane.

b. Find the range of the particle for $\beta = 120^\circ$. $\theta = 60^\circ$

Along the Inclined plane:-



$$T = \frac{2u_v}{g_{\text{eff}}}$$

$$T = \frac{2 \times 5 \sin 60^\circ}{g \sin 60^\circ} = 1 \text{ sec.}$$



$$R = (5 \cos 60^\circ)T + \frac{1}{2}(g \cos 60^\circ)T^2$$

$$R = (5 \times \frac{1}{2} \times 1) + \frac{1}{2} \times 10 \times \frac{1}{2} \times (1)^2$$

$$R = \frac{5}{2} + \frac{5}{2} = \underline{5 \text{ m}} \checkmark$$



(b)



$$T = \frac{2 \times \left(\frac{u}{2}\right)}{\frac{\sqrt{3}g}{2}}$$


$$T = \frac{2u}{\sqrt{3}g} = \left(\frac{1}{\sqrt{3}} \text{ Sec}\right) \checkmark$$

$$R = \left(\frac{\sqrt{3}u}{2}\right)T - \frac{1}{2} \times 5 \times T^2$$

$$\begin{aligned} R &= \left(\frac{\sqrt{3}}{2} \times 5 \times \frac{1}{\sqrt{3}}\right) - \frac{1}{2} \times 5 \times \left(\frac{1}{3}\right) \\ &= \left(\frac{5}{2} - \frac{5}{6}\right) = \frac{15-5}{6} = \frac{10}{6} = \left(\frac{5}{3} \text{ m}\right) \end{aligned}$$

Projectile Motion

Q. A body has maximum range R_1 when projected up the plane. The same body when projected down the inclined plane, it has maximum range R_2 . Find the maximum horizontal range. Assume equal speed of projection in each case and the body is projected onto the inclined plane in the line of the greatest slope.



$$R_1 = \frac{u^2}{g(1+\sin\alpha)}, \quad R_2 = \frac{u^2}{g(1-\sin\alpha)}$$


$$1+\sin\alpha = \left(\frac{u^2}{R_1 g}\right) = \frac{R}{R_1} \quad \text{--- (1)}$$

$$1-\sin\alpha = \left(\frac{u^2}{g R_2}\right) = \frac{R}{R_2} \quad \text{--- (2)}$$

$$\frac{(1)+(2)}{2} = \frac{R\left(\frac{1}{R_1} + \frac{1}{R_2}\right)}{2} \Rightarrow R = \left(\frac{2 R_1 R_2}{R_1 + R_2}\right)$$

Projectile Motion

- P.W.*
Q. A small sphere is projected with a velocity of 3 ms^{-1} in a direction 60° from the horizontal y -axis, on the smooth inclined plane (Fig.) The motion of sphere takes place in the $x - y$ plane. Calculate the magnitude v of its velocity after 2 s.



Projectile Motion


Q. A body is projected up along a smooth inclined plane with velocity u from the point A as shown in Fig. The angle of inclination is 45° and the top is connected to a well of diameter 40 m. If the body just manages to cross the well, what is the value of u ? The length of inclined plane is $20\sqrt{2}$ m.

(A) 40 ms^{-1}

(B) $40\sqrt{2} \text{ ms}^{-1}$

(C) 20 m s^{-1}

(D) $20\sqrt{2} \text{ ms}^{-1}$



Projectile Motion

$$T = \left(\frac{2u_y}{g_{\text{eff}}} \right)$$

Q. In Fig., the time taken by the projectile to reach from A to B is t . Then the distance AB is equal to

✓ (A) $\frac{ut}{\sqrt{3}}$ ✓

(B) $\frac{\sqrt{3}ut}{2}$

(C) $\sqrt{3}ut$

(D) $2ut$

$$R = \left(\frac{\sqrt{3}u}{2} \right) t - \frac{1}{2} \times 5 \times t^2$$


$$R = \left(\frac{\sqrt{3}u}{2} \right) t - \frac{5t^2}{2}$$

$$R = \frac{\sqrt{3}}{2} (ut) - \frac{5t(t)}{2}$$

$$R = \frac{\sqrt{3}}{2} (ut) - \frac{5t}{2} \left(\frac{u}{5\sqrt{3}} \right)$$

$$R = \frac{\sqrt{3}}{2} (ut) - \left(\frac{ut}{2\sqrt{3}} \right) = \frac{ut}{2} \left(\sqrt{3} - \frac{1}{\sqrt{3}} \right) = \frac{ut}{2\sqrt{3}} (3-1) = \frac{ut}{\sqrt{3}}$$

$$\left[\begin{array}{l} \theta = 60 - 30 \\ \theta = 30^\circ \end{array} \right]$$




$$t = \frac{2 \times \frac{u}{2}}{\left(\frac{\sqrt{3} \times 10}{2} \right)} = \frac{u}{10 \times \frac{\sqrt{3}}{2}} = \left(\frac{u}{5\sqrt{3}} \right)$$

Projectile Motion

H.W. I.I.T Adv.

Q. Two inclined planes OA and OB having inclinations 30° and 60° with the horizontal respectively intersect each other at O, as shown in figure. A particle is projected from point P with velocity $u = 10\sqrt{3} \frac{\text{m}}{\text{s}}$ along a direction perpendicular to plane OA. If the particle strikes plane OB perpendicular at Q. Calculate

- (A) time of flight
- (B) velocity with which the particle strikes the plane OB.
- (C) height h of point P from point O.
- (D) distance PQ. (Take $g = 10 \text{ m/s}^2$)



Projectile Motion


Q. A projectile is fired with a velocity u at right angles to the slope, which is inclined at an angle θ with the horizontal. Derive an expression for the distance R to the point of impact.

$$T = \left[\frac{2u}{g \cos \theta} \right] \quad u_y = u$$

$$R = \frac{1}{2} \times (g \sin \theta) \times T^2$$


$$R = \frac{1}{2} \times g \sin \theta \times \frac{4u^2}{g^2 \cos^2 \theta}$$

$$R = \left(\frac{2u^2 \sin \theta}{g \cos^2 \theta} \right) \Rightarrow R = \frac{2u^2}{g} (\tan \theta) (\sec \theta)$$



Projectile Motion

- Q.** Cannon A is located on a plain a distance L from a wall of height H . On top of this wall is an identical cannon (cannon B). Ignore air resistance throughout this problem. Also ignore the size of the cannons relative to L and H . The two groups of gunners aim the cannons directly at each other. They fire at each other simultaneously, with equal muzzle speed v_0 . What is the value of v_0 for which the two cannon balls collide just as they hit the ground?



Projectile Motion


Q. A particle is thrown at time $t = 0$ with a velocity of 10 m/s at an angle of 60° with the horizontal from a point on an incline plane, making an angle of 30° with the horizontal. The time when the velocity of the projectile becomes parallel to the incline is:

(A) $\frac{2}{\sqrt{3}} \text{ sec}$

✓ (B) $\frac{1}{\sqrt{3}} \text{ sec}$

(C) $\sqrt{3} \text{ sec}$

(D) $\frac{1}{2\sqrt{3}} \text{ sec}$



Projectile Motion

Q. Find range of projectile which is projected perpendicular to the incline plane with velocity 20 m/s as shown in figure:

(A) 75 m ✓

(B) 40 m

(C) 45 m

(D) 50 m

$$T = \frac{2u_y}{g_{\text{eff}}}$$

$$T = \frac{2 \times 20}{10 \times \frac{4}{5}} = 5 \text{ sec}$$

$$R = \left(\frac{1}{2} a_x T^2 \right) = \frac{1}{2} \times 10 \times \frac{3}{5} \times (5)^2$$

$$R = 75 \text{ m}$$

