


- Q.1** A motorcycle is travelling on a curved track of radius 500 m if the coefficient of friction between road and tyres is 0.5 . The speed avoiding skidding, will be :-
 (A) 50 m/s (B) 75 m/s (C) 25 m/s (D) 35 m/s
- Q.2** A string of length 0.1 m cannot bear a tension more than 100 N. It is tied to a body of mass 100 g and rotated in a horizontal circle. The maximum angular velocity is
 (A) 100rad/s (B) 1000rad/s
 (C) 10000/s (D) 0.1rad/s
- Q.3** A car when passes through a convex bridge exerts a force on it which is equal to :-
 (A) $Mg + \frac{Mv}{r}$ (B) $\frac{Mv^2}{r}$ (C) Mg (D) none of these
- Q.4** A small mass of 10gm. lies in a hemispherical bowl of radius 0.5 m at a height of 0.2 m from the bottom of the bowl. The mass will be in equilibrium if the bowl rotates at an angular speed of ($g = 10 \text{ m/sec}^2$): -



- (A) $\frac{10}{\sqrt{3}} \text{ rad./s}$ (B) $10\sqrt{3} \text{ rad/s}$ (C) 10 rad/s (D) $\sqrt{20} \text{ rad./s}$
- Q.5** A particle is acted upon by a force of constant magnitude which is always perpendicular to the velocity of the particle. The motion of the particle takes place in a plane, it follows that-
 (A) Its velocity is constant
 (B) Its acceleration is constant
 (C) Its kinetic energy is constant
 (D) It moves in a straight line
- Q.6** A tube of length L is filled completely with an incompressible liquid of mass M and closed at both the ends. The tube is then rotated in a horizontal plane about one of its ends with a uniform angular velocity ω . The force exerted by the liquid at the other end is

- (A) $\frac{ML\omega^2}{2}$ (B) $ML\omega^2$ (C) $\frac{ML\omega^2}{4}$ (D) $\frac{ML^2\omega^2}{2}$

(Physics)

CIRCULAR MOTION

Q.7 For a particle in uniform circular motion, the acceleration \vec{a} at a point $P(R, \theta)$ on the circle of radius R is (Here θ is measured from the x-axis).

(A) $\frac{v^2}{R} \hat{i} + \frac{v^2}{R} \hat{j}$

(B) $-\frac{v^2}{R} \cos \theta \hat{i} + \frac{v^2}{R} \sin \theta \hat{j}$

(C) $-\frac{v^2}{R} \sin \theta \hat{i} + \frac{v^2}{R} \cos \theta \hat{j}$

(D) $-\frac{v^2}{R} \cos \theta \hat{i} - \frac{v^2}{R} \sin \theta \hat{j}$

Q.8 A gramophone record is revolving with an angular velocity ω . A coin is placed at a distance r from the centre of the record. The static coefficient of friction is μ . The coin will revolve with the record if :-


(A) $r \geq \frac{\mu g}{\omega^2}$

(B) $r = \mu g \omega^2$

(C) $r < \frac{\omega^2}{\mu g}$

(D) $r \leq \frac{\mu g}{\omega^2}$

Q.9 A bead is constrained to move on rod in gravity free space as shown in figure. The rod is rotating with angular velocity ω and angular acceleration α about its end. If μ is coefficient of friction. Mark the correct option. (Rod rotates in the plane of paper.)



(A) If $\mu = \frac{\omega^2}{\alpha}$ friction on bead is static in nature

(B) If $\mu > \frac{\omega^2}{\alpha}$ friction on bead is kinetic in nature

(C) If $\mu < \frac{\omega^2}{\alpha}$ friction is static

(D) If bead does not slide relative to rod. Friction will not exist between bead and rod.

Q.10 A particle is projected with initial speed u and at an angle θ with horizontal. What is the radius of curvature of the parabola traced out by the projectile at a point where the particle velocity makes an angle $\theta/2$ with the horizontal?

(A) $\frac{u^2 \cos^2 \theta}{g}$

(B) $\frac{u^2 \cos^2 \theta}{g \cos^3 \theta/2}$

(C) $\frac{u^2 \sin^2 \theta}{g \cos^3 \theta}$

(D) $N \cdot O \cdot T$

Q.11. A train has to negotiate a curve of radius 400 m. By how much height should the outer rail be raised with respect to inner rail for a speed of 48 km/hr ? The distance between the rails is 1m :

(A) $\frac{1}{15}$

(B) $\frac{2}{25}$

(C) $\frac{2}{45}$

(D) $\frac{3}{17}$



ANSWER KEY

- | | | | | | | | | | | | | | |
|----|-----|----|-----|-----|-----|-----|-----|----|-----|----|-----|----|-----|
| 1. | (A) | 2. | (A) | 3. | (D) | 4. | (A) | 5. | (C) | 6. | (A) | 7. | (D) |
| 8. | (D) | 9. | (A) | 10. | (B) | 11. | (C) | | | | | | |

