

HOMEWORK-2

1. DIFFERENTIATION OF IMPLICIT FUNCTION

1. If $y = \sqrt{\sin x + y}$, then $\frac{dy}{dx} =$

(A) $\frac{\sin x}{2y-1}$

(B) $\frac{\sin x}{1-2y}$

(C) $\frac{\cos x}{1-2y}$

(D) $\frac{\cos x}{2y-1}$

Ans. (D)

Sol. $y = \sqrt{\sin x + y}$

squaring both side

$$y^2 = \sin x + y$$

differentiating w.r.t. to x

$$2yy' = \cos x + y'$$

$$y' = \frac{\cos x}{2y-1}$$

2. If $ax^2 + 2hxy + by^2 = 0$, then $\frac{dy}{dx}$ equals-

(A) $\frac{ax+hy}{hx+by}$

(B) $-\frac{ax+hy}{hx+by}$

(C) $\frac{hx+by}{ax+hy}$

(D) $-\frac{hx+by}{ax+hy}$

Ans. (B)

Sol. $ax^2 + 2hxy + by^2 = 0$

$$2ax + 2hx \frac{dy}{dx} + 2hy \cdot 1 + 2by \frac{dy}{dx} = 0$$

$$\frac{dy}{dx}(hx + by) = -(ax + hy)$$

$$\frac{dy}{dx} = -\left(\frac{ax + hy}{hx + by}\right)$$

3. If $x\sqrt{y} + y\sqrt{x} = 1$, then $\frac{dy}{dx}$ equals-

(A) $-\frac{y+2\sqrt{xy}}{x+2\sqrt{xy}}$

(B) $-\sqrt{\frac{x}{y}} \left(\frac{y+2\sqrt{xy}}{x+2\sqrt{xy}}\right)$

(C) $-\sqrt{\frac{y}{x}} \left(\frac{y+2\sqrt{xy}}{x+2\sqrt{xy}}\right)$

(D) $\frac{x}{y}$

Ans. (C)

Sol. $x\sqrt{y} + y\sqrt{x} = 1$

$$x \cdot \frac{1}{2\sqrt{y}} \frac{dy}{dx} + \sqrt{y} \cdot 1 + y \frac{1}{2\sqrt{x}} + \sqrt{x} \frac{dy}{dx} = 0$$

$$\frac{dy}{dx} \left[\frac{x}{2\sqrt{y}} + \sqrt{x} \right] = - \left(\frac{y}{2\sqrt{x}} + \sqrt{y} \right)$$

$$\frac{dy}{dx} \left(\frac{x + 2\sqrt{xy}}{2\sqrt{y}} \right) = - \left(\frac{y + 2\sqrt{xy}}{2\sqrt{x}} \right)$$

$$\frac{dy}{dx} = - \frac{\sqrt{y}}{\sqrt{x}} \left(\frac{y + 2\sqrt{xy}}{x + 2\sqrt{xy}} \right)$$

4. If $e^x \sin y - e^y \cos x = 1$, then $\frac{dy}{dx}$ equals-

(A) $\frac{e^x \sin y + e^y \sin x}{e^y \cos x - e^x \cos y}$

(B) $\frac{e^x \sin y + e^y \sin x}{e^y \cos x + e^x \cos y}$

(C) $\frac{e^x \sin y - e^y \sin x}{e^y \cos x - e^x \cos y}$

(D) $\frac{e^x}{e^y}$

Ans. (A)

Sol. $e^x \sin y - e^y \cos x = 1$

$$e^x \cdot \cos y \frac{dy}{dx} + \sin y e^x - e^y (-\sin x) - \cos x.$$

$$e^y \cdot \frac{dy}{dx} = 0$$

$$\frac{dy}{dx} (e^x \cos y - e^y \cos x) = -(e^x \sin y + e^y \sin x)$$

$$\frac{dy}{dx} = - \left(\frac{e^x \sin y + e^y \sin x}{e^x \cos y - e^y \cos x} \right)$$

$$= \frac{e^x \sin y + e^y \sin x}{e^y \cos x - e^x \cos y}$$

5. If $\sqrt{1-x^2} + \sqrt{1-y^2} = a(x-y)$, then the value of dy/dx is -

(A) $\frac{\sqrt{1-x^2}}{\sqrt{1-y^2}}$

(B) $\frac{\sqrt{1-y^2}}{\sqrt{1-x^2}}$

(C) $-\frac{\sqrt{1-x^2}}{\sqrt{1-y^2}}$

(D) $-\frac{\sqrt{1-y^2}}{\sqrt{1-x^2}}$

Ans. (B)

Sol. $\frac{\sqrt{1-y^2}}{\sqrt{1-x^2}}$

Substituting $x = \sin \theta$ and $y = \sin \phi$ in the given equation, we get

$$\cos \theta + \cos \phi = a(\sin \theta - \sin \phi)$$

$$\Rightarrow 2 \cos \frac{\theta + \phi}{2} \cdot \cos \frac{\theta - \phi}{2} = 2a \cos \frac{\theta + \phi}{2} \cdot \sin \frac{\theta - \phi}{2}$$

$$\Rightarrow \cot \frac{\theta - \phi}{2} = a \Rightarrow \theta - \phi = 2 \cot^{-1} a$$

$$\Rightarrow \sin^{-1} x - \sin^{-1} y = 2 \cot^{-1} a$$

Differentiating with respect to x , we get

$$\frac{1}{\sqrt{1-x^2}} - \frac{1}{\sqrt{1-y^2}} \frac{dy}{dx} = 0 \Rightarrow \frac{dy}{dx} = \frac{\sqrt{1-y^2}}{\sqrt{1-x^2}}$$

6. If $x^3 \cos(xy) + y^3 \sin(xy) + 1 = 0$, then $\frac{dy}{dx}$ equals

(A) $\frac{x^3 y \tan(xy) - (3x^2 + y^4)}{xy^3 + (3y^2 - x^4) \tan xy}$

(B) $\frac{x^3 y \tan(xy) + (3x^2 + y^4)}{xy^3 - (3y^2 - x^4) \tan xy}$

(C) $\frac{x^3 y - (3x^2 + y^4) \tan(xy)}{xy^3 \tan(xy) + (3y^2 - x^4)}$

(D) $x^2 y \tan(xy) + 3x^2$

Ans. (A)

Sol. $f(x) = x^3 \cos xy + y^3 \sin xy + 1$

$$\frac{\partial f}{\partial y} = x^3 \cdot (-\sin xy) \cdot y + \cos xy \cdot 3x^2 + y^3 \cdot \cos xy \cdot y$$

$$= -x^3 y \sin xy + 3x^2 \cos xy + y^4 \cos xy$$

$$\& \frac{\partial f}{\partial y} = x^3 (-\sin xy) \cdot x + y \cos xy \cdot x + \sin xy \cdot 3y^2$$

$$= -x^4 \sin xy + y^3 x \cos xy + 3y^2 \sin xy$$

$$\therefore \frac{dy}{dx} = - \frac{\partial f / \partial x}{\partial f / \partial y}$$

$$= - \left(\frac{-x^3 \sin xy + 3x^2 \cos xy + y^4 \cos xy}{-x^4 \sin xy + y^3 x \cos xy + 3y^2 \sin xy} \right)$$

$$= - \left(\frac{-x^3 y \tan xy + 3x^2 + y^4}{-x^4 \tan xy + y^3 x + 3y^2 \tan xy} \right)$$

$$= \frac{x^3 y \tan xy - (3x^2 + y^4)}{y^3 x + (3y^2 - x^4) \tan xy}$$

7. If $y = 2^{\log_2 x^{2x}} + \left(\tan \frac{\pi x}{4} \right)^{\frac{4}{\pi x}}$ then $\left. \frac{dy}{dx} \right|_{x=1}$ is

(A) 4

(B) 1

(C) 0

(D) Not defined

Ans. (A)

Sol. $y = x^{2x} + \left(\tan \left(\frac{\pi x}{4} \right) \right)^{4/\pi x}$

$$y = u + v$$

$$\Rightarrow \frac{dy}{dx} = \frac{du}{dx} + \frac{dv}{dx}$$

$$u = x^{2x}, v = \left(\tan \left(\frac{\pi x}{4} \right) \right)^{4/\pi x}$$

2. DIFFERENTIATION OF LOGARITHMIC FUNCTION

8. If $x^m \cdot y^n = (x + y)^{m+n}$, then $\frac{dy}{dx}$ is

(A) $\frac{x+y}{xy}$

(B) xy

(C) $\frac{x}{y}$

(D) $\frac{y}{x}$

Ans. (D)

Sol. Given that, $x^m y^n (x + y)^{m+n}$

Taking log on both sides, we get

$$m \log x + n \log y = (m + n) \log(x + y)$$

On differentiating w.r.t. x , we get

$$\frac{m}{x} + \frac{n}{y} \frac{dy}{dx} = \frac{(m+n)}{(x+y)} \left(1 + \frac{dy}{dx} \right)$$

$$\Rightarrow \frac{dy}{dx} \left(\frac{m+n}{x+y} - \frac{n}{y} \right) = \frac{m}{x} - \frac{m+n}{x+y}$$

$$\Rightarrow \frac{dy}{dx} \left(\frac{my + ny - nx - ny}{y(x+y)} \right)$$

$$= \frac{mx + my - mx - nx}{x(x+y)} \Rightarrow \frac{dy}{dx} = \frac{y}{x}$$

9. If $f(x) = |x|^{\sin x}$ then $f'(\pi/4)$ equals

(A) $\left(\frac{\pi}{4} \right)^{1/\sqrt{2}} \left(\frac{\sqrt{2}}{2} \ln \frac{4}{\pi} - \frac{2\sqrt{2}}{\pi} \right)$

(B) $\left(\frac{\pi}{4} \right)^{1/\sqrt{2}} \left(\frac{\sqrt{2}}{2} \ln \frac{4}{\pi} + \frac{2\sqrt{2}}{\pi} \right)$

(C) $\left(\frac{\pi}{4} \right)^{1/\sqrt{2}} \left(\frac{\sqrt{2}}{2} \ln \frac{\pi}{4} - \frac{2\sqrt{2}}{\pi} \right)$

(D) $\left(\frac{\pi}{4} \right)^{1/\sqrt{2}} \left(\frac{\sqrt{2}}{2} \ln \frac{\pi}{4} + \frac{2\sqrt{2}}{\pi} \right)$

Ans. (D)

Sol. $f(x) = |x|^{\sin x}$

$$x > 0, f(x) = x^{\sin x} \Rightarrow y = x^{\sin x}$$

$$\ell ny = \sin x \ln x$$

$$y' = x^{\sin x} \left\{ \frac{\sin x}{x} + \cos x \ln x \right\}$$

$$y|_{x=\pi/4} = \left(\frac{\pi}{4} \right)^{\frac{1}{\sqrt{2}}} \left\{ \frac{1}{\sqrt{2}} \times \frac{4}{\pi} + \frac{1}{\sqrt{2}} \ln \frac{\pi}{4} \right\}$$

$$x < 0 \quad y = -x^{-\sin x}$$

$$\ln y = -\sin x \ln(-x)$$

10. If $y = (1+x)(1+x^2)(1+x^4) \dots (1+x^{2n})$, then $\frac{dy}{dx}$ at $x = 0$ is
 (A) -1 (B) 1 (C) 0 (D) 2^n

Ans. (B)

Sol. $y = (1+x)(1+x^2) \dots (1+x^{2n})$

$$y = \frac{(1-x^2)(1+x^2)(1+x^4) \dots (1+x^{2n})}{(1-x)}$$

$$y = \frac{1-x^{4n}}{1-x}$$

$$\frac{dy}{dx} = \frac{(1-x)(-4nx^{4n-1}) + (1-x^{4n})}{(1-x)^2}$$

$$\frac{dy}{dx} = \frac{-4nx^{4n-1} + 4nx^{4n} + 1 - x^{4n}}{(1-x)^2}$$

$$\left. \frac{dy}{dx} \right|_{x=0} = \frac{-4n \times 0 + 0 + 1 - 0}{1} = 1$$

3. DIFFERENTIATION OF INFINITE SERIES

11. If $x = e^{y+e^y+\dots \text{upto } \infty}$, $x > 0$, then $\frac{dy}{dx}$
 (A) $\frac{x}{1+x}$ (B) $\frac{1}{x}$ (C) $\frac{1-x}{x}$ (D) $\frac{1+x}{x}$

Ans. (C)

Sol. $x = e^y + e^{y+e^y+\dots \infty}$

$$x = e^{y+x}$$

$$x = e^{(y+x)} \left\{ \frac{dy}{dx} + 1 \right\}$$

$$\frac{dy}{dx} = \frac{1 - e^{x+y}}{e^{x+y}} = \frac{1-x}{x}$$

12. If $y = \sqrt{x}^{\sqrt{x}^{\sqrt{x}^{\dots \infty}}}$, then the value of $\frac{dy}{dx}$ is
 (A) $\frac{xy^2}{2-y \log x}$ (B) $\frac{x^2}{y(2-y \log x)}$ (C) $\frac{y^2}{x(2-y \log x)}$ (D) $\frac{y^2}{x(2+y \log x)}$

6

(MATHEMATICS)

METHOD OF DIFFERENTIATION

14. Let $f(x) = x + \frac{1}{2x + \frac{1}{2x + \frac{1}{2x + \dots \infty}}}$. Compute the value of $f(100) \cdot f'(100)$

Ans. 100

Sol. $2y = \frac{1}{2x + (y-x)} + y - x + 2x$

$$y = \frac{1}{y+x} + x$$

$$y^2 - x^2 = 0 \Rightarrow yy' - x = 0$$

$$f(100) \cdot f'(100) = 100$$

15. If $y = x + \frac{1}{x + \frac{1}{x + \frac{1}{x + \dots}}}$, prove that $\frac{dy}{dx} = \frac{1}{2 - \frac{x}{x + \frac{1}{x + \dots}}}$

Sol. $y = x + \frac{1}{x + \frac{1}{x + \dots}}$

$$y = x + \frac{1}{y}$$

$$y^2 = xy + 1$$

$$2yy' = xy' + y \Rightarrow y' = \frac{y}{2y - x}$$

$$y' = \frac{1}{2 - \frac{x}{y}} \quad y' = \frac{1}{2 - \frac{x}{x + \frac{1}{x + \dots}}}$$

16. If $y = \tan^{-1} \frac{1}{x^2+x+1} + \tan^{-1} \frac{1}{x^2+3x+3} + \tan^{-1} \frac{1}{x^2+5x+7} + \tan^{-1} \frac{1}{x^2+7x+13} + \dots$ to n terms.

Find dy/dx , expressing your answer in 2 terms,

Ans. $\frac{1}{1+(x+n)^2} - \frac{1}{1+x^2}$

Sol. $y = \tan^{-1}(x+1) - \tan^{-1}x$

$$+ \tan^{-1}(x+2) - \tan^{-1}(x+1)$$

$$+ \tan^{-1}(x+n) - \tan^{-1}(x+(n-1))$$

$$y = \tan^{-1}(x+n) - \tan^{-1}x$$

$$\Rightarrow \frac{dy}{dx} = \frac{1}{1+(x+n)^2} - \frac{1}{1+x^2}$$

4. DIFFERENTIATION OF DETERMINANT

17. If $f(x), g(x), h(x)$ are polynomials in x of degree 2

and $F(x) = \begin{vmatrix} f & g & h \\ f' & g' & h' \\ f'' & g'' & h'' \end{vmatrix}$, then $F'(x)$ is equal to

- (A) 1 (B) 0 (C) -1 (D) $f(x) \cdot g(x) \cdot h(x)$

Ans. (B)

Sol. $F'(x) = \begin{vmatrix} f' & g' & h' \\ f'' & g'' & h'' \\ f''' & g''' & h''' \end{vmatrix} + \begin{vmatrix} f & g & h \\ f'' & g'' & h'' \\ f''' & g''' & h''' \end{vmatrix} + \begin{vmatrix} f & g & h \\ f' & g' & h' \\ f''' & g''' & h''' \end{vmatrix} = 0$

18. If $y = \sin mx$ then the value of $\begin{vmatrix} y & y_1 & y_2 \\ y_3 & y_4 & y_5 \\ y_6 & y_7 & y_8 \end{vmatrix}$

(where subscripts of y shows the order of derivative) is

- (A) independent of x but dependent on m (B) dependent of x but independent of m
(C) dependent on both m & x (D) independent of m & x

Ans. (D)

Sol. $\begin{vmatrix} \sin mx & m \cos mx & -m^2 \sin mx \\ -m^3 \cos mx & m^4 \sin mx & m^5 \cos mx \\ -m^6 \sin mx & -m^7 \cos mx & m^8 \sin mx \end{vmatrix}$

by expanding $= 0$

19. If $f(x) = \begin{vmatrix} \cos(x+x^2) & \sin(x+x^2) & -\cos(x+x^2) \\ \sin(x-x^2) & \cos(x-x^2) & \sin(x-x^2) \\ \sin 2x & 0 & \sin 2x^2 \end{vmatrix}$ then

- (A) $f(-2) = 0$ (B) $f'(-1/2) = 0$
(C) $f'(-1) = -2$ (D) $f''(0) = 4$

Ans. (B, C, D)

Sol. $f(x) = \begin{vmatrix} \cos(x+x^2) & \sin(x+x^2) & -\cos(x+x^2) \\ \sin(x-x^2) & \cos(x-x^2) & \sin(x-x^2) \\ \sin 2x & 0 & \sin 2x^2 \end{vmatrix}$

5. HIGHER ORDER DERIVATIVE

20. If $y = a \cos(\ln x) + b \sin(\ln x)$, then $x^2 \frac{d^2 y}{dx^2} + x \frac{dy}{dx}$

- (A) 0 (B) y (C) $-y$ (D) xy

Ans. (C)

Sol. $y = a \cos \ell n x + b \sin \ell n x$

differentiating w.r.t. to x

$$y' = -\frac{a}{x} \sin \ell n x + \frac{b}{x} \cos \ell n x$$

$$xy' = -a \sin \ell n x + b \cos \ell n x$$

$$xy'' + y' = -\frac{a \cos \ell n x}{x} - \frac{b \sin \ell n x}{x}$$

$$x^2 y'' + xy' = -y$$

21. $\frac{d^2 x}{dy^2}$ equals :

(A) $\left(\frac{d^2 y}{dx^2}\right)^{-1}$

(B) $\left(\frac{d^2 y}{dx^2}\right)^{-1} \left(\frac{dy}{dx}\right)^{-3}$

(C) $\left(\frac{d^2 y}{dx^2}\right) \left(\frac{dy}{dx}\right)^{-2}$

(D) $\left(\frac{d^2 y}{dx^2}\right) \left(\frac{dy}{dx}\right)^{-3}$

Ans. (D)

Sol. Here, $\frac{dy}{dx} = \left(\frac{dy}{dx}\right)^{-1}$

Differentiating both sides w.r.t. y , we get

$$\frac{d^2 x}{dy^2} = \left(\frac{dy}{dx}\right)^{-2} \cdot \frac{d}{dy} \cdot \left(\frac{dy}{dx}\right)$$

$$= -\left(\frac{dy}{dx}\right)^{-2} \cdot \frac{d}{dy} \left(\frac{dy}{dx}\right) \cdot \frac{dx}{dx}$$

$$= -\left(\frac{dy}{dx}\right)^{-2} \cdot \frac{d}{dx} \cdot \left(\frac{dy}{dx}\right) \cdot \frac{dx}{dy}$$

$$= -\left(\frac{dy}{dx}\right)^{-2} \cdot \frac{d^2 y}{dx^2} \cdot \left(\frac{dy}{dx}\right)^{-1}$$

$$= -\left(\frac{dy}{dx}\right)^{-3} \cdot \left(\frac{d^2 y}{dx^2}\right)$$

6. TRIGONOMETRIC SUBSTITUTIONS

22. The derivative of $\sec^{-1} \left(\frac{1}{2x^2-1}\right)$ w.r.t. $\sqrt{1-x^2}$ at $x = \frac{1}{2}$ is

(A) 4

(B) $1/4$

(C) 1

(D) 7

Ans. (A)

Sol. Let $u = \sec^{-1} \left(\frac{1}{2x^2-1}\right)$

$$= \cos^{-1}(2x^2 - 1) \text{ Let } x = \cos \theta$$

$$= \cos^{-1}(2\cos^2\theta - 1)$$

$$= \cos^{-1}(\cos 2\theta)$$

$$u = 2\theta \Rightarrow u = 2\cos^{-1}x$$

$$\Rightarrow \frac{du}{dx} = 2 \times \frac{-1}{\sqrt{1-x^2}} = \frac{-2}{\sqrt{1-x^2}}$$

$$\text{Let } v = \sqrt{1-x^2}$$

$$\Rightarrow \frac{dv}{dx} = \frac{1}{2\sqrt{1-x^2}} \cdot (-2x) = \frac{-x}{\sqrt{1-x^2}}$$

$$\therefore \frac{du/dx}{dv/dx} = \frac{2/\sqrt{1-x^2}}{x/\sqrt{1-x^2}} = \frac{2}{x}$$

$$\Rightarrow \frac{du}{dv} = \frac{2}{x} \Rightarrow \left(\frac{du}{dv}\right) = \frac{2}{1/2} = 2 \times 2 = 4$$

$$\therefore \left(\frac{du}{dv}\right)_{x=1/2} = 4$$

23. If $y = \sin^{-1} \frac{2x}{1+x^2}$ then $\left.\frac{dy}{dx}\right|_{x=-2}$ is

(A) $\frac{2}{5}$

(B) $\frac{2}{\sqrt{5}}$


(C) $-\frac{2}{5}$

(D) $\frac{\sqrt{5}}{2}$

Ans. (C)


Sol. $y = \sin^{-1} \left(\frac{2x}{1+x^2} \right) \cdot \left.\frac{dy}{dx}\right|_{x=-2}$

$$x = \tan\theta \Rightarrow y = \sin^{-1}(\sin 2\theta)$$



$$\theta < -\frac{\pi}{4}$$

$$2\theta < -\frac{\pi}{2}$$



$$y = \pi - 2\theta = \pi - 2\tan^{-1}x$$

$$\left. \frac{dy}{dx} \right|_{x=-2} = \frac{-2}{(1+x^2)} = \frac{-2}{5}$$

24. $\frac{d}{dx} \left[\tan^{-1} \left(\frac{\sqrt{x^2+a^2}+x}{\sqrt{x^2+a^2}-x} \right)^{1/2} \right]$, is equal to -

(A) $\frac{a}{2(x^2+a^2)}$

(B) $\frac{a}{x^2+a^2}$

(C) $1/2$

(D) None of these

Ans. (A)

Sol. $y = \tan^{-1} \left(\frac{\sqrt{a^2+x^2}+x}{\sqrt{a^2+x^2}-x} \right)^{1/2}$

put $x = a \tan \theta$

$$= \tan^{-1} \left(\frac{\sqrt{a^2 + a^2 \tan^2 \theta} + a \tan \theta}{\sqrt{a^2 + a^2 \tan^2 \theta} - a \tan \theta} \right)^{1/2}$$

$$= \tan^{-1} \left(\frac{\sec \theta + \tan \theta}{\sec \theta - \tan \theta} \right)^{1/2}$$

$$= \tan^{-1} \left(\frac{1 + \sin \theta}{1 - \sin \theta} \right)^{1/2}$$

$$= \tan^{-1} \left\{ \frac{\cos \frac{\theta}{2} + \sin \frac{\theta}{2}}{\cos \frac{\theta}{2} - \sin \frac{\theta}{2}} \right\}^{-1/2}$$

$$= \tan^{-1} \left(\frac{1 + \tan \theta/2}{1 - \tan \theta/2} \right) = \tan^{-1} \tan(\pi/4 + \theta/2)$$

$$= \pi/4 + \theta/2$$

$$y = \frac{\pi}{4} + \frac{1}{2} \tan^{-1} \frac{x}{a}$$

$$\frac{dy}{dx} = 0 + \frac{1}{2} \times \frac{1}{1+\frac{x^2}{a^2}} \times \frac{1}{a} = \frac{fa^2}{2(x^2+a^2)}$$

25. $\frac{d}{dx} \left[\sin^2 \cot^{-1} \frac{1}{\sqrt{\frac{1+x}{1-x}}} \right]$, is equal to -

(A) 0

(B) 1/2

(C) -1/2

(D) -1]

Ans. (B)

Sol. $y = \sin^2 \cot^{-1} \frac{1}{\sqrt{\frac{1+x}{1-x}}}$

$$= \sin^2 \tan^{-1} \sqrt{\frac{1+x}{1-x}} \left\{ \because \tan^{-1} x = \sin^{-1} \frac{x}{\sqrt{1+x}} \right\}$$

$$= \sin^2 \sin^{-1} \frac{\sqrt{\frac{1+x}{1-x}}}{\sqrt{1+x}} = \sin^2 \sin^{-1} \frac{\sqrt{1+x}}{\sqrt{2}}$$

$$= \left(\sin^{-1} \sin^{-1} \frac{\sqrt{1+x}}{\sqrt{2}} \right)^2 = \left(\frac{\sqrt{1+x}}{\sqrt{2}} \right)^2$$

$$\frac{dy}{dx} = 0 + \frac{1}{2}$$

26. If $y = \tan^{-1} \frac{u}{\sqrt{1-u^2}}$ & $x = \sec^{-1} \frac{1}{2u^2-1}$

$u \in \left(0, \frac{1}{\sqrt{2}}\right) \cup \left(\frac{1}{\sqrt{2}}, 1\right)$ prove that $2 \frac{dy}{dx} + 1 = 0$

Sol. $y = \tan^{-1} \frac{u}{\sqrt{1-u^2}}$ & $x = \sec^{-1} \frac{1}{\sqrt{2u^2-1}}$

$u \in \left(0, \frac{1}{\sqrt{2}}\right) \cup \left(\frac{1}{\sqrt{2}}, 1\right)$

$u = \sin \theta \quad 0 < u < \frac{1}{\sqrt{2}} \text{ OR } \frac{1}{\sqrt{2}} < u < 1$

$\left(\frac{\pi}{4}, \frac{\pi}{2}\right) \text{ OR } \left(0, \frac{\pi}{4}\right) \rightarrow 1^{\text{st}} \text{ Q}^{\text{nt}}$

$y = \tan^{-1} \left(\frac{\sin \theta}{\cos \theta} \right) = \theta$

$x = \sec^{-1} = \sec^{-1} \left(-\frac{1}{\cos 2\theta} \right)$

$\Rightarrow \sec^{-1}(-\sec 2\theta)$

$x = \pi - 2\theta$

$\frac{dy}{dq} = 1, \frac{dx}{dq} = -2$

$\frac{dy}{dx} = -\frac{1}{2} \Rightarrow 2 \cdot \frac{dy}{dx} + 1 = 0$

7. MIXED PROBLEMS

27. If $y = \sec(\tan^{-1}x)$, then $\frac{dy}{dx}$ at $x = 1$ is equal to: [JEE Main 2013]

- (A) 1 (B) $\sqrt{2}$ (C) $\frac{1}{\sqrt{2}}$ (D) $\frac{1}{2}$

Ans. (C)

Sol. $y^{-1} = \sec(\tan^{-1}x) \tan(\tan^{-1}x) \times \frac{1}{1+x^2}$

at $x = 1$

$$y' = \sqrt{2} \times \frac{1}{2} = \frac{1}{\sqrt{2}}$$

28. For $x \in \mathbb{R}$, $f(x) = |\log 2 - \sin x|$ and $g(x) = f(f(x))$, then: [JEE MAIN 2016]

- (A) $g'(0) = \cos(\log 2)$
 (B) $g'(0) = -\cos(\log 2)$
 (C) g is differentiable at $x = 0$ and $g'(0) = -\sin(\log 2)$
 (D) g is not differentiable at $x = 0$

Ans. (A)

Sol. We have, $f(x) = |\log 2 - \sin x|$
 and $g(x) = f(f(x))$, $x \in \mathbb{R}$
 Note that, for $x \rightarrow 0$, $\log 2 > \sin x$

$$f(x) = \log 2 - \sin x$$

$$\Rightarrow g(x) = \log 2 - \sin(f(x))$$

$$= \log 2 - \sin(\log 2 - \sin x)$$

Clearly, $g(x)$ is differentiable at $x = 0$ as $\sin x$ is differentiable.

$$\text{Now, } g'(x) = -\cos(\log 2 - \sin x)(-\cos x)$$

$$= \cos x \cdot \cos(\log 2 - \sin x)$$

$$\Rightarrow g'(0) = 1 \cdot \cos(\log 2)$$

29. If for $x \in (0, \frac{1}{4})$, then derivative of $\tan^{-1} \left(\frac{6x\sqrt{x}}{1-9x^3} \right)$ is $\sqrt{x} \cdot g(x)$, then $g(x)$ equals

- (A) $\frac{9}{1+9x^3}$ (B) $\frac{3x\sqrt{x}}{1-9x^3}$ [JEE MAIN 2017]
 (C) $\frac{3x}{1-9x^3}$ (D) $\frac{3x}{1+9x^3}$

Ans. (A)

Sol. Let $y = \tan^{-1} \left(\frac{6x\sqrt{x}}{1-9x^3} \right) = \tan^{-1} \left[\frac{2 \cdot (3x^{3/2})}{1-(3x^{3/2})^2} \right]$

$$= 2 \tan^{-1}(3x^{3/2}) \left[\because 2 \tan^{-1} x = \tan^{-1} \frac{2x}{1-x^2} \right]$$

$$\frac{dy}{dx} = 2 \cdot \frac{1}{1+(3x^{3/2})^2} \cdot 3 \times \frac{3}{2} (x)^{1/2}$$

$$= \frac{9}{1+9x^3} \cdot \sqrt{x}$$

$$\therefore g(x) = \frac{9}{1+9x^3}$$

30. Let $f(\theta) = \sin \left(\tan^{-1} \left(\frac{\sin \theta}{\sqrt{\cos 2\theta}} \right) \right)$ where $-\frac{\pi}{4} < \theta < \frac{\pi}{4}$, Then the value of $\frac{d}{d(\tan \theta)} (f(\theta))$ is

[JEE 2011]

Ans. 1

Sol. Given $f(\theta) = \sin \left(\tan^{-1} \left(\frac{\sin \theta}{\sqrt{\cos 2\theta}} \right) \right)$

$$\text{Let } \tan^{-1} \left(\frac{\sin \theta}{\sqrt{\cos 2\theta}} \right) = y$$

$$\Rightarrow \tan y = \frac{\sin \theta}{\sqrt{2\cos^2 \theta - 1}}$$

Refer to the figure below,

$$\Rightarrow \sin y = \frac{\sin \theta}{\cos \theta}$$

$$f(\theta) = \sin y = \frac{\sin \theta}{\cos \theta} = \tan \theta$$

$$\Rightarrow \frac{d}{d(\tan \theta)} (f(\theta)) = 1$$

ANSWER KEY

1. DIFFERENTIATION OF IMPLICIT FUNCTION

1. (D) 2. (B) 3. (C) 4. (A) 5. (B) 6. (A) 7. (A)

2. DIFFERENTIATION OF LOGARITHMIC FUNCTION

8. (D) 9. (D) 10. (B)

3. DIFFERENTIATION OF INFINITE SERIES

11. (C) 12. (C) 13. (B) 14. 100 16. $\frac{1}{1+(x+n)^2} - \frac{1}{1+x^2}$

4. DIFFERENTIATION OF DETERMINANT

17. (B) 18. (D) 19. (B,C,D)

5. HIGHER ORDER DERIVATIVE

20. (C) 21. (D)

6. TRIGONOMETRIC SUBSTITUTIONS

22. (A) 23. (C) 24. (A) 25. (B)

7. MIXED PROBLEMS

27. (C) 28. (A) 29. (A) 30. 1