

HOMework-3

DIFFERENTIATION OF A FUNCTION RESPECT TO ANOTHER FUNCTION

- If $x = e^{\sin^{-1}t}$, $y = \tan^{-1}t$, then $\frac{dy}{dx} =$
 (A) $\frac{1}{1+t^2} e^{-\sin^{-1}t} \sqrt{1-t^2}$ (B) $\frac{1}{1+t^2} e^{-\sin^{-1}t}$
 (C) $(1+t^2)e^{-\sin^{-1}t} \sqrt{1-t^2}$ (D) None of these
- Find derivative of $f(\tan x)$ w.r.t. $g(\sec x)$ at $x = \frac{\pi}{4}$
 where $f'(1) = 2$ and $g'(\sqrt{2}) = 4$ is
 (A) 3 (B) -4 (C) $\frac{2}{19}$ (D) $\frac{1}{\sqrt{2}}$
- Differential coefficient of $\sin^{-1}x$ with respect to $3 \cdot \sin^{-1}(3x - 4x^3)$ is
 (A) $\frac{1}{3}$ if $-\frac{\pi}{8} < x < \frac{\pi}{8}$ (B) 3 if $-\frac{\pi}{8} < x < \frac{\pi}{8}$
 (C) $\frac{1}{3}$ if $-\frac{\pi}{9} < x < \frac{\pi}{9}$ (D) $\frac{1}{3}$ if $-\frac{1}{2} < x < \frac{1}{2}$
- The differential coefficient of $\sin^{-1} \frac{t}{\sqrt{1+t^2}}$ w.r.t. $\cos^{-1} \frac{1}{\sqrt{1+t^2}}$ is
 (A) $1 \forall t > 0$ (B) $-1 \forall t < 0$ (C) $1 \forall t \in \mathbb{R}$ (D) $2 \forall t > 0$

LOGARITHMIC FUNCTION/TRIGONOMETRIC SUBSTITUTIONS

- $y = \cos^{-1} \sqrt{\frac{\sqrt{1+x^2}+1}{2\sqrt{1+x^2}}}$ then $\frac{dy}{dx}$ is
 (A) $\frac{1}{2(1+x^2)}$, $x \in \mathbb{R}$ (B) $\frac{1}{2(1+x^2)}$, $x > 0$
 (C) $\frac{-1}{2(1+x^2)}$, $x < 0$ (D) $\frac{1}{2(1+x^2)}$, $x < 0$

INFINITE SERIES

- If $y = \sqrt{x + \sqrt{y + \sqrt{x + \dots}}}$, then $\frac{dy}{dx} =$
 (A) $\frac{x-y^2}{2y^3-2xy-1}$ (B) $\frac{x-y^2}{2y^3-2xy-1}$ (C) $\frac{x+y^2}{2y^3-2xy-1}$ (D) $\frac{y^2-x}{(y^2-x)^3-1}$

DIFFERENTIATION OF PARAMETRIC EQUATIONS

- If $x = \frac{1+t}{t^3}$, $y = \frac{3}{2t^2} + \frac{2}{t}$ then, $x \left(\frac{dy}{dx} \right)^3 - \frac{dy}{dx} =$
 (A) 0 (B) -1 (C) 1 (D) 2
- If $\sin x = \frac{2t}{1+t^2}$ and $\cot y = \frac{1-t^2}{2t}$. Then value of $\frac{d^2x}{dy^2}$ is equal to
 (A) 0 (B) 1 (C) -1 (D) $\frac{1}{2}$

MIXED PROBLEMS

9. If $y = \tan^{-1} \left(\frac{\ln \frac{e}{x^2}}{\ln ex^2} \right) + \tan^{-1} \frac{3+2\ln x}{1-6\ln x}$ then
- (A) $\frac{dy}{dx} = 0$ (B) $\frac{d^2y}{dx^2} = 0$ (C) $\frac{dy}{dx} = \frac{2}{x(1+\ln^2 x)}$ (D) $\frac{dy}{dx} = 1$

JEE MAIN

10. If $x^m \cdot y^n = (x+y)^{m+n}$, then $\frac{dy}{dx}$ is - [AIEEE-2006]
- (A) $\frac{x+y}{xy}$ (B) xy (C) $\frac{x}{y}$ (D) $\frac{y}{x}$
11. Let y be an implicit function of x defined by $x^{2x} - 2x^x \cot y - 1 = 0$. then $y'(1)$ equals : [AIEEE-2009]
- (A) $\log 2$ (B) $-\log 2$ (C) -1 (D) 1
12. Let $f: (-1,1) \rightarrow \mathbb{R}$ be a differentiable function with $f(0) = -1$ and $f'(0) = 1$. Let $g(x) = [f(2f(x) + 2)]^2$. Then $g'(0)$:- [AIEEE-2010]
- (A) 4 (B) -4 (C) 0 (D) -2
13. $\frac{d^2x}{dy}$ equals :- [AIEEE-2011]
- (A) $\left(\frac{d^2y}{dx^2} \right) \left(\frac{dy}{dx} \right)^{-2}$ (B) $-\left(\frac{d^2y}{dx^2} \right) \left(\frac{dy}{dx} \right)^{-3}$
- (C) $\left(\frac{d^2y}{dx^2} \right)^{-1}$ (D) $-\left(\frac{d^2y}{dx^2} \right)^{-1} \left(\frac{dy}{dx} \right)^{-3}$
14. If $y = \sec(\tan^{-1} x)$, then $\frac{dy}{dx}$ at $x = 1$ is equal to : [JEE-MAIN-2013]
- (A) $\frac{1}{\sqrt{2}}$ (B) $\frac{1}{2}$ (C) 1 (D) $\sqrt{2}$
15. If g is the inverse of a function f and $f'(x) = \frac{1}{1+x^5}$, then $g'(x)$ is equal to : [JEE-MAIN-2014]
- (A) $1 + x^5$ (B) $5x^4$ (C) $\frac{1}{1+\{g(x)\}^5}$ (D) $1 + \{g(x)\}^5$
16. If for $x \in \mathbb{R}$, $f(x) = |\log 2 - \sin x|$ and $g(x) = f(f(x))$, then : [JEE(Main)-2016]
- (A) g is differentiable at $x = 0$ and $g'(0) = -\sin(\log 2)$
- (B) g is not differentiable at $x = 0$
- (C) $g'(0) = \cos(\log 2)$
- (D) $g'(0) = -\cos(\log 2)$
17. If for $x \in \left(0, \frac{1}{4}\right)$, the derivative of $\tan^{-1} \left(\frac{6x\sqrt{x}}{1-9x^3} \right)$ is $\sqrt{x} \cdot g(x)$ then $g(x)$ equals : [JEE (Main)2017]
- (A) $\frac{3}{1+9x^3}$ (B) $\frac{9}{1+9x^3}$ (C) $\frac{3x\sqrt{x}}{1-9x^3}$ (D) $\frac{3x}{1-9x^3}$

(MATHEMATICS)

METHOD OF DIFFERENTIATION

18. If $x = 3 \tan t$ and $y = 3 \sec t$, then the value of $\frac{d^2y}{dx^2}$ at $t = \frac{\pi}{4}$, is : [JEE (Main)2019]
 (A) $\frac{3}{2\sqrt{2}}$ (B) $\frac{1}{6\sqrt{2}}$ (C) $\frac{1}{3\sqrt{2}}$ (D) $\frac{1}{6}$
19. Let $f: \mathbb{R} \rightarrow \mathbb{R}$ be a function such that $f(x) = x^3 + x^2 f'(A) + x f''(B) + f f'''(C)$, $x \in \mathbb{R}$. Then $f(B)$ equals: [JEE (Main)2019]
 (A) 8 (B) 30 (C) -4 (D) -2
20. If $x \log_e(\log_e x) - x^2 + y^2 = 4$ ($y > 0$), then $\frac{dy}{dx}$ at $x = e$ is equal to : [JEE (Main)2019]
 (A) $\frac{e}{\sqrt{4+e^2}}$ (B) $\frac{(2e-1)}{2\sqrt{4+e^2}}$ (C) $\frac{(1+2e)}{2\sqrt{4+e^2}}$ (D) $\frac{(1+2e)}{\sqrt{4+e^2}}$
21. For $x > 1$, if $(2x)^{2y} = 4e^{2x-2y}$, then $(1 + \log_e 2x)^2 \frac{dx}{dy}$ is equal to : [JEE (Main)2019]
 (A) $\frac{x \log_e 2x - \log_e 2}{x}$ (B) $x \log_e 2x$
 (C) $\log_e 2x$ (D) $\frac{x \log_e 2x + \log_e 2}{x}$
22. If $2y = \left(\cot^{-1} \left(\frac{\sqrt{3} \cos x + \sin x}{\cos x - \sqrt{3} \sin x} \right) \right)^2$, $x \in \left(0, \frac{\pi}{2} \right)$, then $\frac{dy}{dx}$ is equal to [JEE (Main)2019]
 (A) $\frac{\pi}{6} - x$ (B) $x - \frac{\pi}{6}$ (C) $2x - \frac{\pi}{3}$ (D) $\frac{\pi}{3} - x$
23. The derivative of $\tan^{-1} \left(\frac{\sin x - \cos x}{\sin x + \cos x} \right)$, with respect to $\frac{x}{2}$, where $\left(x \in \left(0, \frac{\pi}{2} \right) \right)$ is : [JEE (Main)2019]
 (A) 2 (B) $\frac{2}{3}$ (C) 1 (D) $\frac{1}{2}$
24. Let $(x)^k + (y)^k = (a)^k$ where $a, k > 0$ and $\frac{dy}{dx} + \left(\frac{y}{x} \right)^{\frac{1}{3}} = 0$, then find k - [JEE (Main)2020]
 (A) $\frac{1}{3}$ (B) $\frac{2}{3}$ (C) $\frac{4}{3}$ (D) 2
25. If $y^{14} + y^{-14} = 2x$, and $(x^2 - 1) \frac{d^2y}{dx^2} + \alpha x \frac{dy}{dx} + \beta y = 0$ then $|\alpha - \beta|$ is equal to . [JEE (Main)2021]
26. If $y(x) = (x^{x^x})$, $x > 0$ then $\frac{d^2x}{dy^2} + 20$ at $x = 1$ is equal to: [JEE (Main)2022]
27. Let $y(x) = (1+x)(1+x^2)(1+x^4)(1+x^8)(1+x^{16})$ [JEE (Main)2023]
 Then $y' - y''$ at $x = -1$ is equal to
 (A) 976 (B) 464 (C) 496 (D) 944

JEE ADVANCED

SECTION-1

28. For $x > 0$, $\lim_{x \rightarrow 0} ((\sin x)^{1/x} + (1/x)^{\sin x})$ is :- [JEE 2006, 3]
 (A) 0 (B) -1 (C) 1 (D) 2
29. $\frac{d^2x}{dy^2}$ equals :-
 (A) $\left(\frac{d^2y}{dx^2}\right)^{-1}$ (B) $-\left(\frac{d^2y}{dx^2}\right)^{-1} \left(\frac{dy}{dx}\right)^{-3}$ (C) $\left(\frac{d^2y}{dx^2}\right) \left(\frac{dy}{dx}\right)^{-2}$ (D) $-\left(\frac{d^2y}{dx^2}\right) \left(\frac{dy}{dx}\right)^{-3}$
30. (a) Let $g(x) = \ln f(x)$ where $f(x)$ is a twice differentiable positive function on $(0, \infty)$ such that $f(x+1) = xf(x)$. then for $N = 1, 2, 3, \dots$, $g''\left(N + \frac{1}{2}\right) - g''\left(\frac{1}{2}\right) =$
 (A) $-4 \left\{1 + \frac{1}{9} + \frac{1}{25} + \dots \cdot \frac{1}{(2N-1)^2}\right\}$ (B) $4 \left\{1 + \frac{1}{9} + \frac{1}{25} + \dots \cdot \frac{1}{(2N-1)^2}\right\}$
 (C) $-4 \left\{1 + \frac{1}{9} + \frac{1}{25} + \dots \cdot \frac{1}{(2N+1)^2}\right\}$ (D) $4 \left\{1 + \frac{1}{9} + \frac{1}{25} + \dots \cdot \frac{1}{(2N+1)^2}\right\}$
- (b) Let f and g be real valued functions defined on interval $(-1, 1)$ such that $g(x)$ is continuous, $g(0) \neq 0$, $g'(0) = 0$, $g''(0) \neq 0$, and $f(x) = g(x)\sin x$.
- Statement-1 :** $\lim_{x \rightarrow 0} [g(x)\cot x - g(0)\operatorname{cosec} x] = f''(0)$
 And
Statement-2 : $f'(0) = g(0)$
- (A) Statement-1 is true, statement-2 is true and statement-2 is correct explanation of statement-1.
 (B) Statement-1 is true, statement-2 is true and statement-2 is NOT the correct explanation for statement-1
 (C) Statement-1 is true, statement-2 is false.
 (D) Statement-1 is false, statement-2 is true. [JEE 2008, 3 + 3]

31. If the function $f(x) = x^3 + e^{\frac{x}{2}}$ and $g(x) = f^{-1}(x)$, then the value of $g'(1)$ is [JEE 2009, 4]
32. Let $f(\theta) = \sin \left(\tan^{-1} \left(\frac{\sin \theta}{\sqrt{\cos 2\theta}} \right) \right)$, where $-\frac{\pi}{4} < \theta < \frac{\pi}{4}$. then the value of $\frac{d}{d(\tan \theta)} (f(\theta))$ is [JEE 2011, 4]
33. The slope of the tangent to the curve $(y - x^5)^2 = x(1 + x^2)^2$ at the point $(1, 3)$ is [JEE(Advanced)-2014, 3]

SECTION-2

34. Let $f: \mathbb{R} \rightarrow \mathbb{R}$, $g: \mathbb{R} \rightarrow \mathbb{R}$ and $h: \mathbb{R} \rightarrow \mathbb{R}$ be differentiable functions such that $f(x) = x^3 + 3x + 2$, $g(f(x)) = x$ and $h(g(g(x))) = x$ for all $x \in \mathbb{R}$. Then-
 (A) $g'(2) = \frac{1}{15}$ (B) $h'(1) = 666$
 (C) $h(0) = 16$ (D) $h(g(3)) = 36$ [JEE(Advanced)-2016, 4(-2)]

35. For any positive integer n , define $f_n: (0, \infty) \rightarrow \mathbb{R}$ as

[JEE(Advanced)-2018, 4(0)]

$$f_n(x) = \sum_{j=1}^n \tan^{-1} \left(\frac{1}{1+(x+j)(x+j-1)} \right) \text{ for all } x \in (0, \infty)$$

(Here, the inverse trigonometric function $\tan^{-1} x$ assumes values in $\left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$)

(A) $\sum_{j=1}^5 \tan^2 (f_j(0)) = 55$

(B) $\sum_{j=1}^{10} (1 + f'_j(0)) \sec^2 (f_j(0)) = 10$

(C) For any fixed positive integer n , $\lim_{x \rightarrow \infty} \tan (f_n(x)) = \frac{1}{n}$

(D) For any fixed positive integer n , $\lim_{x \rightarrow \infty} \sec^2 (f_n(x)) = 1$

ANSWER KEY

DIFFERENTIATION OF A FUNCTION RESPECT TO ANOTHER FUNCTION

1. (A) 2. (D) 3. (A,C,D) 4. (A,B)

LOGARITHMIC FUNCTION/TRIGONOMETRIC SUBSTITUTIONS

5. (B,C)

INFINITE SERIES

6. (D)

DIFFERENTIATION OF PARAMETRIC EQUATIONS

7. (C) 8. (A)

MIXED PROBLEMS

9. (A,B)

JEE MAIN

10. (D) 11. (C) 12. (B) 13. (B) 14. (A) 15. (D) 16. (C)
17. (B) 18. (B) 19. (D) 20. (B) 21. (A) 22. (B) 23. (A)
24. (B) 25. (17) 26. (16) 27. (C)

JEE ADVANCED

SECTION-1

28. (C) 29. (D) 30. (a) A; (b) A 31. (2) 32. (1) 33. (8)

SECTION-2

34. (B,C) 35. (D)