



Q. find $[H^+]$, $[H_2PO_4^-]$, $[HPO_4^{2-}]$ & $[PO_4^{3-}]$ in 0.1M HCl & 0.1M H_3PO_4 (aq) soln.



Given $K_{a1} = 10^{-4}$, $K_{a2} = 10^{-7}$, $K_{a3} = 10^{-11}$.



$$10^{-4} = \frac{(0.1-x)x}{0.1-x} \quad x = 10^{-4}$$




$$10^{-7} = \frac{0.1 \times y}{x} \Rightarrow y = 10^{-10}$$




$$K_{a1} \times K_{a2} \times K_{a3} = 10^{-22} = \frac{[H^+]^3 [PO_4^{3-}]}{[H_3PO_4]}$$

$$10^{-22} = \frac{10^{-3} \times z}{0.1}$$

pH of a solution containing salt: \rightarrow





Salt



Salt






Acc. to Arrhenius





$$K_a = \frac{[\text{CH}_3\text{COO}^-][\text{H}^+]}{[\text{CH}_3\text{COOH}]}$$

Acc. to Bronsted




⇒ Na⁺, K⁺, Cl⁻, Br⁻, I⁻, NO₃⁻
do not react with H₂O





$$K_b = \frac{K_w}{K_a}$$

$$K_a K_b = K_w$$



$$\frac{K_w}{K_a} = K_b = \frac{\alpha \cdot \alpha}{c - \alpha} = \frac{\alpha^2}{c - \alpha} = \frac{c\alpha^2}{1 - \alpha}$$




Q. find pH of .

① 0.1M CH_3COONa (Given K_a of $\text{CH}_3\text{COOH} = 10^{-5}$)

$$K_b = \frac{10^{-14}}{10^{-5}} = 10^{-9}$$

② 0.01M KCN (Given K_a of $\text{HCN} = 10^{-10}$ M)

③ 0.1M NH_4Cl [Given K_a of $\text{NH}_4^+ = 10^{-9}$ M]



$$x = 10^{-5}$$

$$\text{pOH} = 5$$

$$\text{pH} = 9$$

1. Acetic acid (CH_3COOH) is partially dimerised to $(\text{CH}_3\text{COOH})_2$ in the vapour phase. At a total pressure of 0.200 atm, acetic acid is 92.0% dimerized at 298 K. The value of equilibrium constant (K_p) of dimerisation under these conditions is

(A) 57.5

(B) 9.7

(C) 97

(D) 194



2. Which of the following statement(s) is/are correct?

(a) The ratio of the radii of the first three Bohr orbits of hydrogen atom is 1: 8: 27.

T (b) The ratio of magnitude of total energy: kinetic energy: potential energy for electron in any orbit of hydrogen atom is 1: 1: 2.

(c) The frequency of a green light is 6×10^{14} Hz, then its wavelength is 500 nm.

(d) The ratio of de-Broglie wavelength of a H-atom, He-atom and CH_4 -molecule moving with equal kinetic energy is 4: 2: 1.

$$\lambda = \frac{h}{\sqrt{2mK.E}}$$

$$\frac{1}{1} : \frac{1}{\sqrt{4}} : \frac{1}{\sqrt{16}}$$

$$1 : \frac{1}{2} : \frac{1}{4}$$

3. Consider the following equilibrium



and the following graph :


Select the correct statements :

~~F~~ (A) Effect (I) represent the removal of reactant A

~~F~~ (B) Effect (II) represent the insertion of inert gas at constant volume

~~T~~ (C) Effect (III) represent the decrease in volume of container

~~T~~ (D) Effect (IV) represent the addition of moles of gas 'B'



4. A mixture containing substance 'X' & 'Y' is mixed with a substance 'Z'. If 'Z' reacts with both X & Y then select the correct option(s)

(A) Moles of X + moles of Y = Total moles of Z reacted "is always applicable"

(B) Equivalents of X + equivalents of Y = Total equivalents of Z reacted "is always applicable"

(C)
$$\frac{\text{Equivants. of X}}{n - \text{factor of Z in reaction with X}} + \frac{\text{Equivants of Y}}{n - \text{factor of Z in reaction with Y}} = \text{Total moles of}$$

Z reacted ; is always applicable

(D)
$$\frac{\text{moles of Z}}{\text{reacted in reaction with X}} + \frac{\text{moles of Z}}{\text{reacted in reaction with Y}} = \text{Total moles of Z reacted ; is}$$

always applicable

C. P


$$x + z$$

$$eq\ q\ x = \textcircled{eq\ q\ z}$$

$$y + z$$

$$eq\ q\ y = \textcircled{eq\ q\ z}$$

6. The value of n in the following processes: $\text{AO}_4^{n-} + 2e^- \rightarrow \text{HAO}_n^{2-}$ is




7. The minimum mass (in g) of CaCO_3 required to establish the equilibrium:

$\text{CaCO}_3(\text{s}) \rightleftharpoons \text{CaO}(\text{s}) + \text{CO}_2(\text{g})$, $K_c = 0.05\text{M}$ at a certain temperature in a 1.0 L container is

JEE ADVANCED PAPER-2

1. Isotherms of carbon dioxide at various temperatures are represented in Figure. The number of correct statement is/ are(s):
- (A) CO_2 exist only in gaseous state between the points 'a' and 'b' at temperature T_1 ?
- (B) At point 'c' CO_2 start liquefying when temperature is T_1 ?
- (C) No condensation takes place when the temperature is T_3 whatsoever the pressure.
- (D) Between point 'c' and 'd' at temperature T_1 both liquid and gaseous CO_2 exist at equilibrium?



Paragraph (Match list):

For the reaction: $\text{CaCO}_3(\text{s}) \rightleftharpoons \text{CaO}(\text{s}) + \text{CO}_2(\text{g})$; $K_p = 2.463 \text{ atm}$ at 900 K. The reaction is performed in a rigid vessel of 15.0 litre capacity maintained at 900 K, starting with the following amounts of $\text{CaCO}_3(\text{s})$ (given in Column I) Given $R = 0.0821 \text{ L}\cdot\text{atm}\cdot\text{K}^{-1}\cdot\text{mol}^{-1}$. Match the correct information given in Column II for the amount given in Column I.

$2.463 \times 15 = n \times 0.0821 \times 360 \times 3$
 $n = 1/2$

Column I		Column II	
(A)	100 g	(P)	50% dissociation of CaCO_3
(B)	50 g	(Q)	100% dissociation of CaCO_3
(C)	25 g	(R)	Addition of small amount of $\text{CO}_2(\text{g})$ may result shift of equilibrium in backward direction
		(S)	Addition of 10 g of $\text{CaCO}_3(\text{s})$ may result increase in moles of $\text{CO}_2(\text{g})$