

Collision b/w block C & A  
is perfectly inelastic.  
Find Time period and  
amplitude after Collision.

L.M.C

$$\frac{m}{4}u = m \cdot v$$




$$v = \left(\frac{u}{4}\right)$$

$$T = 2\pi \sqrt{\frac{\mu}{K}} = \left(2\pi \sqrt{\frac{m}{2K}}\right) \checkmark$$


$$\mu = \frac{m}{2} \quad v_{com} = \frac{mv}{2m} = \frac{v}{2}$$

$$A = \frac{v}{2\omega} = \frac{u}{8\omega} = \frac{u}{8 \times \frac{2\pi}{T}} = \frac{u}{16\pi} \times 2\pi \sqrt{\frac{m}{2K}}$$


$$A = \frac{u}{8} \sqrt{\frac{m}{2K}} \checkmark$$



$$\frac{u}{2} = A\omega$$



# Case of Vertical Spring



$$F_n = -[K(x_0 + x) - mg]$$

$$F_n = -[(Kx_0 - mg) + Kx]$$

$$F_n = -Kx$$

$$a = -\frac{K}{m}x$$

$$a = -\omega^2 x$$

$$\omega = \sqrt{\frac{K}{m}}$$

$$T = 2\pi\sqrt{\frac{m}{K}}$$

$$mgy = \frac{1}{2}Ky^2$$


$$y = \left(\frac{2mg}{K}\right)$$



# Extra force Method

Pully Massless, String & Spring Massless.



$$T = kx$$



$$F_y = T = -ky$$

$$a = -\frac{k}{m}y$$


$$T = \left(2\pi\sqrt{\frac{m}{k}}\right)$$





# Extra force Method

Pully Massless, String & Spring Massless.



$$T = 2Ky_1 \text{ (Pully Massless)}$$

$$T = -2K(2y)$$

$$T = -4Ky$$


$$a = -\frac{4K}{m}y$$

$$a = -\omega^2 y$$

$$T = 2\pi\sqrt{\frac{m}{4K}}$$

$$T = \pi\sqrt{\frac{m}{K}}$$

$$y = 2y_1$$



$$2T = Ky_1$$

$$T = \frac{K}{2}y_1 = \frac{Ky}{4}$$


$$F_y = -T = -\frac{K}{4}y$$

$$a = -\frac{K}{4m}y$$


$$a = -\omega^2 y$$

$$\omega = \sqrt{\frac{K}{4m}}$$

$$T = 2\pi\sqrt{\frac{4m}{K}} \quad \text{A}$$



All pulley and strings are massless.



$$y = 2(y_1 + y_2 + y_3 + y_4)$$

$$T \propto y$$

$$2T = K_1 y_1 = K_2 y_2 = K_3 y_3 = K_4 y_4 \Rightarrow \text{Series Combination.}$$

$$y_1 = \frac{2T}{K_1}, \quad y_2 = \frac{2T}{K_2}, \quad y_3 = \frac{2T}{K_3}, \quad y_4 = \frac{2T}{K_4}$$

$$y = 4T \left[ \frac{1}{K_1} + \frac{1}{K_2} + \frac{1}{K_3} + \frac{1}{K_4} \right]$$

$$T = \frac{y}{4 \left[ \frac{1}{K_1} + \frac{1}{K_2} + \frac{1}{K_3} + \frac{1}{K_4} \right]}$$


$$\omega^2 = \frac{1}{4m \left[ \frac{1}{K_1} + \frac{1}{K_2} + \frac{1}{K_3} + \frac{1}{K_4} \right]}$$

$$T = 2\pi \sqrt{\frac{4m \left( \frac{1}{K_1} + \frac{1}{K_2} + \frac{1}{K_3} + \frac{1}{K_4} \right)}{\frac{1}{K_{eq}}}} \Rightarrow T = 2\pi \sqrt{4m \left( \frac{1}{K_{eq}} \right)}$$


$$Q = \frac{y}{4m \left[ \frac{1}{K_1} + \frac{1}{K_2} + \frac{1}{K_3} + \frac{1}{K_4} \right]}$$



The whole system is placed on smooth horizontal table.  
Find time period of block if block is compressed along one of the spring and released.  
All the springs are at its natural length.




$x$  is Very Small So.  
angle b/w two Spring  
remain same after  
block is displaced from  
mean position.



$$F_y = -\left[Kx + 2 \cdot \left(\frac{Kx}{2} \cos 60^\circ\right)\right]$$

$$F_y = -\left[Kx + \frac{Kx}{2}\right] = -\left[\frac{3Kx}{2}\right]$$

$$a = -\frac{3K}{2m}x$$



$$a = -\omega^2 x$$

$$\omega = \sqrt{\frac{3K}{2m}}$$

$$T = 2\pi \sqrt{\frac{2m}{3K}} \quad \checkmark$$



Find the time period of cylinder.  
Cylinder rolls without slipping.

M-1

$$E_T = C, \quad \frac{dE_T}{dt} = 0.$$

→ Write total energy of the system at instant  $t=t$  at any intermediate position i.e b/w mean or extreme position.



$$E_T = \frac{1}{2} M \underline{v}^2 + \frac{1}{2} \left( \frac{MR^2}{2} \right) \underline{\omega}^2 + \frac{1}{2} K (2x)^2$$

$$v = R\omega \Rightarrow \omega = \frac{v}{R}$$

$$E_T = \frac{1}{2} M v^2 + \frac{MR^2}{4} \times \frac{v^2}{R^2} + 2Kx^2$$

$$E_T = \frac{Mv^2}{2} + \frac{Mv^2}{4} + 2Kx^2$$

$$E_T = \frac{3}{4} M v^2 + 2Kx^2$$



$$\begin{aligned} \vec{x}_{P/E} &= \vec{x}_{P/Com} + \vec{x}_{Com/E} & x &= R\theta \\ &= R\theta + x \\ &= \underline{2x} \end{aligned}$$



$$E_T = \frac{3}{4} M v^2 + 2Kx^2$$

$$\frac{dE_T}{dt} = \frac{3}{4} M \frac{d(v^2)}{dt} + 2K \frac{d(x^2)}{dt}$$

$$0 = \frac{3}{4} M \left( 2v \frac{dv}{dt} \right) + 2K (2x) \left( \frac{dx}{dt} \right)$$

$$0 = \frac{3Mv}{2} \left( \frac{dv}{dt} \right) + 4Kx \left( \frac{dx}{dt} \right)$$

$$\cancel{\frac{3Mv}{2} \left( \frac{dv}{dt} \right)} = -4Kx \left( \cancel{\frac{dx}{dt}} \right)$$

$$\Downarrow$$

$$\left( \frac{3M}{2} \right) a = -4Kx$$

$$a = -\frac{8K}{3M} x$$

$$a = -\frac{8K}{3M} x$$

$$a = -\omega^2 x$$

$$\omega = \sqrt{\frac{8K}{3M}}$$

$$T = 2\pi \sqrt{\frac{3M}{8K}} \checkmark$$

Find the time period of cylinder.  
Cylinder rolls without slipping. M-2

$$T_{res} = \text{---}$$

$$\alpha = \frac{T_{res}}{I}$$

$$\alpha = -\omega^2 \theta$$


$$\sin \theta = \frac{x}{2R}$$

$$x = 2R \sin \theta$$

$$x = 2R(\theta) \leftarrow$$

$$\omega^2 = \frac{8K}{3M}$$

$$T = 2\pi \sqrt{\frac{3M}{8K}}$$



$$\theta \rightarrow 0$$



$$\cos \theta \rightarrow 1$$

$$T_r = -(Kx) 2R \cos \theta$$

$$T_r = -K(2R\theta) \times 2R$$

$$T_r = -4KR^2\theta$$

$$\alpha = -\frac{4KR^2\theta}{\frac{3MR^2}{2}} = -\frac{8K}{3M}\theta$$



$$I_0 = \frac{MR^2}{2} + MR^2$$