


Link to View Video Solution: [Click Here](#)

1. \therefore for inclined projectile



$$T = \frac{2u_y}{a_y} = \frac{2u \sin \theta}{g \cos \beta}.$$

If it will hit the ground with 90°

i.e. $v_x = 0$.

$$\therefore v_x = u_x - a_x T \Rightarrow 0 = u_x - a_x T.$$


$$u_x = a_x T$$

$$u \cos \theta = g \sin \beta \cdot \frac{2u \sin \theta}{g \cos \beta}$$

$$\theta = \cot^{-1} [2 \tan \beta]$$


i.e. $k = 2$.

2.



$$h = x \tan \theta - \frac{gx^2}{2u^2 \cos^2 \theta} \text{ [standard eq.]}$$

$$h = x \tan \theta - \frac{gx^2 \sec^2 \theta}{2u^2}$$

Link to View Video Solution:  [Click Here](#)

$$h = x \tan \theta - \frac{gx^2}{2u^2} (1 + \tan^2 \theta)$$

$$2u^2h = 2u^2x \tan \theta - gx^2(1 + \tan^2 \theta).$$

$$gx^2 \tan^2 \theta - 2u^2x \tan \theta + 2u^2h + gx^2 = 0 \quad \text{---(1)}$$

for crossing the wall roots of above equation should be real.

$$\text{i.e. } 4u^4x^2 - 4gx^2(2u^2h + gx^2) \geq 0$$

$$u^4 - 2u^2gh - g^2x^2 \geq 0$$

$$u^4 - 2u^2gh + h^2g^2 - g^2x^2 - h^2g^2 \geq 0$$

$$(u^2 - gh)^2 \geq g^2(x^2 + h^2).$$

$$u \geq \sqrt{hg + g\sqrt{x^2 + h^2}}$$

$$u \geq \sqrt{g(h + \sqrt{h^2 + x^2})}$$

3. $h = l \sin 53 = 10 \sin 53$

$$h = 8 \text{ meter}$$

$$u_y = 10 \cos 53 = 6$$


$$\therefore h = u_y t + \frac{1}{2} g t^2$$


$$\therefore -8 = 6t + \frac{1}{2} \times 10 \times t^2$$

$$5t^2 - 6t - 8 = 0$$

$$(t - 2)(5t + 4) = 0$$

$$t = 2 \text{ seconds}$$




Link to View Video Solution:  [Click Here](#)

4. If particle P and Q are collide to each other after 4 seconds i.e. flight time of P is 4 seconds.

$$T = \frac{2u}{g \cos 60} = \frac{2u}{10 \times \frac{1}{2}}$$

$$4 = \frac{4u}{10}$$

$$u = 10 \text{ m/seconds}$$



5. $h = l \sin 53 = 10 \sin 53$

$$h = 8 \text{ meter}$$

$$u_y = 10 \cos 53 = 6$$


$$\therefore h = u_y t + \frac{1}{2} g t^2$$

$$\therefore -8 = 6t + \frac{1}{2} \times 10 \times t^2$$


$$5t^2 - 6t - 8 = 0$$

$$(t - 2)(5t + 4) = 0$$

$$t = 2 \text{ seconds}$$




- 6.



$$\text{Range} = AO = u_x \times t = u \cos 60 \times t = \frac{ut}{2}.$$

$$AB = \frac{AO}{\cos 30} = \frac{ut}{\sqrt{3}}$$

Link to View Video Solution:  [Click Here](#)

$$7. \quad \therefore T = \frac{2u_y}{a_y}$$

$$T = \frac{2 \times 40 \sin 53}{g \cos 37}$$

$$T = 8 \text{ seconds}$$

$$\therefore a_x = g \sin 37 = 6$$

$$a_y = g \cos 37^\circ = 8$$

$$u_y = 40 \sin 53 = 32$$

$$u_x = 40 \cos 53 = 24$$

$$R = u_x T + \frac{1}{2} a_x T^2$$

$$= 24 \times 8 + \frac{1}{2} \times 6 \times 8^2$$

$$= 192 + 3 \times 64$$


$$R = 384 \text{ meter}$$


particle will cover maximum height in half flight of time.

$$\text{i.e. } t = 4 \text{ seconds}$$

$$h = 32 \times 4 - \frac{1}{2} \times 8 \times 4^2$$

$$h = 64$$



Link to View Video Solution:  [Click Here](#)

8. $T = \frac{2u_y}{a_y} = \frac{2 \times 24}{8}$

$T = 6 \text{ seconds}$

$R = u_x T + \frac{1}{2} a_x T^2$

$= 32 \times 6 + \frac{1}{2} \times (-6) \times 6^2$

$R = 84$

$h = 24 \times 3 - \frac{1}{2} \times 8 \times (3)^2$

$h = 36 \text{ m}$

