


6.

$$(13\sin\theta - 5)(5\sin\theta - 3) = 0 \quad \text{and} \quad \tan\theta = \frac{12}{5}$$

$\sin\theta = \frac{5}{13}, \frac{3}{5}$

$$8\sin\theta = 4 + \cos\theta, \quad \sin\theta = ?$$

$$8\sin\theta - 4 = \cos\theta$$

$$64\sin^2\theta + 16 - 64\sin\theta = \cos^2\theta = 1 - \sin^2\theta$$

$$65\sin^2\theta - 64\sin\theta + 15 = 0$$

$$-39\sin\theta + 25\sin\theta$$

14:

$$2 - 2\sin\theta = \cos\theta$$

$$4(1-\sin\theta)^2 = 1-\sin^2\theta$$

$$(-\sin\theta) \left( 4 - 4\sin\theta - 1 - \sin\theta \right) = 0$$

$$(-\sin\theta)(3 - 5\sin\theta) = 0$$

$$\sin\theta = 1, \frac{3}{5}$$

$$y = x^2 = f(x)$$

$$R_f = [0, \infty)$$

$$D_f = R$$

$$f(x) = x^3$$

$$D_f = R$$

$$R_f = R$$

$$(-\sqrt[3]{16})$$

$$\begin{aligned} -2\sqrt[3]{2} &= -2 \cdot 2^{\frac{1}{3}} \\ y &= \pi \\ x &= \sqrt{\pi} \\ x &= -\sqrt{\pi} \end{aligned}$$

$$\frac{\cos x}{\sin x} = \cot x$$

$$y = \cot x = f(x)$$

$$D_f = R - \{n\pi\}, n \in \mathbb{I}$$

$$R_f = R$$

$$\tan x \rightarrow \begin{cases} \frac{\pi}{2} \rightarrow \pi \\ -\infty \rightarrow 0 \\ 0 \rightarrow -\infty \end{cases}$$

$$x \rightarrow 0 \text{ to } \frac{\pi}{2}$$

$$\cot x \rightarrow \begin{cases} \infty \rightarrow 0 \\ 0 \rightarrow \infty \end{cases}$$


$$\tan x \rightarrow \begin{cases} \infty \rightarrow 0 \\ 0 \rightarrow \infty \end{cases}$$


$$x \rightarrow 0.000 \dots$$


$$\tan x \rightarrow 0.000 \dots$$

$$\cot x \rightarrow \frac{1}{0.000 \dots} \rightarrow \infty$$

$$\cot \frac{\pi}{2} = \frac{\cos \frac{\pi}{2}}{\sin \frac{\pi}{2}} = \frac{0}{1} = 0$$







$$f(x) = \sec x$$

$$\cos x \neq 0$$

$$D_f = \mathbb{R} - \left\{ \left( 2n+1 \right) \frac{\pi}{2} \right\}, n \in \mathbb{I}$$

$$R_f = (-\infty, -1] \cup [1, \infty)$$

$$\sec^2 x \geq 1$$

$$\sec^2 x = 1 + \tan^2 x \quad | \quad 1 + \tan^2 x \leq 1 + \tan^2 x < \infty$$

$$\begin{aligned} t^2 &\geq 1 \\ (t-1)(t+1) &\geq 0 \end{aligned}$$

$\begin{array}{c} \text{---} \\ \text{+} \end{array}$        $\begin{array}{c} \text{+} \\ \text{---} \\ \text{+} \end{array}$        $\begin{array}{c} \text{+} \\ \text{+} \\ \text{+} \end{array}$

$$\begin{aligned} \tan x &\in \mathbb{R} \\ \tan^2 x &\in [0, \infty) \\ 1 + \tan^2 x &\in [1, \infty) \end{aligned}$$

$$f(x) = \operatorname{cosec} x$$

$$D_f = R - \{n\pi\}, n \in \mathbb{I}$$

$$R_f = (-\infty, -1] \cup [1, \infty)$$

$$\operatorname{cosec}^2 x = 1 + \cot^2 x \geq 1$$

$$(\operatorname{cosec} x - 1)(\operatorname{cosec} x + 1) \geq 0$$

$$\begin{array}{ccccccc} & + & - & + & & & \\ \hline -1 & & & & & & \end{array}$$

Note →

$$\sin(n\pi) = 0 \quad , n \in \mathbb{I}$$

$$\sin\left(2n\pi + \frac{\pi}{2}\right) = 1 \quad , n \in \mathbb{I}$$

$$\sin\left(2n\pi + \frac{3\pi}{2}\right) = -1 \quad , n \in \mathbb{I}$$

$$\cos\left((2n+1)\frac{\pi}{2}\right) = 0$$

$$\cos(2n\pi) = 1, n \in \mathbb{I}$$

$$\cos((2n+1)\pi) = -1$$

$$\tan(n\pi) = 0 \quad n \in \mathbb{I}$$


$$\tan\left((2n+1)\frac{\pi}{2}\right) = \text{not defined}$$

# Reduction formulae

$$\sin\left(\frac{\pi}{2} - \theta\right) = \cos\theta$$

$$\cos\left(\frac{\pi}{2} - \theta\right) = \sin\theta$$

$$\tan\left(\frac{\pi}{2} - \theta\right) = \cot\theta$$





$$\sin \theta = -\frac{y}{r}$$

$$\cos \theta = -\frac{x}{r}$$

$$\sin\left(\frac{\pi}{2} - \theta\right) = \frac{y}{r} = \cos \theta$$

$$\cos\left(\frac{\pi}{2} - \theta\right) = \frac{x}{r} = \sin \theta$$



$$\sin\left(\frac{\pi}{2} + \theta\right) = \cos\theta$$

$$\cos\left(\frac{\pi}{2} + \theta\right) = -\sin\theta$$

$$\tan\left(\frac{\pi}{2} + \theta\right) = -\cot\theta$$

$$\sin(\pi - \theta) = \sin\theta$$

$$\cos(\pi - \theta) = -\cos\theta$$

$$\tan(\pi - \theta) = -\tan\theta$$

$$\sin(\pi + \theta) = -\sin\theta$$

$$\cos(\pi + \theta) = -\cos\theta$$

$$\tan(\pi + \theta) = \tan\theta$$

$$\sin\left(\frac{3\pi}{2} - \theta\right) = -\cos\theta$$

$$\cos\left(\frac{3\pi}{2} - \theta\right) = -\sin\theta$$

$$\tan\left(\frac{3\pi}{2} - \theta\right) = \cot\theta$$

$$\sin\left(\frac{3\pi}{2} + \theta\right) = -\cos\theta$$

$$\cos\left(\frac{3\pi}{2} + \theta\right) = \sin\theta$$

$$\tan\left(\frac{3\pi}{2} + \theta\right) = -\cot\theta$$

$$\sin(-\theta) = \sin(2\pi - \theta) = -\sin\theta$$

$$\cos(-\theta) = \cos(2\pi - \theta) = \cos\theta$$

$$\tan(-\theta) = \tan(2\pi - \theta) = -\tan\theta$$

$$\sin(2\pi + \theta) = \sin\theta$$

$$\cos(2\pi + \theta) = \cos\theta$$

$$\tan(2\pi + \theta) = \tan\theta$$

