



$$4\sqrt{3} = \frac{1}{2} 2 \times 5 \frac{\sqrt{3}}{2} + \frac{1}{2} xy \frac{\sqrt{3}}{2}$$

$$xy = 7$$

$$2^2 + 5^2 - 2(2)(5)\cos 60^\circ = x^2 + y^2 - 2xy \cos 120^\circ$$

1. Simplify

$$\begin{vmatrix} 23 & 66 & 11 \\ 36 & 55 & 26 \\ 63 & 143 & 37 \end{vmatrix} = 11 \begin{vmatrix} 23 & 6 & 11 \\ 36 & 5 & 26 \\ 63 & 13 & 37 \end{vmatrix}$$

$\ominus =$

$$\begin{aligned} & \xrightarrow{C_1 \rightarrow C_1 - 4C_2} \downarrow 11 \begin{vmatrix} -1 & 6 & 1 \\ 16 & 5 & 16 \\ 11 & 13 & 11 \end{vmatrix} \\ & \xrightarrow{C_3 \rightarrow C_3 - 2C_2} \begin{vmatrix} -1 & 6 & 1 \\ 16 & 5 & 16 \\ 11 & 13 & 11 \end{vmatrix} \\ & = abc \left(1 + \frac{1}{a} + \frac{1}{b} + \frac{1}{c} \right) \end{aligned}$$

2. Show that

$$\begin{vmatrix} 1+a & 1 & 1 \\ 1 & 1+b & 1 \\ 1 & 1 & 1+c \end{vmatrix}$$

$$= abc \begin{vmatrix} 1+\frac{1}{a} & \frac{1}{b} & \frac{1}{c} \\ \frac{1}{a} & 1+\frac{1}{b} & \frac{1}{c} \\ \frac{1}{a} & \frac{1}{b} & 1+\frac{1}{c} \end{vmatrix} \xrightarrow{C_1 \rightarrow C_1 + C_2 + C_3} \left(1 + \frac{1}{a} + \frac{1}{b} + \frac{1}{c} \right) abc$$

$$\begin{aligned} & R_2 \rightarrow R_2 - R_1 \\ & R_3 \rightarrow R_3 - R_1 \end{aligned}$$

$$\begin{vmatrix} 1 & \frac{1}{b} & \frac{1}{c} \\ 0 & 1 & \frac{1}{c} \\ 0 & \frac{1}{b} & 1 \end{vmatrix}$$

$$abc \left(1 + \frac{1}{a} + \frac{1}{b} + \frac{1}{c} \right) = abc \left(1 + \frac{1}{a} + \frac{1}{b} + \frac{1}{c} \right) \begin{vmatrix} 1 & \frac{1}{b} & \frac{1}{c} \\ 0 & 1 & \frac{1}{c} \\ 0 & \frac{1}{b} & 1 \end{vmatrix}$$

3.
Simplify

$$\begin{vmatrix} 1 & \sin 3\theta & \sin^3 \theta \\ 2\cos \theta & \sin 6\theta & \sin^3 2\theta \\ \underbrace{4\cos^2 \theta - 1}_{3 - 4\sin^2 \theta} & \sin 9\theta & \sin^3 3\theta \end{vmatrix} = 0$$

↓

$$C_2 \rightarrow C_2 - 3\sin \theta C_1 + 4C_3$$

4. Let A28, 3B9 and 62C are 3 digit numbers divisible by integer K, then P.T. the determinant

$$\begin{vmatrix} A & 3 & 6 \\ 8 & 9 & C \\ 2 & B & 2 \end{vmatrix} \text{ is also divisible by } K.$$

$$\downarrow R_2 \rightarrow R_2 + 10R_3 + 100R_1$$

$$\begin{vmatrix} A & 3 & 6 \\ \underline{A28} & \underline{3B9} & \underline{62C} \\ 2 & B & 2 \end{vmatrix} = K \begin{vmatrix} A & 3 & 6 \\ 2 & B & 2 \\ 2 & B & 2 \end{vmatrix}$$

5.

Let

 $D_r =$

$$\begin{vmatrix} r & x & \frac{n(n+1)}{2} \\ 2r-1 & y & n^2 \\ 3r-2 & z & \frac{n(3n-1)}{2} \end{vmatrix}$$

, find $\sum_{r=1}^n D_r$

$$\sum_{r=1}^n D_r = D_1 + D_2 + \dots + D_n =$$

$$\begin{vmatrix} \sum_{r=1}^n r & x & \frac{n(n+1)}{2} \\ \sum_{r=1}^n (2r-1) & y & n^2 \\ \sum_{r=1}^n (3r-2) & z & \frac{n(3n-1)}{2} \end{vmatrix} = 0$$

6. If $\begin{vmatrix} p & b & c \\ a & q & c \\ a & b & r \end{vmatrix} = 0$, find the value of

$$\frac{p}{p-a} + \frac{q}{q-b} + \frac{r}{r-c} = 2$$

$$\begin{aligned} R_2 &\rightarrow R_2 - R_1 \\ R_3 &\rightarrow R_3 - R_1 \end{aligned}$$

$$\begin{vmatrix} p & b & c \\ a-p & q-b & 0 \\ a-p & 0 & r-c \end{vmatrix} = p(q-b)(r-c) + b(p-a)(r-c) + c(p-a)(q-b) = 0$$

$$0 \Rightarrow \frac{p}{p-a} + \frac{b-q+q}{q-b} + \frac{c-r+r}{r-c} \Rightarrow \frac{p}{p-a} + \frac{b}{q-b} + \frac{c}{r-c} = 0$$

7. Simplify

$$\begin{vmatrix} (b+c)^2 & a^2 & bc \\ (c+a)^2 & b^2 & ca \\ (a+b)^2 & c^2 & ab \end{vmatrix} \quad \downarrow C_1 \rightarrow C_1 + C_2 - 2C_3$$

$$\begin{pmatrix} 2 & 2 & 2 \\ a & b & c \end{pmatrix} \begin{vmatrix} 1 & a & a^3 \\ 1 & b & b^3 \\ 1 & c & c^3 \end{vmatrix}$$

//

$$= (a^2 + b^2 + c^2)$$

$$\begin{vmatrix} 1 & a^2 & bc \\ 1 & b^2 & ca \\ 1 & c^2 & ab \end{vmatrix}$$

$$\begin{pmatrix} 2 & 2 & 2 \\ a & b & c \end{pmatrix} \begin{vmatrix} a & a^3 & 1 \\ b & b^3 & 1 \\ c & c^3 & 1 \end{vmatrix} = \frac{(a^2 + b^2 + c^2)}{abc} \begin{vmatrix} a & a^3 & abc \\ b & b^3 & abc \\ c & c^3 & abc \end{vmatrix}$$

$$\begin{aligned} & (a^2 + b^2 + c^2)(a-b)(b-c) \\ & (c-a)(a+b+c) \end{aligned}$$

8. P.T. $\begin{vmatrix} a & b-c & b+c \\ a+c & b & c-a \\ a-b & a+b & c \end{vmatrix} = (a^2+b^2+c^2)(a+b+c)$

$\frac{\sum a(\sum a^2)}{a^2bc}$ $\begin{vmatrix} 1 & b^2 & c^2 \\ 1 & b^2 & c^2-ac \\ 1 & b^2 & c^2 \end{vmatrix}$

$\frac{(a+b+c)}{a^2bc}$ $\begin{vmatrix} a^2 & b^2 & c^2 \\ a^2+ac & b^2 & c^2-ac \\ a^2-ab & b^2 & c^2 \end{vmatrix}$

$\frac{(a+b+c)}{a^2bc}$ $\begin{vmatrix} a^2 & b^2 & c^2 \\ a^2+ac & b^2 & c^2-ac \\ a^2-ab & b^2 & c^2 \end{vmatrix}$

$\begin{vmatrix} a^2 & ab-ac & ab+ac \\ bat+bc & b^2 & bc-ba \\ ca-cb & catcb & c^2 \end{vmatrix}$

$R_1 \rightarrow R_1 + R_2 + R_3$

$\frac{(a+b+c)}{a}$ $\begin{vmatrix} a & b & c \\ a+c & b & c-a \\ a-b & a+b & c \end{vmatrix}$

$\frac{(a+b+c)}{abc}$ $\begin{vmatrix} a & b & c \\ ab+bc & b^2 & bc-ba \\ ca-cb & catcb & c^2 \end{vmatrix}$

$\frac{\sum a \sum a^2}{a^2}$ $\begin{vmatrix} 1 & b & c \\ 1 & b & c-a \\ 1 & a+b & c \end{vmatrix}$

$\frac{\sum a \sum a^2}{a^2}$ $\begin{vmatrix} 1 & b & c \\ 0 & 0 & -a \\ 0 & a & 0 \end{vmatrix}$

$$\begin{vmatrix} a & b-c & b+c \\ a+c & b & c-a \\ a-b & a+b & c \end{vmatrix} \xrightarrow{C_1 \rightarrow aC_1 + bC_2 + cC_3}$$

$$\boxed{11-25}$$

Ex-11

$$\frac{1}{a} \begin{vmatrix} a^2+b^2+c^2 & b-c & b+c \\ a^2+b^2+c^2 & b & c-a \\ a^2+b^2+c^2 & a+b & c \end{vmatrix} = \frac{(a^2+b^2+c^2)}{a}$$

$$\frac{(a^2+b^2+c^2)}{a} \begin{vmatrix} 1 & b-c & b+c \\ 0 & c & -a-b \\ 0 & a+c & -b \end{vmatrix}$$

$$\begin{vmatrix} 1 & b-c & b+c \\ 1 & b & c-a \\ 1 & a+b & c \end{vmatrix}$$

$$\begin{aligned} R_2 &\rightarrow R_2 - R_1 \\ R_3 &\rightarrow R_3 - R_1 \end{aligned}$$