

Indefinite Integration

Antiderivative / Primitive / Integral of a function

reverse process of differentiation.

$$\int \frac{d}{dx} x = x + C$$

$$\frac{d}{dx} f(x) = g(x)$$

$$\Rightarrow d(f(x)) = g(x) dx$$

$$d(\sin x) = \cos x dx$$

$$d(x^4) = 4x^3 dx$$

$$\frac{d}{dx} (f(x) + 2) = g(x)$$

Integrand.

$$\int g(x) dx = f(x) + C$$

arbitrary constant

Indefinite

integration constant.

$$\int x^n dx = \frac{x^{n+1}}{n+1} + C, \quad n \neq -1$$

$$\int (ax+b)^n dx = \frac{(ax+b)^{n+1}}{a(n+1)} + C, \quad n \neq -1$$

$$\int \frac{dx}{x} = \ln|x| + C$$

$$\int \frac{dx}{ax+b} = \frac{1}{a} \ln|ax+b| + C$$

$$\frac{d}{dx} \frac{(ax+b)^{n+1}}{a(n+1)} = (ax+b)^n$$

$$\int e^x dx = e^x + C$$

;

$$\int e^{px+q} dx = \frac{1}{p} e^{px+q} + C$$

$$\int a^x dx = \frac{a^x}{\ln a} + C$$

;

$$\int a^{px+q} dx = \frac{a^{px+q}}{p \ln a} + C$$

$$\int \sin x dx = -\cos x + C$$

;

$$\int \sin(ax+b) dx = -\frac{1}{a} \cos(ax+b) + C$$

$$\int \cos x dx = \sin x + C$$

;

$$\int \cos(ax+b) dx = \frac{1}{a} \sin(ax+b) + C$$

$$\int \sec^2 x dx = \tan x + C \quad ; \quad \int \sec^2(ax+b) dx = \frac{1}{a} \tan(ax+b) + C$$

$$\int \csc^2 x dx = -\cot x + C \quad ; \quad \int \csc^2(ax+b) dx = -\frac{1}{a} \cot(ax+b) + C$$

$$\int \sec x \tan x dx = \sec x + C \quad ; \quad \int \sec(ax+b) \tan(ax+b) dx = \frac{1}{a} \sec(ax+b) + C$$

$$\int \csc x \cot x dx = -\csc x + C \quad ; \quad \int \csc(ax+b) \cot(ax+b) dx = -\frac{1}{a} \csc(ax+b) + C.$$

$$\int \frac{dx}{\sqrt{1-x^2}} = \sin^{-1} x + C$$

$$; \int \frac{dx}{\sqrt{a^2-x^2}} = \sin^{-1}\left(\frac{x}{a}\right) + C$$

$$\int \frac{dx}{1+x^2} = \tan^{-1} x + C$$

$$; \int \frac{dx}{a^2+x^2} = \frac{1}{a} \tan^{-1}\left(\frac{x}{a}\right) + C$$

$$\int \frac{dx}{|x| \sqrt{x^2-1}} = \sec^{-1} x + C$$

$$; \int \frac{dx}{|x| \sqrt{x^2-a^2}} = \frac{1}{a} \sec^{-1}\left(\frac{x}{a}\right) + C$$

$$\underline{1.} \quad \int 2^{\ln x} dx = \int x^{\ln 2} dx = \frac{x^{\ln 2 + 1}}{(\ln 2 + 1)} + C.$$

$$\int \frac{(x+5)+(x-2)}{(x+5)(x-2)} dx = \int \left(\frac{1}{x-2} + \frac{1}{x+5} \right) dx = \ln|x-2| + \ln|x+5| + C.$$

$$\underline{2.} \quad \int \frac{\sqrt{x^4 + x^{-4} + 2}}{x^3} dx = \int \frac{x^2 + \frac{1}{x^2}}{x^3} dx = \int \left(\frac{1}{x} + \frac{1}{x^5} \right) dx$$

$$\underline{3.} \quad \int \frac{x dx}{(x^2 + 2x + 1)} = \int \frac{(x+1-1)dx}{(x+1)^2} = \ln|x| - \frac{1}{4x^4} + C.$$

$$= \int \left(\frac{1}{x+1} - \frac{1}{(x+1)^2} \right) dx$$

$$\underline{4.} \quad \int \frac{(2x+3)dx}{(x^2+3x-10)} = \ln|x+1| + \frac{1}{(x+1)} + C.$$

$$\underline{5.} \quad \int \frac{x \, dx}{(x^2 - 5x + 6)} = \int \frac{(3(x-2) - 2(x-3))}{(x-2)(x-3)} \, dx = \int \left(\frac{3}{x-3} - \frac{2}{x-2} \right) dx$$

$$\underline{6.} \quad \int (2^x + 3^x)^2 \, dx = \int (4^x + 9^x + 2 \cdot 6^x) \, dx = 3 \ln|x-3| - 2 \ln|x-2| + C.$$

$$= \frac{4^x}{\ln 4} + \frac{9^x}{\ln 9} + \frac{2 \cdot 6^x}{\ln 6} + C$$

$$\underline{7.} \quad \int \frac{(e^{3x} + e^{5x})}{(e^x + e^{-x})} \, dx = \int \frac{e^x e^{3x} (1 + e^{2x})}{(1 + e^{2x})} \, dx = \int e^{4x} \, dx = \frac{e^{4x}}{4} + C.$$

$$\underline{8.} \quad \int \cos 2x \cos 3x dx = \int \frac{1}{2} (\cos 5x + \cos x) dx = \frac{1}{2} \left(\frac{\sin 5x}{5} + \sin x \right) + C.$$

$$\underline{9.} \quad \int \sin^4 x dx = \int \left(\frac{1 - \cos 2x}{2} \right)^2 dx = \frac{1}{4} \int (1 + \cos^2 2x - 2\cos 2x) dx$$

$$= \frac{1}{4} \int \left(1 + \frac{1 + \cos 4x}{2} - 2\cos 2x \right) dx = \frac{1}{4} \int \left(\frac{3}{2} + \frac{\cos 4x}{2} - 2\cos 2x \right) dx$$

$$= \frac{1}{4} \left(\frac{3}{2}x + \frac{\sin 4x}{8} - \sin 2x \right) + C.$$

$$\underline{10.} \quad \int \frac{(\cos 5x + \cos 4x) dx}{(1 - 2\cos 3x)}$$

$$= \int -2\cos \frac{3x}{2} \cos \frac{x}{2} dx$$

$$= - \int (\cos x + \cos 2x) dx$$

$$= \int \frac{2\cos \frac{3x}{2} \cos \frac{x}{2} dx}{1 - 2\left(2\cos^2 \frac{3x}{2} - 1\right)}$$

$$= \int \frac{2\cos \frac{3x}{2} \cos \frac{x}{2} dx}{3 - 4\cos^2 \frac{3x}{2}}$$

$$\underline{11.} \quad \int \sin(x^\circ) dx = \int \sin\left(\frac{\pi}{180}x\right) dx = -\frac{180}{\pi} \cos \frac{\pi x}{180} + C$$

$$= -\frac{180}{\pi} \cos x^\circ + C$$

$$\cot^2 x \cos^2 x = \cot^2 x - \cos^2 x$$

$$\tan^2 x \sin^2 x = \tan^2 x - \sin^2 x$$

$$\underline{12.} \quad \int \left(\frac{1 - \cos x}{1 + \cos x} \right) dx = \int \tan^2 \frac{x}{2} dx = \int \left(\sec^2 \frac{x}{2} - 1 \right) dx = 2 \tan \frac{x}{2} - x + C$$

$$\underline{13.} \quad \int \sec^2 x \operatorname{cosec}^2 x dx = \int \frac{\sin^2 x + \cos^2 x}{\sin^2 x \cos^2 x} dx = \int (\sec^2 x + \operatorname{cosec}^2 x) dx$$

$$= \tan x - \cot x + C$$

$$\underline{14.} \quad \int \cot^2 x \cos^2 x dx$$

$$= \int (\cot^2 x - \cos^2 x) dx$$

$$= \int \left(\operatorname{cosec}^2 x - 1 - \frac{1 + \cos 2x}{2} \right) dx = -\cot x - \frac{3}{2}x - \frac{\sin 2x}{4} + C$$

$$\underline{15.} \quad \int \frac{(\operatorname{cosec} x + \tan^2 x + \sin^2 x) dx}{\sin x} = \int (\operatorname{cosec}^2 x + \tan x \sec x + \sin x) dx$$

$$= -\cot x + \sec x - \cos x + C.$$

$$\underline{16.} \quad \int \frac{dx}{(1 - \sin 3x)} = \int \frac{\tan x \sec x (1 + \sin 3x)}{\cos^2 3x} dx = \int (\sec^2 3x + \tan 3x \sec 3x) dx$$

$$= \frac{\tan 3x}{3} + \frac{\sec 3x}{3} + C.$$

$$\underline{17.} \quad \int \frac{(x^2 + \cos^2 x) \operatorname{cosec}^2 x dx}{(x^2 + 1)}$$

$$\int \frac{(x^2 + 1 - \sin^2 x) \operatorname{cosec}^2 x dx}{x^2 + 1} = \int \left(\operatorname{cosec}^2 x - \frac{1}{x^2 + 1} \right) dx$$

$$= -\cot x - \tan^{-1} x + C.$$

$$\underline{1.} \quad \int \frac{x^4 dx}{1+x^2}$$

$$\underline{2.} \quad \int \frac{dx}{\sqrt{9-4x^2}}$$

$$\underline{3.} \quad \int \frac{dx}{(x^2-4x+4)(x^2-4x+5)}$$

$$\underline{4.} \quad \int \frac{dx}{(2x-7)\sqrt{(x-3)(x-4)}}$$