





A Vertical Vegetation Structure Model of Europe

Hui Zhang^{1*}, Nico Lang^{1*}, Stefan Oehmcke², Mikolaj Tymon Mazurczyk¹,
Martin Stefan Brandt¹, Rasmus Fensholt¹, Ankit Kariyaa¹, Christian Igel^{1*}

¹University of Copenhagen ²University of Rostock


* {huzh, nila, igel}@di.ku.dk



Highlights

- ✓ **Beyond canopy top height:** We reveal the full relative height profiles of forests, which may provide new insights about biodiversity, carbon sequestration, human activity, etc.
- ✓ **Multi-year coverage:** Unlike existing 2020-only maps, our 3D vegetation structure spans 2019–2022, enabling time series analysis.
- ✓ **Definition-agnostic:** Full-profile modeling allows direct comparison with canopy height maps using RH95 (Meta⁴, UMD³), RH98 (ETH¹), RH100 (UM²).

Data & Method






- ✓ Sampled over **300 million** locations worldwide from 2019 to 2022.
- ✓ Included **slope data** (from Copernicus DEM) to reduce misinterpretation of GEDI signals in steep or low-vegetation areas (e.g., grasslands, moss/lichen).
- ✓ Used **ESA World Cover** to fill GEDI gaps and improve **generalization** in areas without lidar reference.

S2: Optical Images
Dense 2D

Vertical Structure Model (VSM)
Dense 3D

GEDI: Structure Profiles
Sparse 3D



References

- [1] Lang, Jetz, Schindler, Wegner. A high-resolution canopy height model of the Earth. *Nature Ecology & Evolution*, 2023
- [2] Pauls, Zimmer, Kelly, Schwartz, Saatchi, Ciais, Pokutta, Brandt, Gieseke. Estimating canopy height at scale. *ICMI*, 2024
- [3] Potapov, Li, Hernandez-Serna, Tyukavina, Hansen, Kommareddy, Pickens, Turubanova, Tang, Silva, Armston, Dubayah, Blair, Hofton. Mapping global forest canopy height through integration of GEDI and Landsat data. *Remote Sensing of Environment*, 2021
- [4] Tolan, Yang, Nosarzewski, Couairon, Vo, Brandt, Spore, Majumdar, Haziza, Vamaraju, Moutakanni, Bojanowski, Johns, White, Tiecke, Couprie. Very high resolution canopy height maps from RGB imagery using self-supervised vision transformer and convolutional decoder trained on aerial lidar. *Remote Sensing of Environment*, 2024

Table 1. Comparison of canopy top height maps. *Max sampled within a circular footprint corresponding to GEDI (25m diameter)

VSM advantages

- Captures vertical structure details missed by standard canopy height models
- Successfully estimates lower height metrics (understory structure) using GEDI & Sentinel-2
- Provides critical information for diversifying forests under changing climate zones

Limitations & future work

- Current optical models cannot "see through" closed canopies.
- Integrating SAR data (e.g., ESA BIOMASS mission) to resolve canopy penetration issues.

Acknowledgments

This work is supported by the Danish National Research Foundation (DNRF) through TreeSense, the Center for Remote Sensing and Deep Learning of Global Tree Resources (DNRF192), the Novo Nordisk Foundation through the Global Wetland Center (grant number NNF23OC0081089), and the Pioneer Centre for AI, DNRF grant number P1.