
A Vertical Vegetation Structure Model of Europe

Hui Zhang^{1*} Nico Lang^{1*} Stefan Oehmcke² Mikolaj Tymon Mazurczyk¹
Martin Stefan Brandt¹ Rasmus Fensholt¹ Ankit Karirya¹ Christian Igel^{1*}
¹University of Copenhagen ²University of Rostock
*{huzh, nila, igel}@di.ku.dk

Abstract

Vertical vegetation structure data has the potential to reveal nuanced ecosystem response to climate change and disturbances such as from wildfires, droughts, deforestation, and forest degradation. However, existing global-scale studies mainly focus on canopy top height or simplified single descriptors of vertical structure at low spatial resolution. Here, we address this gap by integrating full-waveform lidar observations from the Global Ecosystem Dynamics Investigation (GEDI) and Sentinel-2 optical images into a wall-to-wall vertical structure model (VSM). The model provides a dense map of estimated relative height profiles at 10 meter resolution for Europe. We see great potential in the presented vertical structure model for advancing science and environmental resource management.

1 Introduction

Intact vegetation ecosystems provide a central basis for human and animal life [35] by regulating the climate at local and global scales, regulating the hydrological cycle, and conserving biodiversity. They can reduce the destructive impact of severe weather events and climate change [22], for example, by lowering the risk of soil erosion and land-slides, or by acting as a buffer against hurricanes, storm surges, or avalanches. Even so, pristine ecosystems are globally under immense pressure due to human exploitation of natural resources [37] often associated with the production of commodities such as palm oil, cocoa, soybeans, and cattle [14, 17].

With $\approx 85\%$ of Europe's forested areas being actively managed, often as monocultures with mainly a single or only a few tree species, there are great challenges, but also opportunities ahead to strengthen their climate resilience [11]. These monocultures are vulnerable to extended periods of drought and heat waves, which can lead to vast losses, as for example seen in Germany, where large areas were emergency logged to prevent the further spread of bark beetle pest [32]. In addition, remaining natural ecosystems are fragmented and located in proximity to land with high human activity, such as agriculture or urban areas. This proximity can cause "edge effects" with substantial impacts on ecosystem services even across land use boundaries [34].

Even formally protected areas are impacted by human activities [19]. The urgency of addressing the global challenges of adapting to a changing climate and avoiding biodiversity collapse [15] have led to several international environmental initiatives to preserve the remaining intact ecosystems and to reverse degraded habitats to a natural state. Examples include the UN Decade on Ecosystem Restoration [25] and the Kunming-Montreal Global Biodiversity Framework [1]. A specific target of the latter known as the "30 by 30" commitment aims to protect 30% of the world's land, ocean, and freshwater areas by the year 2030 [1]. Furthermore, in May 2023 the European Union (EU) enacted the EU Deforestation Regulation [13] to limit deforestation caused by the major agricultural commodities sold in the EU.

A crucial step towards implementing such global conservation targets on land is to quantify the conservational value of vegetation ecosystems to guide resources towards protecting the most valuable




Figure 1: Illustration of our approach. Sentinel-2 satellite images are used as a *dense 2D* input to a convolutional neural network (CNN) trained with *sparse 3D* supervision of GEDI vertical profiles. The output is a *dense 3D* Vertical Structure Model (VSM) with a 10m ground sampling distance.

ecosystems [29, 31, 16, 26]. Quantifying this conservational value is challenging as we need to account for a multitude of criteria including carbon stocks [5], biodiversity [36], and the naturalness of ecosystems [3]. Conservation and afforestation of degraded forest ecosystems is discussed as mechanisms to capture carbon from the atmosphere, but high uncertainties of estimated carbon stock potentials remain a challenge [24]. Furthermore, monitoring ecosystem functions is crucial to understand how ecosystems respond to anthropogenic drivers and a changing climate [23].

Vertical vegetation structure is an essential variable [12] that allows to quantify aboveground carbon stocks, biodiversity, and the naturalness of ecosystems [7, 4, 21, 26]. In addition, vegetation structure properties dominate the explanation of the variation of ecosystem functioning [23]. Vertical vegetation structure has been studied to quantify the effectiveness of protected areas [19, 8, 2, 20], but to better understand human drivers influencing three-dimensional forest structure and to facilitate the effective management of protected areas, improved vegetation structure data is needed that is globally consistent, timely, and provide high spatial resolution [19].

Here, we present a globally applicable approach to model the full vertical structure profile of vegetation by combining data from two different space missions: Sentinel-2 satellite images and GEDI space borne lidar profiles. Our approach advances state-of-the-art that are limited to single statistics as proxies for the vertical distribution of biomass (like canopy top height [30, 18, 27] or entropy [4]). We extend the established concept of *canopy height models (CHM)*, i.e., the 2.5D raster data of vegetation height above ground, to the concept of *vertical structure models (VSM)* to represent the full 3D structure of terrestrial ecosystems, capturing both horizontal *and* vertical variation (see illustration in Fig. 1).

We use our developed model to compute annual VSMs for Europe (i.e. countries belonging to the EU, EFTA, and UK). These VSMs are three-dimensional data cubes covering the entire landmass, representing the vertical structure distribution at 10 meter spatial resolution. This data advances the availability of global structure data in spatial coverage, as well as spatial and temporal resolution. While our VSM also provides canopy top heights advancing or matching the accuracy of state-of-the-art, our approach provides full vertical profiles and thus structure variations in the understory that are more sensitive to small canopy gaps.

2 Combining Sentinel-2 optical images and GEDI spaceborne lidar

There is no single satellite sensor that can provide dense global vertical vegetation structure data. While dedicated laser scanner instruments like the Global Ecosystem Dynamics Investigation (GEDI) have advanced the availability of sparse global-scale vertical structure measurements [6], the derived map products that aggregate these sparse measurements fall short in terms of spatial resolution (1 km), global coverage beyond 51.6° N & S, and are not designed for monitoring over time [4]. On the other hand, optical satellite missions like Sentinel-2 or Landsat provide global coverage at spatial resolutions up to 10 m with revisit times of five days, but cannot provide a direct measurement of vertical structure. Modeling vegetation parameters using both types of sensors promises to overcome

the limitations of each sensor alone and the progress of artificial intelligence and modern computer vision methods has led to advanced canopy height products [18, 27]. Nevertheless, the state-of-the-art remains limited by capturing only few statistics of the structure.

Our methodology combines dense 2D satellite imagery from Sentinel-2 and sparse 3D GEDI LiDAR footprints to produce high-resolution dense 3D vertical structure models for Europe (see Fig. 1). We train a convolutional neural network (CNN) to predict the GEDI vertical profile for each Sentinel-2 pixel using sparse supervision and a quantile regression loss.

GEDI lidar data provide sparse height measurements for different canopy layers. We use Relative Height (RH) from the L2A product as reference data. For each GEDI shot, we obtain 101 RH metrics (RH0–RH100) describing the vertical structure of the 3D structure within a 25m footprint. Each RH value represents the height above the ground at which a specific percentage of the total laser energy is returned. In this study, we only retained footprints with a sensitivity value greater than 0.95, following [4], to ensure reliable ground returns in dense canopy cover. Additional quality flags provided in the L2A product were used to remove invalid or low-quality data. To capture vegetation structure under leaf-on conditions, we further restricted the dataset to footprints acquired during the growing season, derived from NPP VIIRS Global Land Surface Phenology product [39].

Sentinel-2. Previous studies on canopy top height mapping [30, 18, 28, 33] have demonstrated that predictive representations can be learned from optical satellite imagery to estimate vegetation height. Building on previous findings in [18], we used 12 optical bands from Sentinel-2 L2A data as input for vegetation structure modeling. To ensure temporal consistency with the GEDI observations, only Sentinel-2 images acquired during the growing season were used. A strict patch-level cloud coverage filter was applied to get a high quality dataset.

Auxiliary data. ESA WorldCover 2021 land cover map [38] was used for multitask learning. The ESA land cover information provides surface type context that helps the model distinguish vegetated from non-vegetated surfaces. Including this land cover mapping task enables the model to learn broader land surface characteristics, thereby reducing unrealistic vertical vegetation profile predictions in regions beyond the GEDI range. Besides, the land cover prediction can enhance the explainability of the model. We also use the slope information derived from the Copernicus DEM [9] to only optimize the RH regression on terrain with $<15^\circ$ slope (threshold following [27]). Steep terrain can lead to a distorted GEDI waveform and yield unreliable vertical profile estimates. However, we optimize the land cover as an auxiliary task in all terrain conditions.

3 Experiments and results

Experimental setup. We sampled a total of 300 million GEDI shots uniformly across the global landmass. After quality filtering and pairing with cloud-free Sentinel-2 image patches, we ended up with a total of 179 million data points. At each location, we sampled a 15×15 pixel patch from the closest, cloud-free Sentinel-2 image within the growing season of that year. To train and evaluate our model, we followed prior work [18] and splitted the data by Sentinel-2 tiles that correspond to approximately $100\text{km} \times 100\text{km}$. Therefore, we held out 12.6 % of randomly selected tiles for testing (i.e. 1392 tiles, with 52 in Europe). We trained our model for 2,234,400 iterations with batch size 4096 using the Adam optimizer.

Deployment and VSM computation. To compute an annual VSM for the year 2020, we sampled up to 20 images with the lowest cloud cover within the growing season for every Sentinel-2 tile. We used a "late" median aggregation strategy following [18]. Hence, each image was processed to predict a date-specific VSM, that was filtered to remove cloudy pixels and pixels with predicted snow and urban landcover before aggregating the individual predictions to a median composite (see Fig. 2).

Evaluation. The RH10, RH25, RH50 and RH98 predictions for Europe are shown in Fig. 2 (top). There are no border artifacts within individual Sentinel-2 tiles, however, deploying the model across adjacent Sentinel-2 tiles remains challenging, as predictions from different dates can lead to tiling artifacts.

We evaluated all RH metrics on the 52 hold-out test tiles in Europe and visualize the residual distribution for each RH (Fig. 2 bottom). While all RH have a low bias, we observe that the variation of residuals increases towards both RH0 and RH100, having the lowest variation around RH35. The




Figure 2: Vertical structure model of Europe. Top: Maps of key metrics RH10, RH25, RH50, RH98. Note, while lower RH metrics can be negative mainly for bare ground landcover, we only visualize positive values. Bottom: Residual error per RH metric on hold-out GEDI data. Boxplots show the median, quartiles and the 5th and 95th percentile for RH0–RH100.




Figure 3: Comparison of RH25 and RH98 in capturing small structures like canopy gaps. Left: Input Sentinel-2 image. Middle: Predicted RH25. Right: Predicted RH98

bias for RH95, RH98, and RH100 compares well with previous studies [30, 18, 27]. An example is provided showing the qualitative differences between predictions for different RH values (Fig. 3).

4 Discussion and conclusions

Our VHM provides additional information not captured by an CHM. The VSM makes full use of the information contained in the GEDI profile measurements. While our estimates of lower RH metrics capture the understory structure, these are dominantly sensitive to small canopy gaps that are closed by the canopy top height proxies (i.e., $>\text{RH95}$) because of GEDI's 25m footprint (Fig. 3). Our model relies on texture structure in Sentinel-2 optical images and cannot "see through" closed canopies. Thus, there is great potential to complement our VSM with SAR data, such as, for example, the ESA BIOMASS mission [10].

The monitoring of resources become more and more important under the changing climate. Beyond forests' importance for biodiversity conservation and carbon storage, Europe is especially relying on several direct ecosystem services in all climate zones. Intact forest ecosystems regulate the local climate which can dampen extended heatwaves in the South and in alpine regions, forests act as protection against avalanches or landslides. However, both intact forest ecosystems and managed forests are under stress. With the changing climate, insects like the bark beetle spread quickly, leading to loss of a large number of trees. European forest management needs to cope with these stress factors by finding new strategies to diversify our forests and make them resilient under the changing climate. We envision that our VSM will be beneficial for monitoring and proactive management, also to reduce the risk of wildfires by identifying fuel material in the understory.

Acknowledgments and Disclosure of Funding

This work is supported by the Danish National Research Foundation (DNRF) through TreeSense, the Center for Remote Sensing and Deep Learning of Global Tree Resources (DNRF192), the Novo Nordisk Foundation through the Global Wetland Center (grant number NNF23OC0081089), and the Pioneer Centre for AI, DNRF grant number P1.

References

- [1] Kunming-montreal global biodiversity framework. In *Convention Biol. Divers. Kunming-Montreal Glob. Biodivers. Framew.(cbd. int)*. Convention on Biological Diversity, 2022.
- [2] Guido Ceccherini, Marco Girardello, Pieter SA Beck, Mirco Migliavacca, Gregory Duveiller, Grégoire Dubois, Valerio Avitabile, Luca Battistella, José I Barredo, and Alessandro Cescatti. Spaceborne lidar reveals the effectiveness of european protected areas in conserving forest height and vertical structure. *Communications Earth & Environment*, 4(1):97, 2023.
- [3] Alessandro Chiarucci and Gianluca Piovesan. Need for a global map of forest naturalness for a sustainable future. *Conservation Biology*, 34(2):368–372, 2020.
- [4] Tiago De Conto, Ralph Dubayah, and John Armston. Characterizing the structural complexity of the Earth’s forests with spaceborne lidar, May 2024. URL <https://www.researchsquare.com/article/rs-4445870/v1>.
- [5] Moreno Di Marco, Stuart HM Butchart, Piero Visconti, Graeme M Buchanan, Gentile F Ficetola, and Carlo Rondinini. Synergies and trade-offs in achieving global biodiversity targets. *Conservation Biology*, 30(1):189–195, 2016.
- [6] Ralph Dubayah, James Bryan Blair, Scott Goetz, Lola Fatoyinbo, Matthew Hansen, Sean Healey, Michelle Hofton, George Hurt, James Kellner, Scott Luthcke, et al. The global ecosystem dynamics investigation: High-resolution laser ranging of the earth’s forests and topography. *Science of remote sensing*, 1:100002, 2020.
- [7] Laura Duncanson, James R Kellner, John Armston, Ralph Dubayah, David M Minor, Steven Hancock, Sean P Healey, Paul L Patterson, Svetlana Saarela, Suzanne Marselis, et al. Above-ground biomass density models for NASA’s Global Ecosystem Dynamics Investigation (GEDI) lidar mission. *Remote Sensing of Environment*, 270:112845, 2022.
- [8] Laura Duncanson, Mengyu Liang, Veronika Leitold, John Armston, SM Krishna Moorthy, Ralph Dubayah, Sébastien Costedoat, BJ Enquist, Lola Fatoyinbo, SJ Goetz, et al. The effectiveness of global protected areas for climate change mitigation. *Nature Communications*, 14(1):2908, 2023.
- [9] European Space Agency / Copernicus Data Space Ecosystem. Copernicus digital elevation model (dem), release 2023_1. <https://dataspace.copernicus.eu/explore-data/data-collections/copernicus-contributing-missions/collections-description/COP-DEM>, 2023. DOI:10.5270/ESA-c5d3d65; accessed: 2025-10-16.
- [10] ESA. Biomass mission: Mission overview. <https://earth.esa.int/eogateway/missions/biomass/description>, 2025. Accessed: 2025-10-16.
- [11] European Environment Agency. Forest information system for europe (fise). <https://forest.eea.europa.eu/>, 2025. Accessed: 2025-10-14.
- [12] GEO BON. What are ebvs? — essential biodiversity variables. <https://geobon.org/ebvs/what-are-ebvs/>, 2025. Accessed: 2025-10-14.
- [13] Christopher L Gilbert. The EU deforestation regulation. *EuroChoices*, 23(3):64–70, 2024.
- [14] Nguyen Tien Hoang and Keiichiro Kanemoto. Mapping the deforestation footprint of nations reveals growing threat to tropical forests. *Nature Ecology & Evolution*, 5(6):845–853, 2021.

- [15] IPBES. *Global Assessment Report on Biodiversity and Ecosystem Services of the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services*. IPBES Secretariat, Bonn, Germany, 2019. doi: 10.5281/zenodo.3831673. URL <https://doi.org/10.5281/zenodo.3831673>.
- [16] Martin Jung, Andy Arnell, Xavier De Lamo, Shaenandhoa García-Rangel, Matthew Lewis, Jennifer Mark, Cory Merow, Lera Miles, Ian Ondo, Samuel Pironon, et al. Areas of global importance for conserving terrestrial biodiversity, carbon and water. *Nature Ecology & Evolution*, 5(11):1499–1509, 2021.
- [17] Nikolai Kalischek, Nico Lang, Cécile Renier, Rodrigo Caye Daudt, Thomas Addoah, William Thompson, Wilma J Blaser-Hart, Rachael Garrett, Konrad Schindler, and Jan D Wegner. Cocoa plantations are associated with deforestation in côte d’ivoire and ghana. *Nature Food*, 4(5):384–393, 2023.
- [18] Nico Lang, Walter Jetz, Konrad Schindler, and Jan Dirk Wegner. A high-resolution canopy height model of the Earth. *Nature Ecology & Evolution*, 7(11):1778–1789, November 2023. ISSN 2397-334X. doi: 10.1038/s41559-023-02206-6. URL <https://www.nature.com/articles/s41559-023-02206-6>. Number: 11 Publisher: Nature Publishing Group TLDR: A probabilistic deep learning model is developed that fuses sparse height data from the GEDI space-borne LiDAR mission with dense optical satellite images from Sentinel-2 to improve the retrieval of tall canopies with typically high carbon stocks and has the potential to foster advances in climate, carbon and biodiversity modelling.
- [19] Wang Li, Wen-Yong Guo, Maya Pasgaard, Zheng Niu, Li Wang, Fang Chen, Yuchu Qin, and Jens-Christian Svenning. Human fingerprint on structural density of forests globally. *Nature Sustainability*, 6(4):368–379, April 2023. ISSN 2398-9629. doi: 10.1038/s41893-022-01020-5. URL <https://www.nature.com/articles/s41893-022-01020-5>. Publisher: Nature Publishing Group.
- [20] Mengyu Liang, Mariano González-Roglich, Patrick Roehrdanz, Karyn Tabor, Alex Zvoleff, Veronika Leitold, Julie Silva, Temilola Fatoyinbo, Matthew Hansen, and Laura Duncanson. Assessing protected area’s carbon stocks and ecological structure at regional-scale using gedi lidar. *Global environmental change*, 78:102621, 2023.
- [21] Suzanne M Marselis, Petr Keil, Jonathan M Chase, and Ralph Dubayah. The use of gedi canopy structure for explaining variation in tree species richness in natural forests. *Environmental Research Letters*, 17(4):045003, 2022.
- [22] Tara G Martin and James EM Watson. Intact ecosystems provide best defence against climate change. *Nature Climate Change*, 6(2):122–124, 2016.
- [23] Mirco Migliavacca, Talie Musavi, Miguel D Mahecha, Jacob A Nelson, Jürgen Knauer, Dennis D Baldocchi, Oscar Perez-Priego, Rune Christiansen, Jonas Peters, Karen Anderson, et al. The three major axes of terrestrial ecosystem function. *Nature*, 598(7881):468–472, 2021.
- [24] Lidong Mo, Constantin M Zohner, Peter B Reich, Jingjing Liang, Sergio De Miguel, Gert-Jan Nabuurs, Susanne S Renner, Johan Van Den Hoogen, Arnan Araza, Martin Herold, et al. Integrated global assessment of the natural forest carbon potential. *Nature*, 624(7990):92–101, 2023.
- [25] Elizabeth Maruma Mrema, Thomas W Crowther, Wanjira Mathai, Thomas Brooks, Sally Archibald, Pedro Brancalion, Edward B Barbier, David Obura, Victoria Reyes-García, William J Bond, et al. Ten years to restore a planet. *One Earth*, 3(6):647–652, 2020.
- [26] Maxim Neumann, Anton Raichuk, Yuchang Jiang, Mélanie Rey, Radost Stanimirova, Michelle J Sims, Sarah Carter, Elizabeth Goldman, Keith Anderson, Petra Poklukar, et al. Natural forests of the world—a 2020 baseline for deforestation and degradation monitoring. *Scientific Data*, 12(1):1715, 2025.
- [27] Jan Pauls, Max Zimmer, Una M. Kelly, Martin Schwartz, Sassan Saatchi, Philippe Ciais, Sebastian Pokutta, Martin Brandt, and Fabian Gieseke. Estimating Canopy Height at Scale, June 2024. URL <http://arxiv.org/abs/2406.01076>. arXiv:2406.01076 [cs].

- [28] Jan Pauls, Max Zimmer, Berkant Turan, Sassan Saatchi, Philippe Ciais, Sebastian Pokutta, and Fabian Gieseke. Capturing Temporal Dynamics in Large-Scale Canopy Tree Height Estimation, January 2025. URL <http://arxiv.org/abs/2501.19328>. arXiv:2501.19328 [cs].
- [29] Stuart L Pimm, Clinton N Jenkins, and Binbin V Li. How to protect half of earth to ensure it protects sufficient biodiversity. *Science advances*, 4(8):eaat2616, 2018.
- [30] Peter Potapov, Xinyuan Li, Andres Hernandez-Serna, Alexandra Tyukavina, Matthew C. Hansen, Anil Kommarreddy, Amy Pickens, Svetlana Turubanova, Hao Tang, Carlos Edibaldo Silva, John Armston, Ralph Dubayah, J. Bryan Blair, and Michelle Hofton. Mapping global forest canopy height through integration of GEDI and Landsat data. *Remote Sensing of Environment*, 253:112165, February 2021. ISSN 0034-4257. doi: 10.1016/j.rse.2020.112165. URL <https://www.sciencedirect.com/science/article/pii/S0034425720305381>.
- [31] Bernardo BN Strassburg, Alvaro Iribarrem, Hawthorne L Beyer, Carlos Leandro Cordeiro, Renato Crouzeilles, Catarina C Jakovac, André Braga Junqueira, Eduardo Lacerda, Agnieszka E Latawiec, Andrew Balmford, et al. Global priority areas for ecosystem restoration. *Nature*, 586(7831):724–729, 2020.
- [32] Frank Thonfeld, Ursula Gessner, Stefanie Holzwarth, Jennifer Kriese, Emmanuel Da Ponte, Juliane Huth, and Claudia Kuenzer. A first assessment of canopy cover loss in germany’s forests after the 2018–2020 drought years. *Remote Sensing*, 2022.
- [33] Jamie Tolan, Hung-I Yang, Benjamin Nosarzewski, Guillaume Couairon, Huy V. Vo, John Brandt, Justine Spore, Sayantan Majumdar, Daniel Haziza, Janaki Vamaraju, Theo Moutakanni, Piotr Bojanowski, Tracy Johns, Brian White, Tobias Tiecke, and Camille Couprie. Very high resolution canopy height maps from RGB imagery using self-supervised vision transformer and convolutional decoder trained on aerial lidar. *Remote Sensing of Environment*, 300:113888, January 2024. ISSN 0034-4257. doi: 10.1016/j.rse.2023.113888. URL <https://www.sciencedirect.com/science/article/pii/S003442572300439X>. TLDR: This paper presents the first high-resolution canopy height map concurrently produced for multiple sub-national jurisdictions, and introduces a post-processing step using a convolutional network trained on GEDI observations.
- [34] Thomas Vanneste, Leen Depauw, Emiel De Lombaerde, Camille Meeussen, Sanne Govaert, Karen De Pauw, Pieter Sanczuk, Kurt Bollmann, Jörg Brunet, Kim Calders, et al. Trade-offs in biodiversity and ecosystem services between edges and interiors in european forests. *Nature ecology & evolution*, 8(5):880–887, 2024.
- [35] James EM Watson, Tom Evans, Oscar Venter, Brooke Williams, Ayesha Tulloch, Claire Stewart, Ian Thompson, Justina C Ray, Kris Murray, Alvaro Salazar, et al. The exceptional value of intact forest ecosystems. *Nature ecology & evolution*, 2(4):599–610, 2018.
- [36] Sarah R Weiskopf, Forest Isbell, Maria Isabel Arce-Plata, Moreno Di Marco, Mike Harfoot, Justin Johnson, Susannah B Lerman, Brian W Miller, Toni Lyn Morelli, Akira S Mori, et al. Biodiversity loss reduces global terrestrial carbon storage. *Nature Communications*, 15(1):4354, 2024.
- [37] Karina Winkler, Richard Fuchs, Mark Rounsevell, and Martin Herold. Global land use changes are four times greater than previously estimated. *Nature communications*, 12(1):2501, 2021.
- [38] Daniele Zanaga, Ruben Van De Kerchove, Dirk Daems, Wanda De Keersmaecker, Carsten Brockmann, Grit Kirches, Jan Wevers, Oliver Cartus, Maurizio Santoro, Steffen Fritz, et al. Esa worldcover 10 m 2021 v200. 2022.
- [39] X Zhang, MA Friedl, and GM Henebry. VIIRS global land surface phenology product user guide; United States Geological Survey: Reston, VA, USA, 2017.