

1
2
3
4 1 **Wild *Zea* as a Model System for Population and Ecological**
5
6
7 2 **Genomics**
8
9

10 3 Matthew B. Hufford¹, Paul Bilinski¹, Tanja Pyhäjärvi¹, Jeffrey Ross-Ibarra^{1,2}
11
12

13 4
14

15 5 ¹ Plant Sciences Department, University of California, Davis, CA 95616, USA
16

17 6 ² The Genome Center and Center for Population Biology, University of California,
18
19 7 Davis, CA 95616, USA
20
21

22 8
23

24 9 *Corresponding author:* Ross-Ibarra, J. (rossibarra@ucdavis.edu)
25
26

27 10
28

29 11 Keywords: population genomics; crop wild relatives; domestication; local
30
31

32 12 adaptation; introgression; *Zea*
33
34

35 13
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Abstract

As the cost of next-generation sequencing diminishes and genomic resources improve, crop wild relatives are well positioned to make major contributions to the field of ecological genomics via full-genome resequencing and reference-assisted *de novo* assembly of genomes of plants from natural populations. The wild relatives of maize, collectively known as teosinte, are a more varied and representative study system than many other model flowering plants. In this review of the population and ecological genomics of the teosintes, we highlight recent advances in the study of maize domestication, introgressive hybridization, and local adaptation and discuss future prospects for applying the genomic resources of maize to this intriguing group of species. The maize/teosinte study system is an excellent example of how crops and their wild relatives can bridge the model/non-model gap.

The Genomic Era and Crop Wild Relatives

Over the past decade, reference genome sequences have been completed for numerous crops (*e.g.*, see Table 1 of [1]). While the purpose of such sequencing efforts has largely been crop improvement, these newly developed resources also hold great promise for population genomic studies in related wild taxa. Genetic tools developed for crops have been successfully applied in ecological and evolutionary studies of wild relatives [2-6]. However, genomic studies of natural populations have trailed behind those of domesticated and other model species. The opportunities afforded by wild relatives' taxonomic proximity to reference genomes

are only beginning to be realized as the diminishing cost of next-generation sequencing technologies and availability of large-scale genotyping arrays make population-scale data feasible. Here we describe recent progress in genomic studies in the genus *Zea* (Family *Poaceae*), which comprises an ecologically diverse collection of species including domesticated maize and its closest wild relatives. These taxa have large, complex genomes and have already contributed much to modern plant genetics (Box 1). The past few years have seen the publication of the maize genome [7], full-genome resequencing of hundreds of domesticated and wild *Zea* individuals [8-10], and characterization of transcriptome-wide patterns of expression [11-14]. In describing recent advances we will: 1) Present wild *Zea* (collectively known as the teosintes) as an ideal study system for ecological and population genomics; 2) Describe long-standing questions that are benefitting from the application of genome-wide data, including maize domestication from teosinte, reciprocal gene flow between teosinte and maize, and local adaptation of teosinte populations; and 3) Discuss future research prospects for this fascinating research system.

The Teosinte Taxa

The teosintes are a group of primarily outcrossing species in the genus *Zea* endemic to a region stretching from northern Mexico to Central America (Figure 1A & 1B). These species occupy a wide range of habitats and have diverse ecologies (Figure 1C; Table 1; Box 1). The genus *Zea* includes five species (Figure 1D). The

1
2
3
4 60 annual species *Zea mays* consists of four subspecies: 1) ssp. *mays*, the domesticated
5
6 61 maize, also known as corn; 2) ssp. *parviglumis*, the direct progenitor of maize [15],
7
8 62 found in the mesic low and middle elevations of southwest Mexico; 3) ssp. *mexicana*,
9
10 63 distributed across the cooler high elevations of the Mexican Central Plateau; and 4)
11
12 64 ssp. *huehuetenangensis*, found only in western Guatemala. The diploid *Zea*
13
14 65 *diploperennis* and tetraploid *Zea perennis* are closely related perennial species
15
16 66 narrowly distributed along the mountain slopes of western Mexico. Finally, *Zea*
17
18 67 *luxurians* and *Zea nicaraguensis* are annual, flood-tolerant species that grow in
19
20 68 southeastern Guatemala and the Pacific coast of Nicaragua [16,17]. We refer to taxa
21
22 69 hereafter solely by species or subspecies name. Exploration continues to uncover
23
24 70 divergent populations of several *Zea* taxa [18], suggesting that there may be more to
25
26 71 learn about the ecological and taxonomic diversity of the genus.
27
28
29
30
31
32
33

34 72 The teosintes garner much attention due to their close relationship with
35
36 73 maize and their relevance to the study of plant domestication, but they also
37
38 74 represent a particularly attractive study system for ecological and evolutionary
39
40 75 genomics in their own right. When compared to other model angiosperms, the
41
42 76 larger size, higher content of repetitive elements, and greater complexity of *Zea*
43
44 77 genomes [8] make them more representative of flowering plants (Box 1; for a
45
46 78 comparison of wild *Zea* to other emerging model systems see Table 1 of [19]).
47
48 79 Additionally, in certain portions of their ranges, numerous large and relatively
49
50 80 undisturbed natural populations of the teosinte species can be found spanning steep
51
52 81 environmental gradients ([20], M. B. Hufford *et al.*, unpublished). These particular
53
54 82 characteristics make the teosintes well suited for studies of local adaptation and
55
56
57
58
59
60
61
62
63
64
65

natural selection. The teosintes also provide opportunities for the study of evolution through hybridization due to the history of diffusion of maize from its domestication center in the Mexican lowlands into sympatry with interfertile, locally adapted teosinte taxa in novel environments [15,21]. Finally, the teosintes include examples of both ancient [22] and recent [23] polyploidy as well as species with varied life-history strategies (*e.g.*, perennial vs. annual growth habit and asexual vs. sexual reproduction; Table 1). As we will describe in the following sections, the promise of genomic studies in this compelling study system is now being realized and answers to numerous ecological and evolutionary questions will soon be within reach due to the opportunities provided by newly emerging genomic technologies.

Population Genomics of Teosinte

Evolutionary Genomics of Maize Domestication

For much of the twentieth century, the identity of the direct progenitor of maize was actively debated (for a summary see [24,25]). Genetic analysis of allozyme [26,27] and microsatellite data [15] largely settled the dispute, convincingly demonstrating a single domestication of maize in the Balsas River Basin of southwest Mexico from the teosinte *parviglumis*. One riddle in the geography of maize domestication remained, however: maize landraces (*i.e.*, varieties grown by local farmers) most genetically similar to *parviglumis* were from the Mexican highlands, where *parviglumis* is not found [15]. This contradiction was

resolved using a sizeable panel of single nucleotide polymorphisms (SNPs) typed in *parviglumis*, *mexicana*, and landraces from across the Americas [28]. *Parviglumis* and *mexicana* were less differentiated from each other than either was from maize, and much more admixture was found between maize and *mexicana* than between maize and *parviglumis*. Somewhat counter-intuitively, introgression from *mexicana* was shown as the likely cause of the genetic similarity between highland maize landraces and *parviglumis*. In contrast to comparisons with *parviglumis*, when inferred allele frequencies of ancestral maize were compared to extant maize landraces, the highest genetic similarity between *parviglumis* and maize was found with landraces from lowland West Mexico, the putative maize center of origin [28].

In addition to these phylogeographic questions, evolutionary biologists studying maize have been keenly interested in identifying the genes responsible for the radical phenotypic transformation that occurred during maize domestication. One of the more notable differences between maize and teosinte is plant architecture: whereas *parviglumis* often produces numerous lateral branches and tillers (basal branches), maize typically possesses an unbranched single stalk. Much of this disparity in plant architecture is caused by the gene *teosinte branched1* (*tb1*), a transcription factor that represses the growth of axillary organs [29]. The causal polymorphism at the *tb1* locus has recently been shown to be a single transposable element insertion in the upstream regulatory region of *tb1* that acts as an enhancer of gene expression [30]. The allele that confers the maize phenotype was also found to be segregating in some teosinte populations [30], suggesting domestication has, at least in certain instances, occurred from standing variation in the wild progenitor

1
2
3
4 129 (for additional potential examples see [31,32]). Maize is also strikingly distinct from
5
6 130 *parviglumis* and other teosintes in the structure of its female infructescence:
7
8
9 131 teosinte possesses numerous spikelets of disarticulating seed that are covered in a
10
11 132 hard protective casing (glume), whereas maize produces a small number of
12
13
14 133 infructescences with naked grain bound to a cob, making it entirely dependent on
15
16 134 humans for propagation. Variation in grain exposure has been shown to be largely
17
18
19 135 due to a single locus, *teosinte glume architecture* (*tga1*; [33]). Several other
20
21 136 domestication loci controlling traits such as flowering time, kernel weight and
22
23
24 137 quality, and prolificacy (*i.e.*, number of ears) have been identified through a
25
26 138 combination of top-down (phenotype to selected genomic region; [34,35]) and
27
28
29 139 bottom-up (selected genomic region to phenotype; [36-39]) genetic approaches (see
30
31 140 [40] for a contrast and discussion of these approaches).

32
33
34 141 Genomic technologies are now revealing that domestication involved more
35
36 142 extensive changes between maize and teosinte than merely the fixation of alternate
37
38
39 143 alleles at a few large-effect loci. The availability of genome-wide resequencing data
40
41 144 from over 100 teosinte, landrace, and modern improved lines as part of the maize
42
43
44 145 HapMap2 project [8] has recently allowed a population genomic analysis of maize
45
46 146 domestication and improvement [12]. This work identified extended regions of
47
48
49 147 differentiation throughout the genome, revealed patterns consistent with stronger
50
51 148 selection during domestication (the transition from teosinte to maize landraces)
52
53
54 149 than during subsequent crop improvement (the transition from open-pollinated
55
56 150 landraces to modern inbred lines) and identified a number of compelling candidate
57
58
59 151 genes with stronger signatures of selection than the canonical loci *tb1* and *tga1* [12].
60
61
62
63
64
65

1
2
3
4 152 Comparison of polymorphism data in maize landraces and teosinte confirmed a
5
6 153 substantial loss (17%) of diversity during the domestication bottleneck, but a
7
8
9 154 genome-wide excess of rare alleles provided evidence that diversity has begun to
10
11
12 155 recover. A slower recovery was found within genes, likely due to the effects of
13
14 156 purifying selection. Several genomic regions in maize landraces showed signs of
15
16 157 introgression from the highland teosinte *mexicana*, suggesting an important role for
17
18
19 158 gene flow during post-domestication maize diffusion. The magnitude and level of
20
21
22 159 variation of gene expression were also shown to have changed significantly during
23
24 160 domestication [12] with substantial re-wiring of coexpression networks in the
25
26 161 maize transcriptome relative to *parviglumis* [14]. The maize HapMap2 project also
27
28
29 162 revealed large-scale, structural changes across the genomes of *Zea* (Box 1).
30
31
32 163 Pervasive structural variants (SVs) were identified as regions showing differences in
33
34 164 the presence or copy number of segments of the genome. Finally, genome size
35
36 165 differences within the species *Z. mays* were found to be caused by segregation of
37
38
39 166 heterochromatic regions known as chromosomal knobs [8], whereas between-
40
41
42 167 species differences (*Z. mays* compared to *luxurians* [41] and the closely-related
43
44 168 taxon *Tripsacum dactyloides* [8]) have been shown to be due largely to transposable
45
46
47 169 element content.

48
49 170

50 51 171 *Introgression Between Teosinte and Maize*

52
53
54 172

55
56 173 Subsequent to domestication, teosinte populations have persisted in both
57
58
59 174 allopatry and sympatry with maize. Hybridization between maize and teosinte is

1
2
3
4 175 well documented where they co-occur [42-44], but genetic barriers to hybridization
5
6 176 are also well known. At least three loci are known to act as pre-zygotic crossing
7
8
9 177 barriers between maize and teosinte: *Teosinte crossing barrier 1 (Tcb1)* and
10
11 178 gametophyte factors (*Ga*) 1 and 2 [45-47]. The biochemical mechanism underlying
12
13
14 179 the action of these loci remains unclear, but alleles conferring incompatibility at *Ga1*
15
16 180 and *Tcb1* appear to be present at higher frequency in *mexicana* than in *parviglumis*
17
18
19 181 populations [47].
20

21
22 182 Hybridization barriers may dramatically limit the rate of gene flow, yet
23
24 183 theory predicts that if certain combinations of genotypes show increased fitness,
25
26 184 introgression can occur despite barriers to gene exchange [48]. Recent evidence
27
28
29 185 suggests reciprocal introgression between the partially incompatible subspecies
30
31
32 186 *mexicana* and maize may be a compelling example of such gene flow. Following its
33
34 187 domestication in the lowlands of southwest Mexico from *parviglumis*, maize diffused
35
36 188 to the highlands of the Mexican Central Plateau where it encountered dramatically
37
38
39 189 different environmental conditions and came into sympatry with the highland
40
41 190 teosinte *mexicana*. Modern highland maize and *mexicana* share multiple phenotypic
42
43
44 191 traits (*e.g.*, abundant macrohairs and leaf pigmentation; [44]). This shared
45
46 192 morphology has been attributed both to introgression from maize into *mexicana*
47
48
49 193 conferring crop mimicry [49] and to *mexicana* introgression into maize conferring
50
51 194 highland adaptation under the assumption that shared traits are beneficial under
52
53
54 195 highland conditions [50]. The application of molecular markers has revealed
55
56 196 considerable evidence for gene flow in both directions, suggesting as much as 20%
57
58
59
60
61
62
63
64
65

of highland maize genomes are derived from *mexicana* and approximately 10% of *mexicana* genomes are derived from maize [28].

Recent work has taken advantage of the availability of genome-wide SNP typing to evaluate the genomic imprint of introgression in sympatric populations of highland Mexican maize and *mexicana* [51]. Numerous genomic regions of introgression were identified in both directions of gene flow yet domestication loci appeared particularly resistant to introgression. Interestingly, regions of *mexicana*-to-maize introgression were shared across populations suggesting either ancient gene flow or multiple independent introgression events at the same loci. These shared regions of introgression from *mexicana* into maize were found to colocalize with previously identified QTL for highland traits [52], suggesting a possible role for *mexicana* in conferring maize adaptation to highland conditions. A growth chamber experiment simulating highland conditions and including maize populations with and without *mexicana* introgression at these shared loci showed significantly higher leaf sheath macrohair counts, extent of pigment, and plant height in populations with a high frequency of *mexicana*-to-maize introgression [51].

Local Adaptation in Teosinte

Local adaptation occurs when resident genotypes have higher fitness than genotypes from other locations [53]. Currently, there are only a handful of examples in which the genetic basis of local adaptation is well understood. Because of their sessile nature, plants are particularly well suited for studies of local adaptation [19],

220 and genome-wide data have recently been analyzed in plant populations across
221 wide geographic ranges [54,55].

222 Among the teosintes, *mexicana* and *parviglumis* are the best candidates for
223 studies of local adaptation. Both have widespread geographic distributions (Figure
224 1A & 1B) consisting of hundreds of distinct populations [2,20,56] that display
225 varying levels of genetic differentiation [2,57], from very fine-scale structure (F_{ST} : 1-
226 2%) within a site [58,59] to more pronounced differentiation between subspecies
227 (F_{ST} : ~15%; [60]). Populations of *parviglumis* and *mexicana* also span a wide
228 climatic range of temperature and precipitation (Table 1). Phenotypically, these
229 subspecies are differentiated based on the smaller spikelets and fruitcases and more
230 open tassels of *parviglumis* in comparison to *mexicana* [16] and by the hairier leaf
231 sheaths and higher pigment content found in *mexicana* [52]. Within subspecies,
232 individual *parviglumis* and *mexicana* populations are known to vary in several
233 morphological and phenological characteristics [56,61,62]. It is not yet clear
234 whether population-level phenotypic differences are due to local adaptation or if
235 they are rather byproducts of genetic drift.

236 Recent studies have investigated the molecular variation underlying local
237 adaptation in teosinte. A study of variation in plant immunity genes across six
238 *parviglumis* populations found little evidence for widespread geographically
239 variable selection but did observe diversity patterns suggesting localized natural
240 selection in the gene *wip1* [63]. *wip1*, a protease inhibitor, is known to interfere
241 with the dietary proteins of insects, and selection at this locus could represent local
242 adaptation to herbivory. A second study of SNP diversity in *mexicana* and

243 *parviglumis* identified *Inv1n*, a 50-Mb chromosomal inversion found segregating in
244 33 *parviglumis* populations [64]. The frequency of *Inv1n* was shown to vary along an
245 altitudinal cline (*i.e.*, a continuous gradient) and associate with environmental and
246 morphological characters. Such inversions are predicted to be beneficial when they
247 capture locally adapted alleles or when the breakpoints of the inversions themselves
248 are beneficial [65]. Most recently, a genome-wide analysis of 36,719 SNPs typed in
249 10 *mexicana* and 11 *parviglumis* populations identified candidate loci underlying
250 local adaptation [66]. Contrary to previous findings from *Arabidopsis thaliana* [55],
251 enrichment of non-genic SNPs was found among candidates, in line with the
252 observation that a substantial proportion of phenotypic associations in maize are
253 found in noncoding regions [8]. In addition, several newly identified inversion
254 polymorphisms showed an excess of differentiation among populations and
255 subspecies along with enrichment for association with environmental variables such
256 as temperature and altitude. The complexity and size of the *Zea* genome (Box 1)
257 may provide ample opportunities for such large-scale chromosomal rearrangements
258 to occur. Finally, the study revealed that factors such as hierarchical population
259 structure and uneven distribution of environmental variation can bias detection of
260 locally adaptive loci based on a single statistical method, underscoring the
261 importance of utilizing multiple approaches.

262

263 **Future Prospects**

264

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

265 The teosintes provide opportunities to examine many compelling questions
266 at the genome level using the vast number of resources developed in maize (Box 2).
267 As the cost of high-throughput sequencing continues to decrease and *de novo*,
268 reference-assisted assembly becomes commonplace in teosinte, genomic studies can
269 move farther beyond maize, delving more deeply into the evolutionary histories of
270 these taxa. For example, the species *luxurians* and *nicaraguensis* are uniquely
271 adapted to the flooded and coastal conditions of Central America. Unlike other
272 teosintes, these taxa develop aerenchyma (air channels that allow gas exchange
273 between submerged and above-water organs of a plant) both in flooded and non-
274 flooded conditions [67], and multiple QTL have already been identified for this trait
275 [68]. Past studies have suggested potential hybridization between these teosintes
276 and local maize landraces [49,50,60], and additional field and genome-wide
277 analyses should assess evidence for adaptive introgression for flood tolerance in
278 landraces at loci encoding these traits.

279 The teosintes also provide an excellent system in which to study the genetics
280 of speciation. The genus appears to have radiated relatively recently, and shared
281 polymorphism and ongoing gene flow are evident across species [60]. Perhaps of
282 greatest interest would be field-based studies of local adaptation (*e.g.*, development
283 of near isogenic lines and reciprocal transplant experiments), incipient speciation,
284 and admixture between *parviglumis* and *mexicana*. These subspecies appear to have
285 diverged only recently, but already exhibit locally adaptive phenotypes, and gene
286 flow between them is limited to a narrow hybrid zone [2,66]. Studies of gene flow

287 and reinforcement within this zone could shed light on the process of divergence in
288 these subspecies.

289 Yet another potential topic of investigation in *Zea* is the recent
290 autopolyploidization (within the last few hundred thousand years; [23]) of *perennis*.
291 Investigation of the more ancient polyploidization event ancestral to all of *Zea* has
292 shown that genome fractionation has been biased toward a single subgenome and is
293 ongoing, distinguishing even modern inbred maize lines [69]. *Perennis* presents an
294 opportunity to investigate incipient fractionation/diploidization in natural
295 populations by utilizing newly developed resources to assess genome-wide patterns
296 of gene expression, DNA methylation, and deletion.

297 Genomic resources could also be brought to bear on the evolution of the
298 perennial life history in *diploperennis* and *perennis*, a feature that distinguishes these
299 species from the remaining annual *Zea* taxa. A previous QTL investigation of
300 *diploperennis* and *parviglumis* did not find large effect loci controlling traits
301 associated with perennialism, and very few genes were linked to the evolution of the
302 perennial growth habit [70]. Full genome resequencing may allow population-
303 genetic-based assessment of selection, but it seems likely that increased marker
304 density would be better exploited by multiparent mapping populations (*e.g.*,
305 [71,72]) of annual/perennial taxa that can take advantage of a large number of
306 recombination events to provide increased resolution. Full transcriptome patterns
307 of gene expression would also be valuable data for identifying loci involved in the
308 contrasting growth habits across *Zea*. Once these full-genome data are generated
309 and we have a better idea about the loci involved in *Zea* perennialism, many

1
2
3
4 310 fascinating ecological and evolutionary questions can be investigated. For example:
5
6 311 Do genes associated with perennialism explain these species' capacity to tolerate
7
8 312 conditions outside the normal growing season of the annual *Zea* (*e.g.*, cold-tolerance
9
10 313 or drought-resistance loci)? To what degree is there overlap with the pathways and
11
12 314 orthologous genes involved in perennialism in closely related taxa such as rice and
13
14 315 sorghum?

15
16 316 While a number of fascinating evolutionary questions can be addressed
17
18 317 within *Zea* on a species-by-species basis, the genus as a whole can also be seen as an
19
20 318 ideal study system for investigating broader and potentially far-reaching topics. For
21
22 319 instance, *Zea* genomes are more representative than the genomes of most current
23
24 320 angiosperm models in that they are comprised primarily of repetitive elements
25
26 321 (transposable elements (TE's), microsatellites, centromere and chromosomal knob
27
28 322 repeats) [73]. Within *Zea*, the evolution of repetitive elements can be studied over
29
30 323 hundreds of thousands of years, revealing epochs of transposition of TE's, the
31
32 324 expansion and contraction of knob and centromere repeat arrays, and the structural
33
34 325 and functional roles of this largely ignored fraction of the genome. *Zea* species also
35
36 326 have a wide range of effective population sizes, allowing for comparisons of the
37
38 327 efficiency of natural selection and the effects of rare variants across closely related
39
40 328 taxa. Association studies in the wild taxa of *Zea* will also provide greater resolution
41
42 329 for the genetic architecture of complex traits than is currently possible in their
43
44 330 domesticated congeners due to the increased number of recombination events in
45
46 331 natural populations and the rapid decay of linkage disequilibrium [8].
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4 332 Finally, much of the proposed evolutionary genomic work in the teosintes
5
6 333 will likely feed back into the improvement of maize (Figure 2). For example,
7
8
9 334 temperature-related highland adaptation loci could be important for maize breeding
10
11 335 in the face of climate change, flood-tolerance loci identified in introgression or
12
13
14 336 continued QTL studies of *luxurians* and *nicaraguensis* could help mitigate crop losses
15
16 337 due to flooding [74], and genomic studies of *diploperennis* and *perennis* may
17
18
19 338 facilitate the development of perennial maize, a resource that could potentially
20
21 339 reduce inputs and prove particularly useful in agroecosystems of the developing
22
23 340 world. In this way, the model/non-model gap will be bridged not only through
24
25 341 application of maize-derived technologies to teosinte, but also through valuable
26
27 342 contributions from the rich evolutionary past of *Zea* to the future of maize.
28
29
30
31
32

33 343

34 344 **Acknowledgements**

35
36 345 We thank John Doebley, Sofiane Mezmouk, Peter Morrell, Anna O'Brien, Shohei
37
38 346 Takuno, and Maud Tenaillon for helpful comments on a previous draft. MBH was
39
40 347 supported by US-NSF grant IOS-0922703, PB was supported by the UC Davis
41
42 348 Beatrice Oberly and S. Atwood McKeehan Fellowship and the UC Davis Plant
43
44 349 Sciences Graduate Student Research Assistantship, and TP received support from
45
46 350 the Academy of Finland. JR-I would like to acknowledge support from US-NSF grant
47
48 351 IOS-0922703 and USDA–National Institute of Food and Agriculture grant 2009-
49
50 352 01864.
51
52
53
54
55
56
57
58

59 353

60 354 **Glossary**

355

356 **Allopatry:** describes geographically and reproductively isolated populations

357 **Chromosomal knob:** heterochromatic DNA in *Zea* comprised primarily of two

358 classes of tandem satellite repeats, 180bp and the 350bp TR-1. Knobs in maize are

359 found both in the middle and on the ends of chromosomal arms, and have been

360 associated with meiotic drive

361 **Fine-scale/cryptic structure:** non-random barriers to exchange of genetic material

362 among individuals in close proximity that are difficult to detect based on observable

363 characters

364 **Fractionation:** the mutational process leading to the reduction of gene number

365 following a whole-genome duplication event

366 **Inferred allele frequency:** frequency of alleles in the ancestral population

367 estimated based on knowledge of current genotypes, population structure, and the

368 process of genetic drift

369 **Introgression:** gene flow and persistence of genetic material from one population

370 or species into the gene pool of another

371 **Purifying selection:** a process of natural selection in which deleterious alleles are

372 removed from the gene pool

373 **Sympatry:** describes populations which inhabit overlapping geographic areas

374 allowing mating

375

376 **Boxes**

377

1
2
3
4 **378 Box 1: The Pan-*Zea* Genome.**
5

6
7 379 Early studies in maize were the first to physically map genes onto
8
9 380 chromosomes and reveal the phenomenon of crossing over [75]. Subsequent
10
11 381 cytological studies showed large-scale structural differences in *Zea* genomes due to
12
13
14 382 varying heterochromatic knob content and chromosomal inversions [T. A. Kato
15
16 383 Yamakake, PhD thesis, University of Massachusetts, 1975; 76]. Array- and
17
18
19 384 sequence-based analyses have also demonstrated substantial genetic and structural
20
21
22 385 variation across taxa [8,12,41,77]. However, the recent origin of the genus [60] and
23
24 386 gene flow among taxa [2], when combined with evidence for structural variation
25
26
27 387 [78,79], suggest researchers will be able to describe core and dispensable fractions
28
29 388 of a Pan-*Zea* genome [80].
30

31
32 389 A large-scale effort is underway to investigate low-copy genic regions in *Zea*
33
34 390 species. Currently, the high-confidence filtered gene set of the 2.1-Gb maize genome
35
36 391 (maizesequence.org; release 5b.60) includes ~40,000 genes. Microarray data
37
38
39 392 indicated ~10% of genes across *Zea mays* accessions were associated with
40
41
42 393 structural variants [81,82]; however, estimates from next-generation sequencing of
43
44 394 103 *Zea* genomes suggested structural variation in as many as 70% of genes and
45
46
47 395 identified significant associations between variants and phenotype [8]. Analyses of
48
49 396 syntenic regions between sorghum and maize show that genes with lower
50
51
52 397 expression are more likely to be lost, suggesting a potential mechanism behind
53
54 398 observed variation in gene content in inbred lines [69,82].
55

56 399 Extensive variation is not limited to the genic fraction of *Zea* genomes.
57
58
59 400 Cytological studies of repetitive elements (knobs, transposable elements (TE's),
60
61
62
63
64
65

centromere repeats), which comprise >85% of the genome, have shown dramatic differences across *Zea* [79,83]. Sequence-based analyses, however, indicate that whereas individual families of TE's have changed in abundance in recent history [12], the relative frequencies of most TE's appear similar between maize and its sister species *luxurians* [41]. Sequence analysis also confirmed predictions [84,85] that chromosomal knob content correlates positively with genome size within *Z. mays*, whereas transposable elements account for genome size differences between *Z. mays* and *Tripsacum* [8]. Fluorescent *in situ* hybridization has previously shown visible differences in centromere repeat (CentC) abundance between *parviglumis* and maize, and these differences can now be confirmed and quantified using low-coverage, whole-genome sequencing (c.f. [41]).

Analysis of shared and non-shared genomic regions across *Zea* individuals has begun to describe the core and dispensable fractions of the Pan-*Zea* genome [80]. The emerging picture is much more fluid than previously expected and challenges the notion of genome conservation even at the intraspecies level. Because of its great diversity, a Pan-*Zea* genome would likely serve as a model for studying genomic evolution.

Box 1 Figure Legends

Box 1, Figure 1. Genomic content of different classes of structural elements in maize and its relatives. Genome size, represented as a line graph, is plotted on the left axis. Repetitive element abundance was standardized against maize. B73 was used as the single reference line for maize, while *parviglumis* (N=9) and *mexicana*

424 (N=2) are averages. A single *Tripsacum dactyloides* individual was also used. Data
425 for figure were obtained from [8].

426

427 **Box 1, Figure 2.** Fluorescence *in situ* hybridization in two closely related *Zea* taxa,
428 *parviglumis* and *mexicana*. Chromosome number is indicated on the upper axis, and
429 homologous chromosomes are shown. Karyotype probes include: 180-bp knob
430 (blue), TR-1 knob (pseudo-colored white), CentC and subtelomere 4-12-1 (green),
431 5S ribosomal gene (yellow), Cent4 (orange), NOR (bluegreen), and TAG
432 microsatellite (1-26-2) and subtelomere 1.1 (red). Regions of overlapping probes
433 exhibit a corresponding color change [83].

434

435 **Box 2:** Genetic tools for studying maize.

436 **Maize reference genome** (maizesequence.org) The version AGPv2 assembly of
437 cultivar B73 consists of 2.1×10^9 bases, defines 39,656 genes and reports 63,540
438 transcripts with high confidence. The genome sequence facilitates both full-genome
439 resequencing of teosinte lines and annotation, which is essential for evaluating the
440 functional consequences of observed genetic variation. The reference genome can
441 also be used in expression analyses through read mapping of sequenced RNA

442 **Genotyping methods** The Illumina MaizeSNP50 single nucleotide polymorphism
443 (SNP) array [86] and Genotyping by Sequencing (GBS) [87] can be used for high
444 throughput genotyping in teosinte.

1
2
3
4 445 **Zinc-finger nuclease genome modification** [88] facilitates precise addition and
5
6 446 disruption of target nucleotides and could be used to transfer, for example, teosinte
7
8
9 447 alleles into maize genomes.
10

11 448

12
13
14 449 **Databases and Resources:**

15
16 450 **MaizeGDB** (maizegdb.org): incorporates maize genetic data from various public
17
18
19 451 resources in a comprehensive database and genome browser.
20

21 452 **Panzea** (panzea.org): numerous databases of genotypic and phenotypic data for
22
23
24 453 association mapping and evolutionary analyses.
25

26 454 **Genomaize** (genomaize.org/): hosts multiple tools for maize biologists, including a
27
28
29 455 genome browser for nucleosome occupancy and a web tool for *in silico* PCR.
30

31 456 **Ac/Ds tagging and Mu insertion databases** (plantgdb.org/prj/AcDsTagging/)
32
33
34 457 (teosinte.uoregon.edu/mu-illumina/): these resources list loci with potentially
35
36
37 458 disruptive transposon insertions and provide seeds of the corresponding maize
38
39 459 lines [89,90].
40

41 460 **The maize transposable element (TE) database** (maizetedb.org): describes
42
43
44 461 hundreds of maize TE families and has been used to estimate abundance of
45
46
47 462 individual TE families in maize and related taxa [8,41].
48

49 463 **Maize transcription atlas** (plexdb.org): describes gene expression in 60 tissues
50
51
52 464 and developmental stages for tens of thousands of genes in the reference cultivar
53
54 465 B73 [13].
55

56 466 **MaizeCyc** (pathway.gramene.org/gramene/maizecyc.shtml): catalogs metabolic
57
58
59 467 and transport pathways of maize.
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

468 **QTeller** (qteller.com): a tool to summarize function, expression, and syntenic
469 information from maize genes in a given genomic interval.
470 **CoGe** (genomevolution.org): tools for plant comparative genomics, including well-
471 known classical genes in maize, and sets of maize orthologs syntenic with other
472 grasses
473 **Maize Stock Center** (maizecoop.cropsci.uiuc.edu): collection of maize genetic
474 stocks, including information on known mutants
475
476

References

- 1 Morrell, P.L. *et al.* (2012) Crop genomics: advances and applications. *Nat. Rev. Genet.* 13, 85–96
- 2 Fukunaga, K. *et al.* (2005) Genetic diversity and population structure of teosinte. *Genetics* 169, 2241–2254
- 3 Martinez-Castillo, J. *et al.* (2006) Structure and genetic diversity of wild populations of lima bean (*Phaseolus lunatus* L.) from the Yucatan Peninsula, Mexico. *Crop Sci.* 46, 1071–1080
- 4 Peleg, Z. *et al.* (2008) Genetic structure of wild emmer wheat populations as reflected by transcribed versus anonymous SSR markers. *Genome* 51, 187–195
- 5 Sapir, Y. *et al.* (2007) Patterns of genetic diversity and candidate genes for ecological divergence in a homoploid hybrid sunflower, *Helianthus anomalus*. *Mol. Ecol.* 16, 5017–5029
- 6 Zheng, X.M. and Ge, S. (2010) Ecological divergence in the presence of gene flow in two closely related *Oryza* species (*Oryza rufipogon* and *O. nivara*). *Mol. Ecol.* 19, 2439–2454
- 7 Schnable, P.S. *et al.* (2009) The B73 maize genome: complexity, diversity, and dynamics. *Science* 326, 1112–1115
- 8 Chia, J.-M. *et al.* (2012) Maize HapMap2 identifies extant variation from a genome in flux. *Nat. Genet.* 44, 803–807
- 9 Gore, M.A. *et al.* (2009) A first-generation haplotype map of maize. *Science* 326, 1115–1117
- 10 Jiao, Y. *et al.* (2012) Genome-wide genetic changes during modern breeding of maize. *Nat. Genet.* 44, 812–815
- 11 Li, P. *et al.* (2010) The developmental dynamics of the maize leaf transcriptome. *Nat. Genet.* 42, 1060–1067
- 12 Hufford, M.B. *et al.* (2012) Comparative population genomics of maize domestication and improvement. *Nat. Genet.* 44, 808–811
- 13 Sekhon, R.S. *et al.* (2011) Genome-wide atlas of transcription during maize development. *Plant J.* 66, 553–563
- 14 Swanson-Wagner, R. *et al.* (2012) Reshaping of the maize transcriptome by domestication. *P. Natl. Acad. Sci. USA* 109, 11878–11883
- 15 Matsuoka, Y. *et al.* (2002) A single domestication for maize shown by multilocus microsatellite genotyping. *P. Natl. Acad. Sci. USA* 99, 6080–6084
- 16 Iltis, H.H. and Doebley, J.F. (1980) Taxonomy of *Zea* (Gramineae). II. Subspecific categories in the *Zea mays* complex and a generic synopsis. *Am. J. Bot.* 67, 994–1004
- 17 Doebley, J. (1990) Molecular systematics of *Zea* (Gramineae). *Maydica* 35, 143–150
- 18 Sanchez, G.J.J. *et al.* (2011) Three new teosintes (*Zea* spp., *Poaceae*) from Mexico. *Am. J. Bot.* 98, 1537–1548
- 19 Anderson, J.T. *et al.* (2011) Evolutionary genetics of plant adaptation. *Trends Genet.* 27, 258–266
- 20 Sanchez, G.J.J. and Corral, J.A.R. (1997) Teosinte distribution in Mexico. In *Gene flow among maize landraces, improved maize varieties and teosinte: implications*

- for transgenic maize (Serratos, J.A., Willcox, M.C., and Castillo-González, F., eds), pp. 18-36, CIMMYT
- 21 Vigouroux, Y. *et al.* (2008) Population structure and genetic diversity of new world maize races assessed by DNA microsatellites. *Am. J. Bot.* 95, 1240–1253
 - 22 Gaut, B.S. and Doebley, J.F. (1997) DNA sequence evidence for the segmental allotetraploid origin of maize. *P. Natl. Acad. Sci. USA* 94, 6809–6814
 - 23 Tiffin, P. and Gaut, B.S. (2001) Sequence diversity in the tetraploid *Zea perennis* and the closely related diploid *Z. diploperennis*: Insights from four nuclear loci. *Genetics* 158, 401–412
 - 24 Bennetzen, J. *et al.* (2001) Genetic evidence and the origin of maize (Biology, archaeology). *Lat. Am. Antiq.* 12, 84–86
 - 25 Doebley, J. (2001) George Beadle's other hypothesis: One-gene, one-trait. *Genetics* 158, 487–493
 - 26 Doebley, J. (1990) Molecular evidence and the evolution of maize. *Econ. Bot.* 44, 6–27
 - 27 Doebley, J. *et al.* (1987) Patterns of isozyme variation between maize and Mexican annual teosinte. *Econ. Bot.* 41, 234–246
 - 28 van Heerwaarden, J. *et al.* (2011) Genetic signals of origin, spread, and introgression in a large sample of maize landraces. *P. Natl. Acad. Sci. USA* 108, 1088–1092
 - 29 Doebley, J. *et al.* (1997) The evolution of apical dominance in maize. *Nature* 386, 485–488
 - 30 Studer, A. *et al.* (2011) Identification of a functional transposon insertion in the maize domestication gene *tb1*. *Nat. Genet.* 43, 1160–U164
 - 31 Gallavotti, A. *et al.* (2004) The role of *barren stalk1* in the architecture of maize. *Nature* 432, 630–635
 - 32 Sigmon, B. and Vollbrecht, E. (2010) Evidence of selection at the *ramosa1* locus during maize domestication. *Mol. Ecol.* 19, 1296–1311
 - 33 Wang, H. *et al.* (2005) The origin of the naked grains of maize. *Nature* 436, 714–719
 - 34 Briggs, W.H. *et al.* (2007) Linkage mapping of domestication loci in a large maize-teosinte backcross resource. *Genetics* 177, 1915–1928
 - 35 Buckler, E.S. *et al.* (2009) The genetic architecture of maize flowering time. *Science* 325, 714–718
 - 36 Tenaillon, M.I. *et al.* (2004) Selection versus demography: A multilocus investigation of the domestication process in maize. *Mol. Biol. Evol.* 21, 1214–1225
 - 37 Vigouroux, Y. *et al.* (2002) Identifying genes of agronomic importance in maize by screening microsatellites for evidence of selection during domestication. *P. Natl. Acad. Sci. USA* 99, 9650–9655
 - 38 Wright, S.I. *et al.* (2005) The effects of artificial selection of the maize genome. *Science* 308, 1310–1314
 - 39 Yamasaki, M. *et al.* (2005) A large-scale screen for artificial selection in maize identifies candidate agronomic loci for domestication and crop improvement. *Plant Cell* 17, 2859–2872
 - 40 Ross-Ibarra, J. *et al.* (2007) Plant domestication, a unique opportunity to identify the genetic basis of adaptation. *P. Natl. Acad. Sci. USA* 104, 8641–8648

- 41 Tenaillon, M.I. *et al.* (2011) Genome size and transposable element content as determined by high-throughput sequencing in maize and *Zea luxurians*. *Genome Biol. Evol.* 3, 219–229
- 42 Baltazar, B.M. *et al.* (2005) Pollination between maize and teosinte: an important determinant of gene flow in Mexico. *Theor. Appl. Genet.* 110, 519–526
- 43 Ellstrand, N.C. *et al.* (2007) Spontaneous hybridization between maize and teosinte. *J. Hered.* 98, 183–187
- 44 Wilkes, H.G. (1967) Teosinte: The Closest Relative of Maize. The Bussey Institute of Harvard University
- 45 Evans, M.M.S. and Kermicle, J.L. (2001) *Teosinte crossing barrier1*, a locus governing hybridization of teosinte with maize. *Theor. Appl. Genet.* 103, 259–265
- 46 Kermicle, J.L. and Evans, M.M.S. (2010) The *Zea mays* sexual compatibility gene *ga2*: naturally occurring alleles, their distribution, and role in reproductive isolation. *J. Hered.* 101, 737–749
- 47 Kermicle, J.L. *et al.* (2006) The *gametophyte-1* locus and reproductive isolation among *Zea mays* subspecies. *Maydica* 51, 219–225
- 48 Barton, N.H. (2001) The role of hybridization in evolution. *Mol. Ecol.* 10, 551–568
- 49 Wilkes, H.G. (1977) Hybridization of maize and teosinte, in Mexico and Guatemala and improvement of maize. *Econ. Bot.* 31, 254–293
- 50 Doebley, J. (1990) Molecular evidence for gene flow among *Zea* species - genes transformed into maize through genetic-engineering could be transferred to its wild relatives, the teosintes. *Bioscience* 40, 443–448
- 51 Hufford, M.B. *et al.* (2012) The genomic signature of crop-wild introgression in maize. arXiv:1208.3894v2 URL: <http://arxiv.org/abs/1208.3894>
- 52 Lauter, N. *et al.* (2004) The inheritance and evolution of leaf pigmentation and pubescence in teosinte. *Genetics* 167, 1949
- 53 Kawecki, T.J. and Ebert, D. (2004) Conceptual issues in local adaptation. *Ecol. Lett.* 7, 1225–1241
- 54 Fournier-Level, A. *et al.* (2011) A map of local adaptation in *Arabidopsis thaliana*. *Science* 334, 86–89
- 55 Hancock, A.M. *et al.* (2011) Adaptation to climate across the *Arabidopsis thaliana* genome. *Science* 334, 83–86
- 56 Weber, A. *et al.* (2007) Major regulatory genes in maize contribute to standing variation in teosinte (*Zea mays* ssp. *parviglumis*). *Genetics* 177, 2349
- 57 Moeller, D.A. *et al.* (2007) Population structure and its effects on patterns of nucleotide polymorphism in teosinte (*Zea mays* ssp. *parviglumis*). *Genetics* 176, 1799
- 58 Hufford, M.B. *et al.* (2011) Influence of cryptic population structure on observed mating patterns in the wild progenitor of maize (*Zea mays* ssp. *parviglumis*). *Mol. Ecol.* 20, 46–55
- 59 van Heerwaarden, J. *et al.* (2010) Fine scale genetic structure in the wild ancestor of maize (*Zea mays* ssp. *parviglumis*). *Mol. Ecol.* 19, 1162–1173
- 60 Ross-Ibarra, J. *et al.* (2009) Historical divergence and gene flow in the genus *Zea*. *Genetics* 181, 1399
- 61 Rodriguez, F.J.G. *et al.* (2006) Characterization of floral morphology and synchrony among *Zea* species in Mexico. *Maydica* 51, 383–398

- 62 Weber, A.L. *et al.* (2008) The genetic architecture of complex traits in teosinte (*Zea mays* ssp. *parviglumis*): new evidence from association mapping. *Genetics* 180, 1221
- 63 Moeller, D.A. and Tiffin, P. (2008) Geographic variation in adaptation at the molecular level: a case study of plant immunity genes. *Evolution* 62, 3069–3081
- 64 Fang, Z. *et al.* Megabase-scale inversion polymorphism in the wild ancestor of maize. *Genetics* 191, 883–894
- 65 Kirkpatrick, M. and Barton, N. (2006) Chromosome inversions, local adaptation and speciation. *Genetics* 173, 419–434
- 66 Pyhäjärvi, T. *et al.* (2012) Complex patterns of local adaptation in teosinte. arXiv:1208.0634v1 URL: <http://arxiv.org/abs/1208.0634>
- 67 Mano, Y. *et al.* (2006) Variation for root aerenchyma formation in flooded and non-flooded maize and teosinte seedlings. *Plant Soil* 281, 269–279
- 68 Mano, Y. *et al.* (2008) A linkage map of maize x teosinte *Zea luxurians* and identification of QTLs controlling root aerenchyma formation. *Mol. Breeding* 21, 327–337
- 69 Schnable, J.C. *et al.* (2011) Differentiation of the maize subgenomes by genome dominance and both ancient and ongoing gene loss. *P. Natl. Acad. Sci. USA* 108, 4069–4074
- 70 Westerbergh, A. and Doebley, J. (2004) Quantitative trait loci controlling phenotypes related to the perennial versus annual habit in wild relatives of maize. *Theor. Appl. Genet.* 109, 1544–1553
- 71 Churchill, G. *et al.* (2004) The Collaborative Cross, a community resource for the genetic analysis of complex traits. *Nat. Genet.* 36, 1133–1137
- 72 Macdonald, S.J. and Long, A.D. (2007) Joint estimates of quantitative trait locus effect and frequency using synthetic recombinant populations of *Drosophila melanogaster*. *Genetics* 176, 1261–1281
- 73 Tenaillon, M.I. *et al.* (2010) A triptych of the evolution of plant transposable elements. *Trends Plant Sci.* 15, 471–478
- 74 Mano, Y. and Omori, F. Breeding for flooding tolerant maize using "teosinte" as a germplasm resource. *Plant Root* 1, 17–21
- 75 Creighton, H.B. and McClintock, B. (1931) A correlation of cytological and genetical crossing-over in *Zea mays*. *P. Natl. Acad. Sci. USA* 17, 492–497
- 76 Ting, Y.C. (1967) Common inversion in maize and teosinte. *Am. Nat.* 101, 87–89
- 77 Springer, N.M. *et al.* (2009) Maize inbreds exhibit high levels of copy number variation (CNV) and presence/absence variation (PAV) in genome content. *PLoS Genet.* 5, e1000734
- 78 Brunner, S. *et al.* (2005) Evolution of DNA sequence nonhomologies among maize inbreds. *Plant Cell* 17, 343–360
- 79 Lamb, J.C. and Birchler, J.A. (2006) Retroelement genome painting: Cytological visualization of retroelement expansions in the genera *Zea* and *Tripsacum*. *Genetics* 173, 1007–1021
- 80 Morgante, M. *et al.* (2007) Transposable elements and the plant pan-genomes. *Curr. Opin. Plant Biol.* 10, 149–155
- 81 Eichten, S.R. *et al.* (2011) B73-Mo17 near-isogenic lines demonstrate dispersed structural variation in maize. *Plant Physiol.* 156, 1679–1690

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

657 82 Swanson-Wagner, R.A. *et al.* (2010) Pervasive gene content variation and copy
658 number variation in maize and its undomesticated progenitor. *Genome Res.* 20,
659 1689–1699
660 83 Albert, P.S. *et al.* (2010) Diversity of chromosomal karyotypes in maize and its
661 relatives. *Cytogenet. Genome Res.* 129, 6–16
662 84 Laurie, D.A. and Bennett, M.D. (1985) Nuclear-DNA content in the genera *Zea* and
663 *Sorghum* - intergeneric, interspecific and intraspecific variation. *Heredity* 55, 307–
664 313
665 85 Poggio, L. *et al.* (1998) Genome size and environmental correlations in maize (*Zea*
666 *mays* ssp. *mays*, *Poaceae*). *Ann. Bot.-London* 82, 107–115
667 86 Ganai, M.W. *et al.* (2011) A large maize (*Zea mays* L.) SNP genotyping array:
668 development and germplasm genotyping, and genetic mapping to compare with the
669 B73 reference genome. *PLoS One* 6, e28334
670 87 Elshire, R.J. *et al.* (2011) A robust, simple genotyping-by-sequencing (GBS) approach
671 for high diversity species. *PLoS One* 6, e19379
672 88 Shukla, V.K. *et al.* (2009) Precise genome modification in the crop species *Zea mays*
673 using zinc-finger nucleases. *Nature* 459, 437–U156
674 89 Vollbrecht, E. *et al.* (2010) Genome-wide distribution of transposed dissociation
675 elements in maize. *Plant Cell* 22, 1667–1685
676 90 Williams-Carrier, R. *et al.* (2010) Use of Illumina sequencing to identify transposon
677 insertions underlying mutant phenotypes in high-copy Mutator lines of maize. *Plant*
678 *J.* 63, 167–177

Figure Legends:

Figure 1: Ecological and Taxonomic Diversity of *Zea*

The wild *Zea* are distributed across a wide climatic range of temperature (a) and precipitation (b). Within-taxon and between-taxa variation has also been observed in traits such as flowering time (summarized based on data from [61]) (c). The phylogeny of *Zea* (d) based on ~1000 SNPs from across the genome (data are from [64]).

Figure 2: Discovery process from population genomics in teosinte to crop improvement in maize. Natural variation among wild teosinte populations can be used to identify genetic variation that correlates with environmental variables. Further, common garden experiments, maize mutation databases, and tissue specific expression data can be used to identify genes potentially underlying adaptation to environment. Finally, causative variants can be transformed into maize to verify phenotypic effects.

Table 1

Table 1: Attributes of the Teosinte Taxa

Taxa	Genomic			Environmental			Life History	
	Chromosomes	Mean Number of Knobs ^a	Genome Size 1C (pg) ^b	Mean Annual Precipitation (mm) ^c	Mean Annual Temperature (°C) ^c	Average Altitude (m) ^c	Life Cycle	Hybridization with other <i>Zea</i> Taxa ^d
<i>Zea mays</i> ssp. <i>mays</i>	2N=2X=10	4	2.73	1141 ± 802	18.04 ± 5.93	1328 ± 1106	Annual	-
<i>Zea mays</i> ssp. <i>parviglumis</i>	2N=2X=10	8.375	2.94	1141 ± 194	22.84 ± 2.49	1203 ± 387	Annual	Viable hybrids commonly formed with maize supported with genetic evidence; forms limited hybrid zone with <i>mexicana</i>
<i>Zea mays</i> ssp. <i>mexicana</i>	2N=2X=10	8.5	2.58	794 ± 149	16.38 ± 2.16	2179 ± 325	Annual	Viable hybrids commonly formed with maize supported with genetic evidence; forms limited hybrid zone with <i>parviglumis</i> ; some genetic evidence for hybridization with <i>luxurians</i>
<i>Zea mays</i> ssp. <i>huehuetenangensis</i>	2N=2X=10	11.5	N/A	2093 ± 275	21.22 ± 1.06	1181 ± 0.3	Annual	Capable of hybridization with maize
<i>Zea diploperennis</i>	2N=2X=10	7	2.65	1235 ± 76	17.94 ± 0.92	1779 ± 152	Perennial	Documented rare but viable hybridization with maize supported with genetic evidence
<i>Zea perennis</i>	2N=4X=20	N/A	5.28	1009 ± 236	16.82 ± 1.98	1946 ± 329	Perennial	No evidence of successful hybridization
<i>Zea luxurians</i>	2N=2X=10	10	4.58	1629 ± 1121	22.88 ± 1.17	858 ± 327	Annual	Limited evidence of hybridization with maize with some supporting genetic evidence
<i>Zea nicaraguensis</i>	2N=2X=10	N/A	N/A	1746 ± 85	27.91 ± 0.1	24 ± 0.1	Annual	N/A


a) For ssp. *mays*, the number of knobs in B73 is reported from [83]. For all other species, mean knobs were calculated from [44].



b) Genome sizes are reported from the Kew (<http://data.kew.org>) Plant DNA C-values database (*mays*, *diploperennis*, *perennis*, *luxurians*) and [8] (*parviglumis* and *mexicana*).

c) Environmental information was estimated from bioclim (<http://www.worldclim.org/bioclim>) variables of sites where populations are documented.




d) Hybridization reports originate within [2,42-44,49,50]

Figure 1






60 candidate SNPs that associate with an environmental variable



Box 1, Figure 1



1

2

3

4

5

6

7

8

9

10

Z. mays ssp. *parviglumis*