


Copyright for test papers and marking guides remains with *West Australian Test Papers*.
Test papers may only be reproduced within the purchasing school according to the advertised Conditions of Sale.
Test papers should be withdrawn after use and stored securely in the school until Friday 5th July 2019.



SEMESTER TWO

MATHEMATICS METHODS UNIT 1

2019

SOLUTIONS


Calculator-free Solutions

1. (a) Since $x = 1$ then $y = \frac{a+b}{4}$ and $\frac{1}{a} + \frac{y}{2a} = 1$ ✓
 $\therefore \frac{1}{a} + \frac{a+b}{8a} = 1$ ✓
 $\therefore 8 + a + b = 8a \rightarrow b = 7a - 8$ ✓
- (b) $y = \frac{ax}{4} + \frac{b}{4} \rightarrow m_1 = \frac{a}{4}$ ✓
 $\frac{x}{a} + \frac{y}{2a} = 1 \rightarrow y = -2x + 2a \rightarrow m_2 = -2$ ✓
 Since $\perp \quad \frac{a}{4} = \frac{1}{2} \rightarrow a = 2$ ✓
 and $b = 14 - 8 = 6$ ✓ [7]
2. (a) $-\frac{2}{2a} = -2$ ✓
 $\therefore -4a = -2 \rightarrow a = \frac{1}{2}$ ✓
- (b) $f_2(x) = -\frac{1}{2}(x^2 + 4x) + b = -\frac{1}{2}[(x+2)^2 - 4] + b$ ✓
 $= -\frac{1}{2}(x+2)^2 + 2 + b$ ✓
- (c) Translation of c up $f_2(x) + c \rightarrow y = -\frac{1}{2}x^2 - 2x + (b+c)$ ✓
 Since tangential, $b^2 - 4ac = 0$
 $\therefore 4 - 4\left(-\frac{1}{2}\right)(b+c) = 0$ ✓
 $\therefore 4 + 2(b+c) = 0$ ✓
 $\therefore c + b = -2 \rightarrow c = -2 - b$ ✓ [8]
3. (a) $y = a(x+1)(x-2)^2$ ✓
 $\therefore (0, 2) \rightarrow 2 = 1(4)a \rightarrow a = \frac{1}{2}$
 $\therefore y = \frac{1}{2}(x+1)(x-2)^2$ ✓
- (b) $y = a\sqrt{x+4}$ ✓
 $\therefore (0, -4) \rightarrow -4 = a\sqrt{4} \rightarrow a = -2$ ✓
 $\therefore y = -2\sqrt{x+4}$ [4]
4. (a) $D_x = \{x \in R : -3 \leq x \leq 3\}$ ✓
 $R_y = \{y \in R : -3 \leq y \leq 3\}$ ✓
- (b) $D_x = \{x \in R : x \leq 9\}$ ✓
 $R_y = \{y \in R\}$ ✓
- (c) There is a one to many mapping or vertical line test fails ✓ [5]

5. (a) (i) $140^\circ = 140 \times \frac{\pi}{180} = \frac{7\pi}{9}$ ✓
- (ii) $-\frac{360^\circ}{\pi} = -\left(\frac{360}{\pi}\right) \times \frac{\pi}{180} = -2^R$ ✓
- (b) (i) $\frac{2\pi}{5} = \frac{2\pi}{5} \times \frac{180}{\pi} = 72^\circ$ ✓
- (ii) $-1.5^R = -1.5 \times \frac{180}{\pi} = -\frac{270^R}{\pi}$ ✓
- (c) $\sin 2^R \approx \sin 115^\circ > \sin 2^\circ$ ✓
 since it has a higher y value in the Unit circle. ✓ [6]

6. (a) $\tan \frac{11\pi}{4} = \tan \left(2\pi + \frac{3\pi}{4} \right) = \tan \frac{3\pi}{4}$ ✓
 $= -1$ ✓
- (b) $\sin^2 30^\circ - \cos^2 60^\circ = \left(\frac{1}{2} \right)^2 - \left(\frac{1}{2} \right)^2$ ✓
 $= 0$ ✓
- (c) $\frac{\tan^2 \frac{\pi}{3} + \cos \pi}{\sin \frac{\pi}{2} - \tan \frac{3\pi}{4}} = \frac{(\sqrt{3})^2 - 1}{1 - (-1)}$ ✓✓
 $= \frac{2}{2} = 1$ ✓ [7]

7. (a) $a = -1$ ✓
 $\frac{360}{b} = 540$ ✓
 $\therefore b = \frac{2}{3}$ ✓



- (c) $0^\circ, 330^\circ, 660^\circ$ ✓✓✓ [8]

8. (a) $f(x) = -2x(2x^2 - 9x + 4)$ ✓
 $\therefore f(x) = -2x(2x - 1)(x - 4)$ ✓
- (b) (i) $18x^2 - 4x^3 - 8x = 0 \rightarrow -2x(2x - 1)(x - 4) = 0$
 $\therefore x = 0 \text{ or } \frac{1}{2} \text{ or } 4$ ✓
- (ii) $36y^4 - 16y^3 - 8y^5 = 0 \rightarrow 2y^2(18y^2 - 4y^3 - 8y) = 0$
 $\therefore y = 0 \text{ or } \frac{1}{2} \text{ or } 4$ ✓✓ [5]


Calculator-Assumed Solutions

9. (a) $n = 5$ ✓
- (b) $(x + by)^5 \rightarrow T_4 = \binom{5}{2} b^3 x^2 y^3 = 80x^2 y^3$ ✓
 $\therefore 10b^3 = 80 \rightarrow b = 2$ ✓
- (c) $(x + 2y)^5 \rightarrow T_5 = \binom{5}{4} 2^4 xy^4$ ✓
 $\therefore 80xy^4$ ✓ [5]
10. (a) (i) $T = \frac{k}{t^2}$ ✓
 $12 = \frac{k}{4} \rightarrow k = 48$ ✓
- (ii) $T = \frac{48}{t^2} \rightarrow T = \frac{48}{(4.75)^2} = 2.127$ ✓
 \therefore Temperature is 2°C ✓
- (b) D ✓ [5]
11. (a) $\sin 43^\circ = \frac{x}{6} \rightarrow x = 6\sin 43^\circ = 4.09$ ✓
 $\therefore \tan a^\circ = \frac{5}{4.09} \rightarrow a = 50.70^\circ$ ✓
- (b) $r \theta = 24 \rightarrow r = \frac{24}{\theta}$ ✓
 $54 = \frac{1}{2} r^2 (\theta - \sin \theta)$ ✓
 $\therefore 54 = \frac{1}{2} \left(\left(\frac{24}{\theta} \right)^2 \right) (\theta - \sin \theta)$ ✓
 $\therefore \theta = 1.211$ ✓
 $\therefore r = 19.8 \text{ cm}$ ✓ [7]

12. (a) $T = 720^\circ = \frac{360}{n} \rightarrow n = \frac{1}{2}$ ✓
 $\therefore y = 2\sin \frac{x}{2}$ ✓
- (b) $T = 360^\circ \rightarrow n = 1$ ✓
 $\therefore y = \cos(x + c)$ ✓
 $(45^\circ, 1) \rightarrow 1 = \cos(45^\circ + c) \rightarrow c = -45^\circ$ ✓
 $\therefore y = \cos(x - 45^\circ)$
- (c) $T = \frac{\pi}{3} \rightarrow \frac{\pi}{3} = \frac{\pi}{n} \rightarrow n = 3$ ✓
 $\therefore y = -\tan 3x + 1$ ✓✓ [8]
13. (a) $\frac{1}{2}x + 5 = 2x - 4$ ✓
 $\therefore x + 10 = 4x - 8 \rightarrow x = 6$ ✓
 $\therefore A(6, 8)$ ✓
- (b) $\left(\frac{6+x}{2}, \frac{8+y}{2}\right) = (2, 6)$ ✓
 $\therefore B(-2, 4)$ ✓
- (c) \overline{CB} has equation $y = -\frac{1}{2}x + c$ (Angle in Semi-circle Theorem)
 $\therefore (-2, 4) \rightarrow c = 3 \rightarrow y = -\frac{x}{2} + 3$ ✓
Hence $-\frac{x}{2} + 3 = 2x - 4 \rightarrow x = 2.8$ ✓
 $\therefore C(2.8, 1.6)$ ✓ [8]
14. (a) (i) $\sqrt{2}\sin(x + 45^\circ) = \sqrt{2}\sin x \cos 45^\circ + \sqrt{2}\sin 45^\circ \cos x$ ✓
 $= \sin x + \cos x$ ✓
- (ii) $\sqrt{2}\sin 75^\circ = \sqrt{2}\sin(45^\circ + 30^\circ) = \sin 30^\circ + \cos 30^\circ$ ✓
 $= \frac{1}{2} + \frac{\sqrt{3}}{2} \text{ or } \frac{1}{2}(1 + \sqrt{3})$ ✓
- (b) $\frac{\cos\left(\frac{\pi}{2} + \theta\right)}{\sin\left(\frac{\pi}{2} - \theta\right)} = \frac{-\sin \theta}{\cos \theta}$ ✓
 $= -\tan \theta$ ✓ [6]
15. (a) It starts in section B. ✓
In section B, $h(0) = 0$ which is the initial height. ✓
- (b) Section A = Section B when
 $0.01t^2 - 2t + 5 = -0.001t^2 - t$ ✓
 $\therefore t = 5.3$ seconds ✓
- (c) Occurs when Section A has its minimum ✓
 $\therefore h = 95$ m ✓
- (d) Coaster back to height of zero when Section A = 0 ✓
 $0.01t^2 - 2t + 5 = 0$ ✓
 $\therefore t = 197$ seconds ✓ [8]

16. (a) $y = \frac{a}{x+1} + 3$ ✓✓
 $\therefore (0, 1) \rightarrow 1 = \frac{a}{1} + 3$ ✓
 $\therefore a = -2$ ✓

(b)




✓✓ [6]

17. (a) (i) 8.6 m ✓
(ii) $t = 18 \rightarrow 6$ pm ✓
(b) $-2\sin\left(\frac{\pi t}{12}\right) + 10 = 9$ ✓
 $\therefore t = 2$ and $t = 10$ ✓
Hence between 2 am and 10 am. ✓ [5]

18. (a) $b = 6$ ✓
 $f = 10$ ✓
(b) $b + c = f$ ✓
(c) 13 ✓
(d) $\begin{pmatrix} 8 \\ 4 \end{pmatrix}$ ✓
(e) $p = \frac{1}{2}$ ✓✓ [6]


19. (a) (i) $\binom{5}{3}\binom{5}{2} = 10 \times 10 = 100$ ✓✓
- (ii) $\binom{2}{2}\binom{3}{1}\binom{5}{2} + \binom{2}{1}\binom{3}{0}\binom{5}{4} + \binom{2}{2}\binom{3}{0}\binom{5}{3}$ ✓
 $= 30 + 10 + 10 = 50$ ✓
- (b) How many different selections containing four blue pens and a black pen are possible? ✓✓
- (c) (i) $P(\text{at least one}) = 1 - P(0) = 1 - \frac{\binom{3}{0}\binom{7}{5}}{\binom{10}{5}}$ ✓
- $= 1 - \frac{21}{252} = \frac{231}{252}$ ✓
- (ii) $P(\text{at most two red} \mid \text{at least one red})$
 $= \frac{\binom{3}{1}\binom{7}{4} + \binom{3}{2}\binom{7}{3}}{231}$ ✓
 $= \frac{105 + 105}{231} = \frac{210}{231}$ ✓ [10]
20. (a) (i) $\frac{DB}{\sin 69^\circ} = \frac{14}{\sin 23^\circ}$ ✓
 $\therefore DB = 33 \text{ m}$ ✓
- (ii) $20^2 = 15^2 + DB^2 - 2(15)(DB)\cos(\text{ABD})$ ✓
 $\therefore \angle \text{ABD} = 19.84^\circ$ ✓
- (b) $\text{Area}(\triangle ABD) = \frac{1}{2} \times 15 \times DB \times \sin(\text{ABD})$ ✓
 $= 85.16 \text{ m}^2$ ✓
 $\text{Area}(\triangle BCD) = \frac{1}{2} \times 14 \times DB \times \sin 88^\circ$ ✓
 $= 234.01 \text{ m}^2$ ✓
 $\therefore \text{Total area} = 319.17 \text{ m}^2$ ✓ [7]

21. (a) (i)



✓✓

(ii)



✓✓

(b) $E' \cap (D \cup F)$

✓✓

[6]

22. (a) $P(L1) = 0.7$, $P(L2) = 0.35$, $P(L2|L1) = 0.35$ and $P(L3) = 0.4$
 Since $P(L2) = P(L2|L1) = 0.35$
 then independent events.

✓

✓

(b) (i) $P(\text{all three}) = P(L1) \times P(L2) \times P(L3)$
 $= 0.7 \times 0.35 \times 0.4 = 0.098$

✓✓

(ii) $P(L1 \cup L2) = P(L1) + P(L2) - P(L1 \cap L2)$
 $= 0.7 + 0.35 - 0.7 \times 0.35 = 0.805$

✓✓

(iii) $P((L1 \cup L2 \cup L3)' | L1')$
 $= P(L2' \cup L3')$
 $= 0.65 + 0.6 - 0.65 \times 0.6 = 0.86$

✓

✓

[8]

23. (a) (i) $\frac{185}{200}$

✓

(ii) $\frac{85}{125}$

✓✓

(b) $P(Y) = \frac{125}{200}$ and $P(7\&8) = \frac{100}{200}$

$P(Y \cap 7\&8) = \frac{85}{200}$

✓

Since $P(Y) \times P(7\&8) \neq P(Y \cap 7\&8)$ then not independent.

✓

[5]