


Copyright for test papers and marking guides remains with *West Australian Test Papers*.
Test papers may only be reproduced within the purchasing school according to the advertised Conditions of Sale.
Test papers should be withdrawn after use and stored securely in the school until Friday July 5th 2019.



MATHEMATICS SPECIALIST UNIT 1


Semester One

2019

SOLUTIONS

Calculator-free Solutions

1.



[6]

2. (a) (i) $\begin{pmatrix} 3 \\ -5 \end{pmatrix} = k \begin{pmatrix} 4 \\ \alpha \end{pmatrix}$ ✓
 $\therefore k = \frac{3}{4} \rightarrow \alpha = -\frac{5}{k} = -\frac{20}{3}$ ✓

(ii) $\begin{pmatrix} 4 \\ \alpha \end{pmatrix} \cdot \begin{pmatrix} 1 \\ 1 \end{pmatrix} = 0$ ✓
 $\therefore 4 + \alpha = 0 \rightarrow \alpha = -4$ ✓

(iii) $\overrightarrow{PQ} = \begin{pmatrix} 1 \\ \alpha + 5 \end{pmatrix}$
 since the x-coordinate is already 1 unit in length, then
 the y-coordinate must be zero. ✓
 $\therefore \alpha = -5$ ✓

(iv) PQ as base $\Rightarrow |OP| = |OQ|$
 $\left| \begin{pmatrix} 3 \\ -5 \end{pmatrix} \right| = \left| \begin{pmatrix} 4 \\ \alpha \end{pmatrix} \right| \rightarrow \sqrt{34} = \sqrt{16 + \alpha^2}$ ✓
 $\therefore \alpha^2 = 18 \rightarrow \alpha = \pm 3\sqrt{2}$ ✓

2. (b) (i) Solving simultaneously (any method, elimination shown below):

$$\begin{array}{rcl} \mathbf{u} = -6\mathbf{i} - 2\mathbf{j} & \times 3 & \\ \mathbf{v} = 2\mathbf{i} + 3\mathbf{j} & \times 2 & \end{array} \rightarrow \begin{array}{rcl} 3\mathbf{u} = -18\mathbf{i} - 6\mathbf{j} & & \\ 2\mathbf{v} = 4\mathbf{i} + 6\mathbf{j} & & \end{array} \downarrow (+)$$

$$\therefore 3\mathbf{u} + 2\mathbf{v} = -14\mathbf{i} \rightarrow \mathbf{i} = -\frac{3}{14}\mathbf{u} - \frac{1}{7}\mathbf{v} \quad \checkmark\checkmark$$

similarly (or by substitution):

$$\begin{array}{rcl} \mathbf{u} = -6\mathbf{i} - 2\mathbf{j} & \times 1 & \\ \mathbf{v} = 2\mathbf{i} + 3\mathbf{j} & \times 3 & \end{array} \rightarrow \begin{array}{rcl} \mathbf{u} = -6\mathbf{i} - 2\mathbf{j} & & \\ 3\mathbf{v} = 6\mathbf{i} + 9\mathbf{j} & & \end{array} \downarrow (+)$$

$$\therefore \mathbf{u} + 3\mathbf{v} = 7\mathbf{j} \rightarrow \mathbf{j} = \frac{1}{7}\mathbf{u} + \frac{3}{7}\mathbf{v} \quad \checkmark\checkmark$$

$$(ii) \mathbf{r} = 14\left(-\frac{3}{14}\mathbf{u} - \frac{1}{7}\mathbf{v}\right) + 7\left(\frac{1}{7}\mathbf{u} + \frac{3}{7}\mathbf{v}\right)$$

$$\therefore \mathbf{r} = -2\mathbf{u} + \mathbf{v} \quad \checkmark\checkmark \quad [14]$$

3. (a) (i) $20! - 18! = 20 \times 19 \times 18! - 18!$ \checkmark

$$= (20 \times 19 - 1) \times 18!$$

$$= (380 - 1)k = 379k \quad \checkmark$$

$$(ii) \frac{{}^{20}P_3}{{}^{21}C_3} = \frac{20!}{3!} \div \frac{21!}{3! \times 18!} \quad \checkmark$$

$$= \frac{20!}{3!} \times \frac{3! \times 18!}{21 \times 20!} = \frac{k}{21} \quad \checkmark$$

$$(b) \text{ RHS} = \binom{n}{n-r} = \frac{n!}{(n-r)! \times [n-(n-r)]!} \quad \checkmark$$

$$= \frac{n!}{(n-r)! [r]!} \quad \checkmark$$

$$= \binom{n}{r} = \text{LHS} \quad [6]$$

4. (a) If $m < 1$, then $m > m^2$. \checkmark

It is NOT always true because it does not work for negatives. \checkmark

$$\text{e.g. } m = -2 < 1 \rightarrow m^2 = 4 > m \quad \therefore \text{false} \quad \checkmark$$

The converse is always true for $0 < m < 1$ \checkmark

- (b) If the parallelogram is not a rectangle, then it does not have congruent diagonals. \checkmark

Yes it is always true as only squares and rectangles have congruent diagonals. \checkmark

- (c) For all rational numbers \checkmark , there exists two integer numbers a and b \checkmark

such that p is the quotient of a and b . [8]

5. (a) (i) $2^6 = 1 + 6 + 15 + 20 + 15 + 6 + 1 = 64$ ✓
- (ii) $11^5 = (10 + 1)^5$
 $= 10^5 + 5 \times 10^4 + 10 \times 10^3 + 10 \times 10^2 + 5 \times 10 + 1^5$ ✓
 $= 100\,000 + 50\,000 + 10\,000 + 1\,000 + 50 + 1$
 $= 161\,051$ ✓
- (b) (i) $x = 3$ since ${}^6C_3 = 20$ ✓
(ii) $x = 7$ since ${}^7C_5 = 21$ ✓
(iii) $x = 8$ since ${}^8C_2 = {}^8C_6$ ✓
- (c) $(2x - y)^5$
 $= (2x)^5 + 5(2x)^4(-y)^1 + 10(2x)^3(-y)^2 + 10(2x)^2(-y)^3 + 5(2x)^1(-y)^4 + (-y)^5$ ✓
 $= 32x^5 - 80x^4y + 80x^3y^2 - 40x^2y^3 + 10xy^4 - y^5$ ✓✓
- (d) (i) ${}^8C_5 = 56$ ✓
(ii) ${}^2C_2 \times {}^6C_3 = 1 \times 20 = 20$ ✓
(iii) ${}^3C_2 \times {}^5C_3 + {}^3C_3 \times {}^5C_2$ ✓
 $= 3 \times 10 + 1 \times 10 = 30 + 10 = 40$ ✓ [13]
6. (a) $\overrightarrow{AD} = \frac{2}{5}\overrightarrow{AB} = \frac{2}{5}(\mathbf{b} - \mathbf{a})$ ✓
 $\overrightarrow{CD} = \overrightarrow{CO} + \overrightarrow{OA} + \overrightarrow{AD} = -\frac{1}{2}\mathbf{b} + \mathbf{a} + \frac{2}{5}(\mathbf{b} - \mathbf{a})$ ✓
 $\therefore \overrightarrow{CD} = \frac{3}{5}\mathbf{a} - \frac{1}{10}\mathbf{b}$ ✓
- (b) $\overrightarrow{OC} + \overrightarrow{CE} = \overrightarrow{OE} \rightarrow \overrightarrow{OC} + \beta\overrightarrow{CD} = \alpha\overrightarrow{OA}$ given
 $\therefore \frac{1}{2}\mathbf{b} + \beta\left(\frac{3}{5}\mathbf{a} - \frac{1}{10}\mathbf{b}\right) = \alpha\mathbf{a}$ ✓
 $\times 10 \rightarrow 5\mathbf{b} + 6\beta\mathbf{a} - \beta\mathbf{b} = 10\alpha\mathbf{a}$
 $\rightarrow (6\beta - 10\alpha)\mathbf{a} = (\beta - 5)\mathbf{b}$
since \mathbf{a} and \mathbf{b} are non-parallel, then:
 $\beta - 5 = 0 \rightarrow \beta = 5$ ✓
 $6\beta - 10\alpha = 0 \rightarrow \alpha = \frac{3}{5}\beta = 3$ ✓ [6]

Calculator-assumed Solutions

7. (a) ABC collinear
- \Rightarrow
- AB // BC

$$\overrightarrow{AB} = \begin{pmatrix} -2 \\ 4 \end{pmatrix} - \begin{pmatrix} 4 \\ -5 \end{pmatrix} = \begin{pmatrix} -6 \\ 9 \end{pmatrix}$$

$$\overrightarrow{BC} = \begin{pmatrix} -6 \\ 10 \end{pmatrix} - \begin{pmatrix} -2 \\ 4 \end{pmatrix} = \begin{pmatrix} -4 \\ 6 \end{pmatrix}$$

$$\therefore \begin{pmatrix} -6 \\ 9 \end{pmatrix} = k \begin{pmatrix} -4 \\ 6 \end{pmatrix} \rightarrow \begin{matrix} k = \frac{-6}{-4} = \frac{3}{2} \\ k = \frac{9}{6} = \frac{3}{2} \end{matrix} \quad \checkmark\checkmark$$

Since k is unique, then AB // BC and hence ABC collinear. \checkmark

$$(b) |AB| = \left| \begin{pmatrix} -6 \\ 9 \end{pmatrix} \right| = 3 \left| \begin{pmatrix} -2 \\ 3 \end{pmatrix} \right| \text{ and } |BC| = \left| \begin{pmatrix} -4 \\ 6 \end{pmatrix} \right| = 2 \left| \begin{pmatrix} -2 \\ 3 \end{pmatrix} \right| \quad \checkmark$$

$$\therefore AB:BC = 3:2 \quad \checkmark \quad [5]$$

8. (a)
- $\angle PFO = 35^\circ$
- \checkmark

Because $\triangle OFP$ is isosceles since $|OP| = |OF| = \text{radii}$ \checkmark

- (b)
- $\angle FEP = 55^\circ$
- \checkmark

Since $\angle FOP = 110^\circ$ from $\triangle OFP$, and the angle at the centre is double the size of the angle at the edge. \checkmark

- (c)
- $\angle PQF = \angle FEP = 55^\circ$
- \checkmark

Angles at the circumference within the same segment are congruent. \checkmark

- (d)
- $\angle CFP = \angle FEP = 55^\circ$
- \checkmark

The alternate segment theorem \checkmark

- (e)
- $|GC| = |CF| = 11 - |FB| = 11 - 8 = 3 \text{ cm}$
- \checkmark

Tangents to a circle from the same external point are congruent. \checkmark

- (f)
- $|AM| \times (|AM| + 2 \times \text{radius}) = |AH|^2$
- \checkmark

$$|AM| \times (|AM| + 8) = 5^2$$

$$|AM|^2 + 8|AM| - 25 = 0$$

$$\text{CAS} \Rightarrow |AM| = -4 \pm \sqrt{41} \quad \checkmark$$

$$\therefore |AM| = \sqrt{41} - 4 \approx 2.40 \text{ cm only solution} \quad \checkmark \quad [13]$$


9. (a) (i) Divisible by 3 and 5 = divisible by 15
 $100 \div 15 = 6.6 \Rightarrow$ only 6 elements are divisible by 15 ✓
 Therefore, assuming every other element is chosen
 instead of those 6, we need $100 - 6 + 1 = 95$ elements ✓
- (ii) Divisible by 3 = $100 \div 3 = 33.3 \Rightarrow$ 33 elements
 Divisible by 5 = $100 \div 5 = 20$ elements ✓
 Divisible by 3 or 5 = $33 + 20 - 6 = 47$ elements ✓
 Assuming the other 53 elements are chosen first,
 then $53 + 1 = 54$ elements must be chosen ✓
- (b) Assuming the highest numbers are chosen first:
 $100 + 99 + 98 + \dots + 91 + 90 = 955$ ✓
 If 89 is chosen next then the sum exceeds 1000. ✓
 Therefore, a maximum of 11 elements must be chosen. ✓ [8]
10. (a) $n(M \cup C) = n(M) + n(C) - n(M \cap C)$ ✓
 $14\,334 \checkmark = 7\,531 + 9\,885 - n(M \cap C)$
 $\therefore n(M \cap C) = 3\,082$ households ✓
- (b) $n(M \cup C \cup B) = n(M) + n(C) + n(B)$
 $- n(M \cap C) - n(M \cap B) - n(C \cap B)$
 $+ n(M \cap C \cap B)$ ✓
 $\therefore n(M \cup C \cup B) = 7\,531 + 9\,885 + 4\,977 - 3\,082 - 2\,252 - 4\,310 + 1\,724$
 $= 14\,473$ that have all three ✓
 Therefore, $16\,366 - 14\,473 = 1\,893$ households have neither ✓ [6]
11. (a) (i) ${}^{36}P_4 = 1\,413\,720$ OR $({}^{36}C_4 \times 4!)$ ✓
 (ii) ${}^{10}C_2 \times {}^{26}C_2 \times 4! = 351\,000$ ✓✓
 (iii) ${}^5C_1 \times {}^{34}P_2 \times {}^4C_1 = 22\,440$ ✓✓
- (b) II and III ✓✓
- (c) ${}^{x+1}P_3 = {}^4C_3 \times {}^xP_2$
 $\frac{(x+1)!}{(x+1-3)!} = 4 \times \frac{x!}{(x-2)!}$ ✓
 $\frac{(x+1) \times x!}{(x-2)!} = 4 \frac{x!}{(x-2)!}$ ✓
 $(x+1) = 4 \rightarrow x = 3$ ✓

$$\begin{aligned}
 11. \quad (d) \quad \text{LHS} &= \frac{n!}{(n-2)!} + 2n \times \frac{(n-1)!}{(n-1)!} && \checkmark \\
 &= \frac{n!}{(n-2)!} \times \frac{(n-1)}{(n-1)} + \frac{2n!}{(n-1)!} && \checkmark \\
 &= \frac{n! \times (n-1) + 2n!}{(n-1)!} = \frac{n!(n-1+2)}{(n-1)!} && \checkmark \\
 &= \frac{n! \times (n+1)}{(n+1-2)!} = \frac{(n+1)!}{(n+1-2)!} = {}^{n+1}P_2 = \text{RHS} && \checkmark
 \end{aligned}$$

[14]

$$\begin{aligned}
 12. \quad (a) \quad \mathbf{w} &= \begin{pmatrix} -20 \cos 10^\circ \\ -20 \sin 10^\circ \end{pmatrix} \\
 \text{Hovering speed} &= -\mathbf{w} = \begin{pmatrix} 20 \cos 10^\circ \\ 20 \sin 10^\circ \end{pmatrix} && \checkmark\checkmark
 \end{aligned}$$

(b) (i)



$$(ii) \quad \mathbf{w} = \begin{pmatrix} -20 \cos 10^\circ \\ -20 \sin 10^\circ \end{pmatrix} \quad \mathbf{d} = \begin{pmatrix} -25 \cos \theta \\ 25 \sin \theta \end{pmatrix} \quad \mathbf{r} = \begin{pmatrix} -r \cos 24.23^\circ \\ r \sin 24.23^\circ \end{pmatrix} \quad \checkmark\checkmark\checkmark$$

$$\begin{aligned}
 (iii) \quad & \begin{aligned} -r \cos 24.23^\circ &= -20 \cos 10^\circ - 25 \cos \theta \\ r \sin 24.23^\circ &= -20 \sin 10^\circ + 25 \sin \theta \end{aligned} \\
 & \therefore \begin{aligned} 25^2 \cos^2 \theta &= (r \cos 24.23^\circ + 20 \cos 10^\circ)^2 \\ 25^2 \sin^2 \theta &= (r \sin 24.23^\circ + 20 \sin 10^\circ)^2 \end{aligned} \\
 & \rightarrow 25^2 = (r \cos 24.23^\circ + 20 \cos 10^\circ)^2 + (r \sin 24.23^\circ + 20 \sin 10^\circ)^2 \quad \checkmark\checkmark \\
 \text{CAS} \rightarrow & \quad r = 38.8621 \text{ m/s} \quad \text{OR} \quad r = -5.7897 \text{ m/s} \quad \checkmark \\
 & \rightarrow \theta = 46.64^\circ \quad \text{OR} \quad \theta = 2.94^\circ \quad \checkmark \\
 \therefore \text{time} &= \frac{d}{v} = \frac{\sqrt{1200^2 + 540^2}}{38.8621} = 33.86 \text{ seconds} \quad \checkmark \\
 \text{bearing} &= 270^\circ + \theta = 308.86^\circ T \quad \checkmark
 \end{aligned}$$


[14]

13. (a) $\sum F_y = 300 \sin 62^\circ + 252 \sin 56^\circ$ ✓
 $= 473.8N$ ✓
 Since $473.8N < 500N$ the machinery is not moving upwards ✓
- (b) No horizontal component needed $\Rightarrow \sum F_x = 0$
 $\therefore 400 \cos 62^\circ = x \cos 56^\circ$ ✓
 $\rightarrow x = \frac{400 \cos 62^\circ}{\cos 56^\circ} = 335.82N$ ✓
- (c) $\sum F_y = 400 \sin 62^\circ + 335.82 \sin 56^\circ$ ✓
 $= 631.59N$ ✓ [7]
14. (a) (i) Let $n \in \mathbb{N}$ with $n = 2k + 1 = \text{odd}$ ✓
 Then $n^2 + 1 = (2k + 1)^2 + 1$
 $= 4k^2 + 4k + 2$ ✓
 $= 2(2k^2 + 2k + 1)$ ✓
 Since 2 is a factor, then $n^2 + 1$ is divisible by 2, and
 hence the conjecture is true $\forall n \in \mathbb{N}$ ✓
- (ii) Contrapositive statement:
 “if $n^2 + 1$ is odd, then n is even.” ✓
 Let $n^2 + 1 = \text{odd} = 2k + 1$
 $\therefore n^2 = 2k$ ✓
 $\Rightarrow n^2 = \text{even} \Rightarrow n = \text{even}$ ✓
 Since the contrapositive statement is true $\forall n \in \mathbb{N}$,
 then the original conjecture is true $\forall n \in \mathbb{N}$ ✓
- (b) $A \Rightarrow B$:
 If the quadrilateral has two diagonals that intersect at right angles,
 then the quadrilateral is a rhombus, which implies it does have
 two pairs of parallel sides.
 $\therefore A \Rightarrow B$ is true ✓
- $B \Rightarrow A$:
 If the quadrilateral has two pairs of parallel sides then it is a parallelogram,
 which does not necessarily imply it is a rhombus, and therefore it does not necessarily
 have diagonals that intersect at right angles.
 $\therefore B \Rightarrow A$ is false ✓
- Therefore, $A \Leftrightarrow B$ is a false statement. ✓

- (c) Assume that n is odd and n^2 is even. ✓
 Then $\exists k \in \mathbb{N}: n = 2k + 1$
 $\rightarrow n^2 = (2k + 1)^2 = 4k^2 + 4k + 1$ ✓
 $= 2(2k^2 + 2k) + 1 = 2m + 1 = \text{odd}$ ✓
 Since n^2 is both even and odd simultaneously, this is a contradiction ✓
 and therefore the original conjecture must be true $\forall n \in \mathbb{N}, n$ even. [15]

15. (a) (i) $|AB|^2 = |OA|^2 + |OB|^2 - 2|OA||OB|\cos\theta$ ✓
 (ii) $\overrightarrow{AB} \cdot \overrightarrow{AB} = |OA|^2 + |OB|^2 - 2|OA||OB|\cos\theta$
 $\text{LHS} = (\overrightarrow{OB} - \overrightarrow{OA}) \cdot (\overrightarrow{OB} - \overrightarrow{OA})$ ✓
 $= \overrightarrow{OB} \cdot \overrightarrow{OB} - \overrightarrow{OB} \cdot \overrightarrow{OA} - \overrightarrow{OA} \cdot \overrightarrow{OB} + \overrightarrow{OA} \cdot \overrightarrow{OA}$ ✓
 $= |OB|^2 + |OA|^2 - 2\overrightarrow{OA} \cdot \overrightarrow{OB}$ ✓
 $\therefore |OB|^2 + |OA|^2 - 2\overrightarrow{OA} \cdot \overrightarrow{OB} = |OA|^2 + |OB|^2 - 2|OA||OB|\cos\theta$
 $\rightarrow -2\overrightarrow{OA} \cdot \overrightarrow{OB} = -2|OA||OB|\cos\theta$ ✓
 $\rightarrow \overrightarrow{OA} \cdot \overrightarrow{OB} = |OA||OB|\cos\theta$ as required

- (b) (i) $\begin{pmatrix} 4 \\ 5 \end{pmatrix} \cdot \begin{pmatrix} 2 \\ -3 \end{pmatrix} = \begin{vmatrix} 4 \\ 5 \end{vmatrix} \times \begin{vmatrix} 2 \\ -3 \end{vmatrix} \cos\theta$
 $8 - 15 = \sqrt{41} \times \sqrt{13} \cos\theta$ ✓
 $\therefore \cos\theta = -\frac{7}{\sqrt{533}}$ ✓
 Since $\cos\theta < 0 \Rightarrow \theta$ is obtuse ✓

- (ii)  $\Rightarrow \sin\theta = \frac{22}{\sqrt{533}}$ ✓

$$\therefore \text{area } \Delta OAB = \frac{1}{2}|OA||OB|\sin\theta$$

$$= \frac{1}{2}\sqrt{41} \times \sqrt{13} \times \frac{22}{\sqrt{533}} = 11 \text{ units}^2 \quad \checkmark \quad [10]$$

16. P, Q, R and S are the midpoints of their respective sides:

$$\overrightarrow{OP} = \begin{pmatrix} -0.5 \\ 4 \end{pmatrix} \quad \overrightarrow{OQ} = \begin{pmatrix} 6 \\ 1.5 \end{pmatrix} \quad \overrightarrow{OR} = \begin{pmatrix} 1.5 \\ -4 \end{pmatrix} \quad \overrightarrow{OS} = \begin{pmatrix} -5 \\ -1.5 \end{pmatrix} \quad \checkmark$$

Therefore:

$$\overrightarrow{PQ} = \begin{pmatrix} 6 \\ 1.5 \end{pmatrix} - \begin{pmatrix} -0.5 \\ 4 \end{pmatrix} = \begin{pmatrix} 6.5 \\ -2.5 \end{pmatrix}$$

$$\overrightarrow{SR} = \begin{pmatrix} 1.5 \\ -4 \end{pmatrix} - \begin{pmatrix} -5 \\ -1.5 \end{pmatrix} = \begin{pmatrix} 6.5 \\ -2.5 \end{pmatrix} \quad \checkmark$$

$$\overrightarrow{PQ} = \overrightarrow{SR} \quad \Rightarrow \therefore \overrightarrow{PQ} \parallel \overrightarrow{SR} \quad \checkmark$$

$$\overrightarrow{PS} = \begin{pmatrix} -5 \\ -1.5 \end{pmatrix} - \begin{pmatrix} -0.5 \\ 4 \end{pmatrix} = \begin{pmatrix} -4.5 \\ -5.5 \end{pmatrix}$$

$$\overrightarrow{QR} = \begin{pmatrix} 1.5 \\ -4 \end{pmatrix} - \begin{pmatrix} 6 \\ 1.5 \end{pmatrix} = \begin{pmatrix} -4.5 \\ -5.5 \end{pmatrix} \quad \checkmark$$

$$\overrightarrow{PS} = \overrightarrow{QR} \quad \Rightarrow \therefore \overrightarrow{PS} \parallel \overrightarrow{QR} \quad \checkmark$$

Since $\overrightarrow{PQ} \parallel \overrightarrow{SR}$ and $\overrightarrow{PS} \parallel \overrightarrow{QR} \Rightarrow PQRS$ is a parallelogram [5]