




A




B



C



Supplemental Figure 1. Comparison Of Global Editing Rate and ADAR1 Isoform Expression Between HT and WT Mice. (A) Comparison of total number of edited sites and site-specific editing rate between HT and WT mice treated with fluoxetine or vehicle and (B) expression of Adar1p110 and Adar1p150. The p-values were obtained using R package clusterProfiler's default Dunn test for non-parametric comparison. As shown in Figure (A), no pairwise comparison was statistically significant; however, an overall increase in editing in HT mice treated with vehicle was observed. While the highest count values for Adar1p150 and Adar1p110 were observed in HT and WT mice treated with fluoxetine, respectively, no significant difference was found in pairwise comparison(B).



Supplemental Figure 2. Significant Gene Ontology (GO) Terms for Genes with Uniquely Edited Sites. Plot shows significant GO terms (p -values ≤ 0.05) for genes with unique edited sites in (A) HT mice treated with fluoxetine, (B) HT mice treated with vehicle (C) WT mice treated with fluoxetine (D) WT mice treated with vehicle. Terms BP, CC and MF indicate Biological Process, Cellular Component (CC), and Molecular Function (MF). Considering the GO landscape of WT-veh mice (D) as normal, the presence of distinctly different GO terms in all other groups highlights the influence of fluoxetine treatment and the autistic phenotype on ADAR editing.