

As needed medications that have been administered in past 7 days:

- **Acetaminophen** 650 mg (20.5 mL) every 6 hours as needed for pain
 - o Receiving ~1 dose per day at various times
- **Celecoxib** 100 mg 2 times daily as needed for pain
 - o has been receiving 1 dose every night at bedtime
- **Hydroxyzine** 25 mg (12.5 mL) every 6 hours as needed for anxiety or itching
 - o Received 4 total doses in past week at various times
- **Melatonin** 6 mg (2 tablets) as needed for sleep
 - o Has been receiving 6 mg every night at 9 pm
- **Midodrine** 5 mg two times daily as needed for low blood pressure (systolic blood pressure < 100 or before physical therapy)
 - o Received 2 total doses in past week
- **Ondansetron** 4 mg every 6 hours as needed for nausea/vomiting
 - o Received 1 total dose in past week
- **Oxycodone** 5 mg (5 mL) as needed for pain
 - o Has received 1-3 doses/per day at various times over the past week
- **Quetiapine** 25 mg at bedtime as needed for sleep or agitation
 - o Has been receiving 1 dose every night at bedtime

As needed medications that have NOT been administered in past 7 days:

- Bisacodyl suppository daily as needed for constipation
- Refresh Plus eye drops 4 times daily as needed for dry eyes
- Genteal Tears Night-Time ointment as needed for eye dryness or irritation
- Lidocaine 4% cream – apply topically as needed for pain
- Lidocaine 4% patch – apply 1 patch daily as needed for pain (on for 12 hours and off for 12 hours)
- Lidocaine jelly 2% (Glydo) – Place 11 mL into the urethra as needed
- Polyethylene glycol (Miralax) 17 g daily as needed for constipation

Reminders for giving medications through your feeding tube:

- Always give each medication alone. Do not mix medications together.
- Flush feeding tube before, between, and after each dose with 15-20 mL of water.
- Dilute liquid medications with 30 mL prior to administering.
- Crush tablets to a powder and mix in 30 mL of water. May use warm water, never hot water, and leave to dissolve for a couple of minutes.
- Open capsules to mix internal powder in 30 mL of water before administering.
- Not all tablets and capsules can be crushed or mixed in water to give by feeding tube. Do not crush tablets that are extended-release, controlled-release, sustained-release, slow-release, or hazardous drugs.
- Call the pharmacy if you have any questions.