
```

syms x y
f=3*y^2-2*y^3-3*x^2+6*x*y;
p=diff(f,x); q=diff(f,y);
[ax,ay]=solve(p,q);
ax=double(ax);
ay=double(ay);
r=diff(p,x);
s=diff(p,y);
t=diff(q,y);
D=r*t-s^2;
for i=1:size(ax)
    T1=subs(D,[x y],[ax(i),ay(i)]);
    T2=subs(r,[x y],[ax(i), ay(i)]);
    T3=subs(f,[x y],[ax(i), ay(i)]);
    if (double(T1)<0)
        sprintf('The point (%f,%f) is a saddle point',ax(i),ay(i))

    else
        if (double(T2)<0)
            sprintf('The maximum value at (x, y) is (%f, %f)',ax(i),ay(i))
            sprintf('and the maximum value of f(x) is %f',T3)

        else
            sprintf('The minimum value at (x, y) is (%f, %f)',ax(i),ay(i))
            sprintf('and the minimum value of f(x) is %f',T3)

        end
    end
end

ans =

    'The point (0.000000,0.000000) is a saddle point'

ans =

    'The maximum value at (x, y) is (2.000000, 2.000000)'

ans =

    'and the maximum value of f(x) is 8.000000'

```

Published with MATLAB® R2023b