

HUMAN IMPACTS

by the numbers

Griffin Chure¹, Rachel Banks², Avi Flamholz², Nicholas S. Sarai³,
Mason Kamb⁴, Tine Valencic¹, Yinon Bar-On⁵, Ron Milo⁵, Rob Phillips^{2,6,*}

California Institute of Technology, Pasadena, CA, USA, 91125:

¹Department of Applied Physics; ²Division of Biology and Biological Engineering; ³Division of Chemistry and Chemical Engineering;

⁴ Department of Physics;


⁵ Chan-Zuckerberg BioHub, San Francisco, CA, 94158

⁶ Weizmann Institute of Science, Rehovot 7610001, Israel: Department of Plant and Environmental Sciences

ABSTRACT

The greatest experiment of the last 10,000 years is the presence and action of modern human beings on planet Earth. At this point, the consequences of this experiment are being felt on many fronts. Yet, many people still hold the view that because the world is so "huge", humans cannot really make a substantial impact. This snapshot represents a collection of essential numbers that summarize the broad reach of human action across the planet, presenting a view of the impact of human presence on Earth.

REFERENCE UNITS	HUMAN IMPACT CATEGORIES
per capita (global) = 1 person	LAND
human population = ≈ 7 × 10 ⁹	ENERGY
area of soccer pitch = ≈ 3000 m ²	FLORA & FAUNA
volume of olympic pool = ≈ 2000 m ³	WATER
power of an incandescent light bulb = ≈ 100 W	ATMOSPHERIC & BIOGEOCHEMICAL CYCLES
mass of a pick-up truck = ≈ 2000 kg	



SIZING UP THE ANTHROPOCENE

Much of our understanding of the scales of things is comparative. When we measure lengths, we do so relative to some standard length that helps us develop intuition for whatever situation we might be thinking about. With the many examples of human impacts presented on Page 1, our aim here is to present some measure of each of those numbers in a ratiometric form that compares the magnitude of a given human impact to some natural scale for that same quantity. For example, in considering the use of land by humans, a natural dimensionless way to characterize that number is by comparing it to the total land area available on our planet, a comparison that yields what we will christen the Terra number. Similarly, when we consider the entirety of human-made materials, a dimensionless version of that mass that provides a sense of perspective is to compare it to the total biomass on our planet. Here we present twelve key human impacts in this kind of dimensionless form that range across the earth, the atmosphere, the oceans and our energy use, with the hope that our readers will be emboldened to consider their own favorite examples in a similar dimensionless format.

THE TERRA NUMBER

$$Te = \frac{\text{land area used by humans}}{\text{total land area of Earth}} = \frac{\text{area with buildings and roads}}{\text{entire globe}} \approx 0.3$$

The Terra Number reflects the fact that, while we have been constrained to the 30% of Earth's surface that is land, we have transformed the terrestrial surface to support our dwellings and, much more importantly, our agriculture. Despite being icons of human action, urban centers occupy between 750 thousand and a few million km² (Huld: 39341, 87575). In contrast, approximately 50 million km² (Huld: 29582) of land on Earth is used either to grow crops or rear livestock. Together, urban and agricultural lands make up ≈ 30% of Earth's terrestrial surface. [Read more about this number.](#)

THE ANTHROPOMASS NUMBER

$$An = \frac{\text{total anthropomass}}{\text{total biomass}} = \frac{\text{humans + concrete + asphalt}}{\text{trees + animals + soil}} \approx 1$$

The Anthropomass Number takes stock of our material production by comparing the total quantity of human-made materials to the entirety of the biomass on planet Earth. Surprisingly, already in 2007, human-made materials added up to more mass than that of the entire biosphere itself (Huld: XXXXX to be added). Concretes and aggregates dominate the anthropomass, with bricks and asphalt coming in a distant second. Though around us everywhere, plastics and metal constitute a tiny fraction of the total anthropomass. The total anthropomass budget amounts to a dizzying 10⁵ kg of human made mass per person on planet earth, with a volume per person that corresponds to roughly the entirety of one lane of an Olympic-sized swimming pool.

THE EXTINCTION NUMBER

$$Ex = \frac{\text{number of known animal extinctions}}{\text{number of expected natural animal extinctions}} = \frac{\text{extinct bird}}{\text{extinct bird + still extant bird}} > 10$$

The Extinction Number reflects the fact that over the past 500 years, far more animal species have gone extinct than would be expected due to natural processes. Since 1500, more than 760 species of animals have been determined to be extinct (Huld: 44641). Recent estimates of ancient rates of animal extinction predict that tenfold fewer (~50) species would have gone extinct over the same period in the absence of human alteration of the environment. It's important to emphasize that these data are incomplete and reflect only a fraction of nature that has been assessed for conservation status. The Extinction Number therefore represents a lower bound on the degree of modern species loss, with the true value likely being much higher.

THE BARNYARD NUMBER

$$By = \frac{\text{mass of terrestrial livestock}}{\text{mass of terrestrial wild animals}} = \frac{\text{cows, chickens, pigs}}{\text{elephants, foxes, pelicans}} \approx 30$$

The Barnyard Number focuses another lens onto this massive transformation of the planet by comparing the total biomass of terrestrial livestock (e.g. cows, chickens, and pigs) to that of wild terrestrial wild animals (e.g. elephants, foxes, and pelicans)². As a result of agricultural intensification throughout the 20th century, livestock now outweigh all wild terrestrial animals by a factor of ≈ 30. While poultry make up the vast majority of terrestrial livestock (≈25 billion individuals, Huld: 94934), they represent a small proportion of the total livestock biomass. This biomass is dominated by cattle which collectively weigh in at ≈ 1.5×10¹² kg, despite having a population of only ≈ 1.5 billion (Huld: 92006). [Read more about this number.](#)

THE DEFORESTATION NUMBER

$$Df = \frac{\text{annual forest loss from human action}}{\text{annual forest loss from wildfire}} \approx \frac{\text{logs}}{\text{burned forest}} \approx 3$$

The Deforestation Number reflects that through direct action, humans deforest and disrupt forested land at three times the rate of natural forest loss. The bulk of this forest loss is due to commodity-driven deforestation (such as logging, Huld: 96098) and forestry (Huld: 38352), whereas expansion of urban areas accounts for < 1% of the total annual forest loss (Huld: 19429). Wildfires account for ≈ 20% or ≈ 5×10¹⁰ m² annually (Huld: XXXXX). [This paragraph will be rewritten after the vignette is finalized.](#)

THE NIAGARA NUMBER

$$Ni = \frac{\text{daily water volume used by humans}}{\text{Niagara Falls daily discharge volume}} \approx \frac{\text{truck with water tank}}{\text{Niagara Falls waterfall}} \approx 30$$

The Niagara Number captures the magnitude of human water usage relative to the scale of Niagara Falls, the largest waterfall in North America by discharge (volumetric flow rate). Agriculture once again defines this aspect of the human interaction with the Earth system as it uses ≈ 4×10⁹ m³ (Huld: 43593) of water per day, accounting for the majority of human water usage. Combining agricultural use with the water volume used for hydroelectric power generation (≈ 7×10⁸ m³/day, Huld: 77411), domestic/municipal use (≈ 10⁸ m³/day, Huld: 69424), and all other uses yields a total daily volume of water ≈ 30 times that which flows over Niagara Falls daily. This is a volume comparable to ½ the daily discharge of the Amazon river.

THE RIVER NUMBER

$$Rv = \frac{\text{river volume controlled by humans}}{\text{free-flowing river volume}} \approx \frac{\text{dammed river}}{\text{natural river}} \approx 1$$

Hydroelectric power generation captures the potential energy stored in water found at elevation, for example snowmelt cascading from alpine glaciers towards the ocean. Capturing this energy, however, requires damming the river – thus interrupting its flow and altering the riverine ecosystem. The River Number quantifies the scale of human interference in river flow by relating the volume of river systems under human control (primarily due to damming) to that of free-flowing rivers. Globally, approximately an equal volume of river water is free flowing (≈ 6×10¹¹ m³, Huld: 55718) as is under direct human control, such as through dams and reservoirs or through man-made channels (≈ 6×10¹¹ m³, Huld: 61661). Of the global free-flowing river volume, approximately 50% of the volume is contained within the Amazon river alone.

THE EARTH MOVER NUMBER

$$Em = \frac{\text{annual mass of earth moved by humans}}{\text{annual mass of earth moved by rivers}} \approx \frac{\text{bulldozer moving dirt}}{\text{rivers}} > 15$$

Humans are becoming formidable rivals to the natural processes by which sediment is generated and moved. This is illustrated by The Earth Mover Number, which reveals that humans move approximately 15 times more earth than is moved by global river systems. Through construction, mining, and agriculture, humans move more than 250 billion tonnes of sediment a year (Huld: XXXXX). Rivers, by comparison, transport ≈ 12 billion tonnes a year when corrected for the increased river sediment load via human action³. This remarkable anthropogenic action rapidly increases erosion rates, leading to increased topsoil loss and turnover, ultimately perturbing natural biogeochemical cycles.

THE NITROGEN NUMBER

$$N_2 = \frac{\text{mass of } N_2 \text{ fixed via the Haber-Bosch Process}}{\text{mass of } N_2 \text{ fixed biologically}} \approx \frac{\text{NH}_3 \text{ tank}}{\text{NH}_3 \text{ plant}} \approx 1$$

Though molecular nitrogen (N₂) makes up nearly 80% of our atmosphere, it must be converted into a more reactive form like ammonia (NH₃) in order for most plants to synthesize biomass. Deemed the “detonator of the population explosion,” the development of the Haber–Bosch process for industrial synthesis of ammonia from molecular nitrogen was critical for supporting a global population above ≈ 3 billion⁴. The Nitrogen Number reveals that humans synthesize as much reactive nitrogen industrially (≈ 1.5×10¹¹ kg per year, Huld: 91803) as is synthesized by nitrogen-fixing microbes in terrestrial ecosystems (≈ 1×10¹¹ kg per year, Huld: 15205). Beyond influencing the balance of reactive nitrogen in the environment, the Haber–Bosch process uses a sizable amount of energy and significantly contributes to global CO₂ emissions.

THE CO₂ NUMBER

$$CO_2 = \frac{\text{annual mass of anthropogenic CO}_2}{\text{annual mass of volcanic CO}_2} \approx \frac{\text{factory smokestack}}{\text{volcanic eruption}} \approx 100$$

The CO₂ Number compares annual amount of human-caused CO₂ emissions to the dominant natural source of CO₂ emissions, volcanism. There are many climate-related consequences of increasing CO₂ emissions. Beyond accelerating climate change, ≈30% of CO₂ released into the atmosphere is absorbed by the oceans, making them appreciably more acidic over time. In recent years, human activities including burning fossil fuels and making concrete have led to the release of ≈ 4×10¹³ kg of CO₂ into the atmosphere each year. Whether through seeps or eruptions, volcanism releases a total of ≈ 10¹¹ kg of CO₂ per year. Thus, the CO₂ number is ≈100, representing the magnitude by which humans outpace nature as a source of CO₂ emissions.

THE CH₄ NUMBER

$$Me = \frac{\text{annual mass of anthropogenic CH}_4}{\text{annual mass of natural CH}_4} \approx \frac{\text{cow and factory}}{\text{wetland}} \approx 1$$

While CO₂ is the most often discussed greenhouse gas, human activities also entail substantial emissions of methane (CH₄), which even more potent greenhouse gas than CO₂. The CH₄ Number compares humanity's methane emissions to all natural sources of methane. Anthropogenic emissions from factories and ruminant livestock (mostly cows) total ≈ 4 billion kg per year, comparable to natural emissions of CH₄ from wetlands, though there is substantial uncertainty regarding the magnitude of natural methane emissions. Thus, human methane emissions roughly equal emissions from all natural sources, indicating that we are approaching nature in our impact on atmospheric methane chemistry.

THE SOLAR NUMBER

$$Su = \frac{\text{annual human power usage}}{\text{annual incident solar power}} \approx \frac{\text{building}}{\text{sun}} \approx 0.0001$$

While humans derive biological energy from food, we derive mechanical and electrical power from various fuel sources like coal, oil, natural gas, and fissile nuclear material. Current global human power usage amounts to an enormous 20 terawatts (20×10¹² W), a number which has been increasing steadily over the last 200 years. The Solar Number puts the 20 TW power consumption of human activities (Huld: 94934) in relief by comparing it to the power incident on our planet from the sun, which represents a power source roughly 10,000 times greater than our current demands. Of course, it is unlikely that we could ever harness 100% of solar energy incident on our planet, but capturing even 1% of this energy would be sufficient to produce 100 times more power than we currently use.

Human Impacts by the Numbers — Supporting Information

About: Here, we present citations and notes corresponding to each quantity assessed here. Each value presented on page 1 is assigned a Human Impacts Database identifier (HuID), accessible via <https://human-impacts.herokuapp.com>. When possible, primary data sources have been collated and stored as files in comma-separated-value (csv) format on the GitHub repository associated with this snapshot, accessible via DOI: XXXXXXX and https://github.com/rpgroup-pboc/human_impacts. A subset of the numbers presented here are investigated in more depth and presented as a collection of vignettes accessible via https://rpgroup.caltech.edu/hi_vignettes (URL subject to change).

A	ANNUAL ICE MELT	HUMAN POPULATION	POWER FROM RENEWABLES
glaciers $\approx 3 \times 10^{11} \text{ m}^3 / \text{yr}$	HuID: 32459	fraction of population that is urban $\approx 55\%$	wind $\approx 0.2 \text{ TW}$
Data Source(s): Intergovernmental Panel on Climate Change (IPCC) 2019 Special Report "The Ocean and Cryosphere in a Changing Climate," Table 2.A.1 on pp. 199–202. Notes: Value corresponds to the trend of annual mass loss from major glaciated regions (2006–2015). Volume loss was calculated from mass loss.		total population $\approx 7.6 \times 10^9$ people	solar $\approx 0.1 \text{ TW}$
ice sheets $\approx 4 \times 10^{11} \text{ m}^3 / \text{yr}$	HuID: 95798; 93137	Data Source(s): Food and Agricultural Organization (FAO) of the United Nations Report on Annual Population, 2019. Notes: Value for total population comes from a combination of direct population reports from country governments as well as inferred measurements from under-reporting or missing data. Fraction of urban population is determined from what each country reports as "urban" which does not follow a singular definition. As explained by the United Nations population division, "When the definition used in the latest census was not the same as in previous censuses, the data were adjusted whenever possible so as to maintain consistency." Rural population is computed from this fraction along with the total human population, implying that the total population is composed only of "urban" and "rural" communities.	hydroelectric $\approx 0.5 \text{ TW}$
arctic sea ice $\approx 3 \times 10^{11} \text{ m}^3 / \text{yr}$	HuID: 89520	Data Source(s): D. N. Wiese et al. 2019 https://doi.org/10.5067/ITEM-SC-3MJ62 Notes: Value corresponds to the trends of combined annual mass loss from the Greenland and Antarctic Ice Sheets (2002–2020). Volume loss was calculated from mass loss.	total $\approx 0.8 \text{ TW}$
total $\approx 1 \times 10^{12} \text{ m}^3 / \text{yr}$	HuID: 89075	Data Source(s): bp Statistical Review of World Energy, 2020. Notes: Sum of glacial, ice sheet, and sea ice melt rate. Notes: Antarctic sea ice loss is not included due to data sparsity. The periods of analysis are not the same, therefore this rate represents an approximation rather than an exact calculation.	Data Source(s): bp Statistical Review of World Energy, 2020. Notes: Reported values correspond to estimates for the 2019 calendar year. Renewable resources are defined as wind, geothermal, solar, biomass and waste. Hydroelectric, while presented here, is not defined as a renewable in the source material.
B	ANNUAL SEA ICE EXTENT LOSS	GREENHOUSE GAS EMISSIONS	FOSSIL FUEL EXTRACTION
arctic sea ice extent loss $\approx 5 \times 10^{10} \text{ m}^2 / \text{yr}$	HuID: 66277	anthropogenic CO ₂ $\approx 4.2 \times 10^{11} \text{ kg / yr}$	natural gas volume $\approx 4 \times 10^{12} \text{ m}^3$
Data Source(s): Fetterer, F., K. Knowles, W. N. Meier, M. Savoie, and A. K. Windnagel. 2017, updated daily. Sea Ice Index, Version 3. [Sea_Ice_Index_Monthly_Data_with_Statistics_G02135_v3.xls]. Boulder, Colorado USA. NSIDC: National Snow and Ice Data Center. doi: https://doi.org/10.7265/NSK072F8 . [Accessed 2020-Oct-19]. Notes: Value corresponds to the average annual loss of Arctic sea ice extent from 1979–2019. Sea ice extent refers to the area of the sea with > 15% ice coverage. The Antarctic sea ice extent trend is not shown because a significant long-term trend over the satellite observation period is not observed and short-term trends are not yet attributable.		Data Source(s): Friedlingstein et al. 2019, DOI: 10.5194/essd-11-1783-2019. Original data sources relevant to this study compiled in Friedlingstein et al.: 1) Gilfillan et al. https://energy.apgstate.edu/CDIAC 2) Average of two bookkeeping models: Houghton and Nassikas 2017 DOI: 10.1020/2016GB005546; Hansis et al. 2015 DOI: 10.5248/zenodo.1000000. Notes: Value corresponds to CO ₂ emissions from fossil fuel combustion, industrial emissions (predominantly cement production), and land-use change during calendar year 2018. CO ₂ was added to the atmosphere at a rate of $\approx 18.6 \text{ Gt / yr}$ in 2018 (HuID: 98043); most of the remainder is taken up by the land sink and ocean sink.	oil volume $\approx 5.5 \times 10^9 \text{ m}^3$
anthropogenic CH ₄ $\approx 4 \times 10^{11} \text{ kg / yr}$	HuID: 96837; 56405; 30725	Data Source(s): Table 2 of Saunois, et al. 2020. DOI: 10.5194/essd-12-1561-2020. Notes: Value corresponds to CH ₄ emissions from anthropogenic sources in the calendar year 2017. Represents emissions from agriculture and waste, fossil fuels, and biomass and biofuel burning. Value is not simply the sum of these sources but is based on a full anthropogenic inventory of emissions. Natural emissions amount to $\approx 0.3 \text{ Gt / yr}$ in 2017. CH ₄ was added to the atmosphere at a rate of $\approx 17 \text{ Mt / yr}$ in 2017; most of the remainder is taken up by chemical loss sink and soil sink.	coal mass $\approx 8 \times 10^9 \text{ t}$
anthropogenic N ₂ O $\approx 7 \times 10^9 \text{ kg / yr}$	HuID: 44575; 68619	Data Source(s): Table 1 of Tian, H., et al. (2020). DOI: 10.1038/s41586-020-2780-z Notes: Value corresponds to annualized N ₂ O emissions from anthropogenic sources in the years 2007–2016.	Data Source(s): bp Statistical Review of World Energy, 2020. Notes: Oil volume includes crude oil, shale oil, oil sands, condensates, and natural gas liquids separate from specific natural gas mining. Natural gas value excludes gas flared or recycled and includes natural gas produced for gas-to-liquids transformation. Coal value includes 2019 value exclusively for solid commercial fuels such as bituminous coal and anthracite, lignite and sub-bituminous coal, and other solid fuels.
C	ANNUAL MATERIAL PRODUCTION	OCEAN WARMING	POWER FROM NUCLEAR FISSION
concrete production $\approx 3 \times 10^{13} \text{ kg / yr}$	HuID: 25488	power deposition $\approx 160 \text{ TW}$	nuclear power $\approx 0.8 \text{ TW}$
steel production $\approx 2 \times 10^{12} \text{ kg / yr}$	HuID: 51453	ocean surface warming $\approx 0.03^\circ \text{C}$	HuID: XXXX
Data Source(s): USGS 2020, Mineral Commodities. DOI:10.3133/mcs2020; Monteiro et al. 2017, DOI:10.138/nmat4930 Notes: Concrete production value corresponds to approximate value from multiple sources. USGS 2020 Mineral Commodity Survey reports mass of cement produced in 2019. This is converted to concrete using a multiplicative conversion factor of ≈ 7 as described in Monteiro et al. 2017. Steel production corresponds to the USGS 2019 value.		Data Source(s): Intergovernmental Panel on Climate Change (IPCC) 2019 Special Report "The Ocean and Cryosphere in a Changing Climate." Table 5.1 on pp. 458 and 459 and 4 on pp. 457. Notes: Value is calculated from the reported annual heat uptake of $\approx 5 \text{ ZJ / yr}$ over the time period of 2005 – 2017. This assumes a constant value for deposition into the ocean surface (0 – 700 m depth) and deep ocean (700 – 2000 m depth) where heat deposition is lower. Ocean surface temperature change is calculated from $\approx 5 \text{ ZJ / yr}$ heat uptake by noting that deposition of $\approx 144 \text{ ZJ / yr}$ raises the temperature of the top 100 m of ocean by $\approx 1^\circ \text{C}$. See the complete report or section 5.2.2 of the source material for more information.	Data Source(s): bp Statistical Review of World Energy, 2020. Notes: Reported values correspond to estimates for the 2019 calendar year.
plastic production $\approx 4 \times 10^{11} \text{ kg / yr}$	HuID: 97241	NUCLEAR FALLOUT	POWER FROM NUCLEAR FISSION
Data Source(s): Table S1 of Geyer et al. 2017. DOI:10.1126/sciadv.1700782. Notes: Value represents the sum total global production of plastic fibers and plastic resin during calendar year 2015.		239+240Pu activity in soil $\approx 1 \text{ Bq / kg}$	nuclear power $\approx 0.8 \text{ TW}$
D	LIVESTOCK POPULATION	EARTH MOVING	HuID: XXXX
chicken $\approx 2 \times 10^{10}$	HuID: 94934	coal mining waste/overburden $\approx 6.5 \times 10^{13} \text{ kg / yr}$	HuID: 59201
cattle $\approx 2 \times 10^9$	HuID: 92006	urbanization $> 1.4 \times 10^{14} \text{ kg / yr}$	HuID: 87228
swine $\approx 1 \times 10^9$	HuID: 21368	Data Source(s): Figure 4 and Figure 5 in Hancock et al. 2014, DOI:10.1144/SP395.15. Figure 3 (col. 2, rows 3 – 5) of Ciszewski and Łokas, 2019, DOI:10.1515/geochr-2015-0111. Notes: Value corresponds to current-day detectable combined radioactivity of ²³⁹ Pu and ²⁴⁰ Pu present in cores of stratified soil with estimated date of $\approx 1963 \text{ CE}$ during the peak of atmospheric nuclear weapons testing. Reported is approximate average activity from sediment samples in SE Australia (Hancock et al.) and Polish river basins (Ciszewski and Łokas).	Data Source(s): The IUCN Red List of Threatened Species. Version 2020–2. Notes: Values correspond to absolute lower-bound measurements of extinctions caused over the past ≈ 520 years. Of the predicted ≈ 8 million animal species, the IUCN databases catalogues only $\approx 900,000$ with only $\approx 75,000$ being assigned a conservation status. Representation of plants and fungi is even more sparse with only $\approx 40,000$ and ≈ 285 being assigned a conservation status, respectively. The number of extinct animal species is undoubtedly higher than these reported values, as signified by an inequality symbol (>).
total $\approx 3 \times 10^{10}$	HuID: 15765	CONTEMPORARY EXTINCTION	HuID: 44641
Data Source(s): Food and Agriculture Organization of the United Nations Statistical Database (FAOSTAT). Notes: Counts correspond to the approximate average of the standing populations reported between 2010 – 2018. Values are reported directly by countries, yet the FAO uses non-governmental statistical sources to address uncertainty and missing (non-reported) data.		animal species > 750	plant species > 120
E	ANNUAL NITROGEN FIXATION	NUCLEAR FALLOUT	HuID: 86866
fixed mass of nitrogen $\approx 1.5 \times 10^{11} \text{ kg / yr}$	HuID: 60580; 78152	239+240Pu activity in soil $\approx 1 \text{ Bq / kg}$	HuID: 38748; 91171
Data Source(s): USGS Mineral Commodities Summaries (Fixed Nitrogen), January 2020; Table 2 of "World fertilizer trends and outlook to 2022" Food and Agricultural Organization of the United Nations, 2019, ISBN: 978-92-5-131894-2. Smit et al. 2010, DOI:10.1039/c9ee02873k. Notes: The approximate mass of contained nitrogen in salnitre ammonia produced globally in 2018 as reported by the USGS is $\approx 144 \text{ Mt}$. This value is in moderate agreement with the forecast of $\approx 160 \text{ Mt}$ of nitrogen-contained ammonia as forecast for 2018 by the FAO. Approximately all of this mass is produced by the Haber-Bosch process ($> 96\%$, Smith et al. 2020).		Data Source(s): Figure 4 and Figure 5 in Hancock et al. 2014, DOI:10.1144/SP395.15. Figure 3 (col. 2, rows 3 – 5) of Ciszewski and Łokas, 2019, DOI:10.1515/geochr-2015-0111. Notes: Value corresponds to current-day detectable combined radioactivity of ²³⁹ Pu and ²⁴⁰ Pu present in cores of stratified soil with estimated date of $\approx 1963 \text{ CE}$ during the peak of atmospheric nuclear weapons testing. Reported is approximate average activity from sediment samples in SE Australia (Hancock et al.) and Polish river basins (Ciszewski and Łokas).	Data Source(s): The IUCN Red List of Threatened Species. Version 2020–2. Notes: Values correspond to absolute lower-bound measurements of extinctions caused over the past ≈ 520 years. Of the predicted ≈ 8 million animal species, the IUCN databases catalogues only $\approx 900,000$ with only $\approx 75,000$ being assigned a conservation status. Representation of plants and fungi is even more sparse with only $\approx 40,000$ and ≈ 285 being assigned a conservation status, respectively. The number of extinct animal species is undoubtedly higher than these reported values, as signified by an inequality symbol (>).
change in [H ⁺] $\approx 0.2\% / \text{yr}$	HuID: 13934	EARTH MOVING	HuID: XXXX
Data Source(s): Figure 2 of European Environment Agency report CLIM 043 (2020). Original data source of report is "Global Mean Sea Water pH" from Copernicus Marine Environment Monitoring Service. Notes: Reported value is calculated from the average annual change in pH over years 1985–2018. Annual change in pH is $\approx 0.001 \text{ pH units}$, corresponding to a change in [H ⁺] of $\approx 0.2\% / \text{yr}$.		coal mining waste/overburden $\approx 6.5 \times 10^{13} \text{ kg / yr}$	HuID: 72899
F	OCEAN pH	urbanization $> 1.4 \times 10^{14} \text{ kg / yr}$	HuID: 59640
agriculture $\approx 5 \times 10^{13} \text{ m}^2 / \text{yr}$	HuID: 29582	Data Source(s): Table 1 of Cooper et al. 2018. DOI: doi.org/gfwfhhd. Notes: Coal mining waste and overburden mass is calculated given commodity-level stripping ratios (mass of overburden/waste per mass of coal resource mined) and reported values of global coal production by type. Urbanization mass is presented as a lower bound estimate of the mass of earth moved from global construction projects. This comes from a conservative estimate that the ratio of the mass of earth moved per mass of cement/concrete used in construction is 2:1. This value is highly context dependent and we encourage the reader to read the source material for a more thorough description of this estimation.	Data Source(s): The IUCN Red List of Threatened Species. Version 2020–2. Notes: Values correspond to absolute lower-bound measurements of extinctions caused over the past ≈ 520 years. Of the predicted ≈ 8 million animal species, the IUCN databases catalogues only $\approx 900,000$ with only $\approx 75,000$ being assigned a conservation status. Representation of plants and fungi is even more sparse with only $\approx 40,000$ and ≈ 285 being assigned a conservation status, respectively. The number of extinct animal species is undoubtedly higher than these reported values, as signified by an inequality symbol (>).
land use		agriculture $\approx 1.2 \times 10^{13} \text{ kg / yr}$	HuID: XXXXX
G	RIVER FRAGMENTATION	ACKNOWLEDGMENTS	
fragmented river volume $\approx 6 \times 10^{11} \text{ m}^3$	HuID: 61661; 15550	This work was supported by the Resnick Institute for Sustainability at the California Institute of Technology. Add financial support information. No specific order, thank the following: Joel Cohen, Alon Shepon, Eric Galbraith, Sean Carroll, Justin Bois, Shyam Saladi, APH150c course, Wati Taylor, Dan Fisher, Gidon Eshel, Greg Huber, Michelle Dan, Brad Marston, Hamza Raniwala, Yue Qin for providing data on water consumption. Suzy Beeler, Soichi Hirokawa, Manuel Razo-Mejia and all members of the Phillips lab at Caltech.	
H	ANNUAL ICE MELT		
glaciers $\approx 3 \times 10^{11} \text{ m}^3 / \text{yr}$	HuID: 32459		
Data Source(s): Intergovernmental Panel on Climate Change (IPCC) 2019 Special Report "The Ocean and Cryosphere in a Changing Climate," Table 2.A.1 on pp. 199–202. Notes: Value corresponds to the trend of annual mass loss from major glaciated regions (2006–2015). Volume loss was calculated from mass loss.			
ice sheets $\approx 4 \times 10^{11} \text{ m}^3 / \text{yr}$	HuID: 95798; 93137		
Data Source(s): D. N. Wiese et al. 2019 https://doi.org/10.5067/ITEM-SC-3MJ62 Notes: Value corresponds to the trends of combined annual mass loss from the Greenland and Antarctic Ice Sheets (2002–2020). Volume loss was calculated from mass loss.			
arctic sea ice $\approx 3 \times 10^{11} \text{ m}^3 / \text{yr}$	HuID: 89520		
Data Source(s): PIOMAS Arctic Sea Ice Volume Reanalysis, original method source: Schweiger et al. 2011 DOI: 10.2029/2011C007084 Notes: Value reported corresponds to the trend of decadal volume loss from Arctic sea ice (1979–2020) which was converted to annual volume loss.			
total $\approx 1 \times 10^{12} \text{ m}^3 / \text{yr}$	HuID: 89075		
Data Source(s): Antarctic sea ice loss is not included due to data sparsity. The periods of analysis are not the same, therefore this rate represents an approximation rather than an exact calculation.			
I	HUMAN POPULATION		
fraction of population that is urban $\approx 55\%$	HuID: 93995		
total population $\approx 7.6 \times 10^9$ people	HuID: 85255		
Data Source(s): Food and Agricultural Organization (FAO) of the United Nations Report on Annual Population, 2019. Notes: Value for total population comes from a combination of direct population reports from country governments as well as inferred measurements from under-reporting or missing data. Fraction of urban population is determined from what each country reports as "urban" which does not follow a singular definition. As explained by the United Nations population division, "When the definition used in the latest census was not the same as in previous censuses, the data were adjusted whenever possible so as to maintain consistency." Rural population is computed from this fraction along with the total human population, implying that the total population is composed only of "urban" and "rural" communities.			
J	GREENHOUSE GAS EMISSIONS		
anthropogenic CO ₂ $\approx 4.2 \times 10^{11} \text{ kg / yr}$	HuID: 47200; 98043		
Data Source(s): Friedlingstein et al. 2019, DOI: 10.5194/essd-11-1783-2019. Original data sources relevant to this study compiled in Friedlingstein et al.: 1) Gilfillan et al. https://energy.apgstate.edu/CDIAC 2) Average of two bookkeeping models: Houghton and Nassikas 2017 DOI: 10.1020/2016GB005546; Hansis et al. 2015 DOI: 10.5248/zenodo.1000000. Notes: Value corresponds to CO ₂ emissions from fossil fuel combustion, industrial emissions (predominantly cement production), and land-use change during calendar year 2018. CO ₂ was added to the atmosphere at a rate of $\approx 18.6 \text{ Gt / yr}$ in 2018 (HuID: 98043); most of the remainder is taken up by the land sink and ocean sink.			
anthropogenic CH ₄ $\approx 4 \times 10^{11} \text{ kg / yr}$	HuID: 96837; 56405; 30725		
Data Source(s): Table 2 of Saunois, et al. 2020. DOI: 10.5194/essd-12-1561-2020. Notes: Value corresponds to CH ₄ emissions from anthropogenic sources in the calendar year 2017. Represents emissions from agriculture and waste, fossil fuels, and biomass and biofuel burning. Value is not simply the sum of these sources but is based on a full anthropogenic inventory of emissions. Natural emissions amount to $\approx 0.3 \text{ Gt / yr}$ in 2017. CH ₄ was added to the atmosphere at a rate of $\approx 17 \text{ Mt / yr}$ in 2017; most of the remainder is taken up by chemical loss sink and soil sink.			
anthropogenic N ₂ O $\approx 7 \times 10^9 \text{ kg / yr}$	HuID: 44575; 68619		
Data Source(s): Table 1 of Tian, H., et al. (2020). DOI: 10.1038/s41586-020-2780-z Notes: Value corresponds to annualized N ₂ O emissions from anthropogenic sources in the years 2007–2016.			
K	WATER WITHDRAWAL & CONSUMPTION (DESCRIPTION TO BE UPDATED)		
agriculture $\approx 1.5 \times 10^{12} \text{ m}^3 / \text{yr}$	HuID: 43593		
power generation $\approx 3 \times 10^{11} \text{ m}^3 / \text{yr}$	HuID: 78784		
domestic $\approx 6 \times 10^{10} \text{ m}^3 / \text{yr}$	HuID: 69424		
total $\approx 2 \times 10^{12} \text{ m}^3 / \text{yr}$	HuID: 27342		
Data Source(s): Figure 1 of Qin et al. 2019, DOI:10.1038/s41893-019-0294-2. Notes: "Agricultural use" is defined as water used for irrigation, maintenance of livestock, and water used in the management of irrigation via damming. "Power generation" is defined as water used for thermal power generation (coal, nuclear, gas, biomass, oil, and other/waste) and hydroelectric generation. "Domestic" is defined as water directly used by humans and water used in the maintenance of municipal water supply. "Total" water use includes the above categories as well as other uses of water in reservoir management including flood control and other unannotated uses. All values pertain to estimates for 2016.			
L	TOTAL POWER CONSUMPTION		
global power consumption $\approx 17 \text{ TW}$	HuID: 31373		
Data Source(s): bp Statistical Review of World Energy, 2020. Notes: represents the sum total energy consumed from oil, natural gas, coal, and nuclear energy and electricity generated from hydroelectric and renewables			
M	FOREST LOSS		
commodity-driven $\approx 5 \times 10^{10} \text{ m}^2 / \text{yr}$	HuID: 96098		
forestry $\approx 6 \times 10^{10} \text{ m}^2 / \text{yr}$	HuID: 38352		
urbanization $\approx 2 \times 10^{10} \text{ m}^2 / \text{yr}$	HuID: 19429		
total $\approx 2 \times 10^{11} \text{ m}^2 / \text{yr}$	HuID: 78576		
Data Source(s): Table 1 and Figure 3 of Curtis et al. 2018 DOI:10.1126/science.aau3445. Hansen et al. 2013 DOI:10.1126/science.1244693. Global Forest Watch, 2020. Notes: Commodity-driven deforestation is defined as "long-term, permanent, conversion of forest and shrubland to a non-forest land use such as agriculture, mining, or energy infrastructure." Forest area disruption due to forestry is defined as large-scale forestry operations occurring within managed forests and tree plantations with evidence of forest regrowth in subsequent years." Forest land disruption due to urbanization is defined as "forest and shrubland conversion for the expansion and intensification of existing urban centers."			
N	SEA LEVEL RISE		
meltwater rise $\approx 1.6 \text{ mm / yr}$	HuID: 97108		
thermal expansion $\approx 1.2 \text{ mm / yr}$	HuID: 97688		
total annual sea-level rise $\approx 3.4 \text{ mm / yr}$	HuID: 81373		
Data Source(s): Table 1 of Frederikse et al. 2020. DOI:10.1088/1742-6596/1068/1/016809. Notes: Values correspond to the average global sea level rise of the years 1993 – 2018. Meltwater is defined as the global annual sea level rise due to melt of glaciers, the Greenland ice sheet, and the Antarctic ice sheet. "Other sources" accounts for sea level change from changes in the terrestrial water shed as well as for other factors not explicitly specified. Total sea level rise of $\approx 3.4 \text{ mm / yr}$ comes directly from measurement while values for meltwater, thermal expansion, and terrestrial water shed storage come from modeling.			
O	POWER FROM FOSSIL FUELS		
natural gas $\approx 4.5 \text{ TW}$	HuID: 49947		
oil $\approx 6.5 \text{ TW}$	HuID: 42121		
coal $\approx 5 \text{ TW}$	HuID: 10400		
total $\approx 16 \text{ TW}$	HuID: XXXXX		
Data Source(s): bp Statistical Review of World Energy, 2020. Notes: Values pertain to 2019 estimates only. Oil volume includes crude oil, shale oil, oil sands, condensates, and natural gas liquids separate from specific natural gas mining. Natural gas value excludes gas flared or recycled and includes natural gas produced for gas-to-liquids transformation. Coal value includes 2019 value exclusively for solid commercial fuels such as bituminous coal and anthracite, lignite and sub-bituminous coal, and other solid fuels. This includes coal used directly in power production as well as coal used in coal-to-liquids and coal-to-gas transformations.			