

White-Lined Gecko / Skunk Gecko Care Guide

Gekko vittatus

Red Rocks Exotic Reptiles · Reviewed October 2026 · Version 2.0

Provide a tall, well-furnished habitat with tight arboreal retreats and multiple routes between warmth and shade. These fast, vocal geckos can bite firmly and are primarily display animals.

Species Profile

Common name: White-Lined Gecko / Skunk Gecko

Scientific name: *Gekko vittatus*

Adult size: About 8-10 in (20-25 cm) total length

Captive lifespan: About 10-15 years is a keeper estimate; firm species-specific averages are limited. Typical captive lifespan is a planning estimate, not a measured statistical average.

Native range: New Guinea and nearby Indo-Pacific islands; verify locality

Habitat and activity: Nocturnal and strongly arboreal; trunks and sheltered forest climbing surfaces

Diet: Varied cultured insects; complete fruit gecko diets are not staples

Care difficulty: Intermediate

Shopping List

- Secure enclosure at least 18 × 18 × 36 in (46 × 46 × 91 cm; L × W × H) for one adult; see Housing for exceptions and group arrangements.
- Digital thermometers for different zones; a suitable hygrometer; timer for a consistent day/night cycle.
- Species-appropriate substrate and multiple stable hides or covered retreats.
- Reliable supply of the staple foods described below, clean feeding tools and a transport container.
- Appropriate overhead heat source if required, controlled by a compatible thermostat; infrared thermometer for surfaces.
- Appropriate visible lighting and, where recommended below, a linear UVB fixture with measured shade and exposure zones.
- Fresh-water dish or cup; insect supplements when applicable; a scale appropriate to the animal's size.

Housing

Our recommended starting adult enclosure is 18 × 18 × 36 in (46 × 46 × 91 cm; L × W × H) for one animal. Larger usable space is encouraged. These dimensions are a husbandry recommendation, not a species-specific experimentally proven threshold.

Fill the height with firmly secured horizontal and diagonal branches, cork, foliage and sheltered paths. Provide several resting places at different heights and enough cover to travel between environmental zones without crossing a bare open box.

Juveniles may use a furnished adult enclosure when food intake, hydration and waste can be monitored. A temporary grow-out enclosure should still offer choice and cover. Quarantine newcomers separately with dedicated tools and a substrate that permits health monitoring; arrange veterinary assessment for unknown-origin animals.

Lighting & UVB

Provide roughly 10-12 hours of visible daylight followed by darkness. A measured UV Index (UVI) around 0.5-1.0 at the most exposed usable site, declining to near zero inside retreats, is our conservative starting recommendation. Low-level UVB is recommended when a usable gradient can be provided; nocturnal activity alone is not a reason to exclude light.

Measure exposure at the animal's back height with an appropriate UVI meter, such as a Solarmeter 6.5. Lamp percentage, wattage and a fixed distance alone do not establish exposure: reflectors, mesh, perch height, plant growth and lamp age change output. Keep glass/plastic out of the UV path, follow fixture safety instructions and provide complete shade. Replace or reassess lamps according to measured output and manufacturer guidance.

Heating & Temperature

- Localized basking surface: 86-90°F (30-32°C).
- Cool-zone air: 72-78°F (22-26°C).
- Nighttime air: 70-75°F (21-24°C) for healthy established animals.

Create the gradient on one side rather than heating every hide to the same temperature. Use a compatible thermostat on each heat source; a manual lamp dimmer does not replace thermostatic protection. Secure the probe and verify temperatures independently: probe thermometers measure air/retreat temperature, while an infrared thermometer measures exposed surfaces.

Use appropriately guarded overhead warmth when needed. Keep animals clear of hot fixtures, avoid heat rocks, and switch visible lights off at night. Night heat is needed only when the room falls below the appropriate range; use a thermostat-controlled lightless source if necessary.

Humidity & Ventilation

55-70% while drying, 75-90% briefly after misting. Treat readings as a starting pattern rather than a number to chase regardless of substrate, airflow or animal behavior.

Measure humidity in a representative occupied zone, away from a freshly sprayed surface. Provide sheltered moisture choices while retaining airflow. Persistent condensation, sour odor, flooded soil or surfaces that never dry indicate a setup problem. Ventilation and hydration must be adjusted together rather than simply sealing all vents.

Water & Hydration

Provide a clean, appropriately shallow dish or accessible elevated water cup at all times. Change it daily and immediately when soiled; wash it regularly with dedicated tools. Potable water is generally suitable, with treatment where local water quality requires it.

For animals that drink droplets, mist foliage or suitable surfaces at the time they become active, then allow partial drying. Droplets supplement the dish; water chemistry claims should be based on evidence rather than a species' trade name.

Substrate

For an established animal, use 2-4 inches of a well-draining tropical substrate such as a pesticide- and fertilizer-free organic topsoil/coco blend, ReptiSoil-style mix, or ABG-type substrate, topped with clean

leaf litter. Bioactive enclosures should have enough drainage capacity to handle repeated misting without leaving the lower soil waterlogged.

Paper towel is appropriate for quarantine, medical monitoring, or when stool must be closely observed.

Avoid reptile carpet, pine or cedar products, large loose bark chunks, and other coarse materials that can trap toes or be difficult to pass if swallowed. Spot-clean feces and dead feeder insects promptly; refresh or replace non-bioactive substrate as it becomes soiled rather than waiting for odor or mold.

Food & Feeding

White-lined geckos are primarily insectivorous. Offer a varied rotation of appropriately sized live invertebrates rather than relying on one feeder species. Good staples include crickets, dubia or discoid roaches, black soldier fly larvae, silkworms, and appropriately sized mealworms; hornworms can be used for variety and hydration. Large, very fatty feeders should be occasional rather than routine.

Juveniles: offer appropriately sized prey daily or nearly daily, adjusting to growth and body condition.

Adults: feed about every other day or roughly 3-4 times per week. Hard-bodied prey should generally be no wider than the space between the gecko's eyes. Remove uneaten insects that could bother the gecko while it rests.

Wild white-lined geckos may opportunistically take small vertebrates, but that does not make rodents or feeder lizards necessary in captivity. Routine vertebrate prey adds excess calories and avoidable risk.

Crested gecko diet, fruit puree, and baby food should not be used as staple foods for this primarily insectivorous species.

Supplements

Lightly dust appropriate insect meals with calcium and use a balanced reptile multivitamin containing suitable vitamin A according to the product's directions. Gut-loading and variety support nutrition but do not eliminate the need to balance calcium-poor prey.

Match D3 supplementation to the actual UVB exposure and to vitamins already included in other products. UVB does not automatically mean zero dietary D3, and lack of UVB does not justify unlimited D3. Follow one coherent program; do not combine multiple strong vitamin products or coat every feeder heavily.

Decor & Enrichment

Decor is functional habitat for this species. Use cork rounds, cork flats, smooth sturdy branches, bamboo-sized perches, vines, ledges, broad leaves, and live or artificial plants to create covered routes through the whole enclosure. Hollow cork tubes are especially useful as secure daytime retreats. Provide several perch diameters and orientations so the gecko can climb, jump, hunt, hide, and choose among different temperatures and humidity levels.

Handling & Behavior

Treat this animal primarily as a display species. For transfers, gently guide it into a cup, tube or transport container instead of chasing and grabbing. Keep doors and escape routes controlled before opening the enclosure.

Allow new animals to settle and eat reliably. Avoid restraint by the tail, repeated uncovering or forced interactions. Track normal activity for the individual so changes in feeding, posture or hiding can be recognized.

Cleaning & Routine Maintenance

Daily: check the animal's access to water/appropriate moisture, temperatures, visible condition and equipment. Remove feces, contaminated material, dead feeders and spoiled food promptly.

Regularly: clean dishes and feeding surfaces, check thermostats and probes, inspect decor stability and maintain ventilation. Bioactive systems still require waste checks, food removal, plant care and drainage maintenance; they are not self-cleaning.

Record appetite, activity, shed/molt history and weight or body condition when safe. Arrange veterinary assessment for unexplained changes. Quarantine new animals with dedicated tools, and disinfect only with a suitable product used at the label dilution/contact time, then rinse and dry as required before return.

Cohabitation

Individual housing is the safest routine default. Shared housing can cause food competition, injury or reproductive pressure even without obvious fighting. Never assume a male/female pair or female group is compatible. Breeding introductions belong in a planned specialist program with separate backup enclosures and individual monitoring.

Shedding, Molting & Seasonal Changes

Maintain access to a clean humid retreat and inspect toes, tail tip and eyes after shedding when observation is possible. Do not peel retained skin or apply oils/adhesives. Persistent constriction, damaged tissue or repeated failed sheds warrants veterinary help.

Seasonal changes in appetite or activity can occur. Check temperatures, hydration and weight before treating prolonged refusal as normal. Deliberate brumation or breeding cycles require healthy animals and species-specific preparation; pet care does not require forcing a seasonal shutdown.

Reproduction

Mature females may develop eggs, including infertile eggs without a current male. Provide a suitable moist laying retreat or substrate access, monitor body condition and nutrition, and seek veterinary advice for prolonged straining, swelling or decline. Breeding, incubation and offspring rearing require a prepared species-specific program.

Common Health Concerns

- Poor body condition or feeding decline: check the full gradient, hydration and food quality; arrange assessment for persistent refusal accompanied by weight loss.
- Calcium/vitamin imbalance: weakness, tremors, a soft or misshapen jaw or abnormal limbs require veterinary assessment; extra high-dose supplements are not a substitute for diagnosis.
- Skin, shed or injury problems: repeated retained skin, wounds, swelling or abnormal discoloration deserve prompt attention.

- Respiratory or systemic illness: labored/open-mouth breathing at rest, marked lethargy, loss of balance, persistent diarrhea or rapid deterioration require an experienced exotic veterinarian.
- Unknown-origin or wild-caught animals may carry parasites. Fecal testing and veterinary-directed treatment are preferable to indiscriminate preventive medication.

Sources & Review Information

Reviewed October 2026 · RRE version 2.0. Care targets are practical starting recommendations informed by natural history, published work and captive experience. For less-studied species, enclosure dimensions, feeding frequencies, UVI ranges and lifespan estimates are not experimentally validated universal thresholds. Adjust for the individual, locality, life stage and veterinary advice.

The Reptile Database. Species taxonomy, distribution and cited natural-history literature.
<https://reptile-database.reptarium.cz/Gekko/vittatus>

Baines et al. (2016). The UV-Tool: choosing UV gradients using natural history and behavior. General lighting framework; not a species-specific dose trial.
<https://doi.org/10.19227/jzar.v4i1.150>

RED ROCKS
 EXOTIC REPTILES
 —★ ★ ★ ★—