
INTEGRATING LAGRANGIAN SUBMANIFOLDS WITH MULTIPLICITY

Ryan O. Van Gelder

Citizen Gardens - The Foundation of Multiplicity

info@citizengardens.org

ABSTRACT

This paper presents a mathematical integration of Lagrangian submanifolds within the framework of multiplicity theory. By extending the classical variational principles to incorporate eigenvalue multiplicity, recursive feedback, and tensor dynamics, this approach models complex systems with applications in quantum mechanics, machine learning, and advanced geometries.

1 Introduction

Lagrangian submanifolds play a fundamental role in symplectic geometry and mechanics. Multiplicity theory enhances this by incorporating eigenvalue dynamics, recursive feedback loops, and tensor coupling. We define the primary components and their interactions below.

2 Mathematical Framework

The dynamic multiplicity equation governing the evolution of densities ρ_k on Lagrangian submanifolds is given as:

$$\frac{\partial \rho_k}{\partial t} = \alpha_k(t)\rho_k + \beta_k(t)I_k + \gamma_k(t) \sum_j T_{kj}\rho_j + \lambda(t)(\Omega_B(\rho) + \Omega_{FS}(\rho)) + \eta_k\rho_k^2 + \zeta_k \sum_{m,n} C_{mn}\rho_m\rho_n + \xi_k(t). \quad (1)$$

Here:

- $\alpha_k(t), \beta_k(t), \gamma_k(t), \lambda(t)$: Time-dependent parameters.
- $\Omega_B(\rho), \Omega_{FS}(\rho)$: Dynamical geometric feedback terms.
- T_{kj} : Coupling tensor for multi-scale interactions.
- ζ_k : Coefficients for quantum entanglement terms.

- $\xi_k(t)$: Stochastic noise term.

3 Lagrangian Submanifolds in Multiplicity

Let $\mathcal{L} \subset \mathcal{M}$ be a Lagrangian submanifold of the symplectic manifold $(\mathcal{M}, \omega)$, with the symplectic form $\omega = d\theta$. The multiplicity-enhanced action integral is:

$$S[\mathcal{L}] = \int_{\mathcal{L}} \left(\frac{1}{2} \omega \wedge \omega + \lambda \Omega_B(\rho) + \Omega_{FS}(\rho) \right) d\mu, \quad (2)$$

where $d\mu$ is the invariant measure on $\mathcal{L}$.

4 Tensor Networks and Feedback Dynamics

The interaction between eigenstates and tensors is represented by:

$$T_{ijk} = \sum_{p,q} \psi_p \psi_q \otimes \rho_k. \quad (3)$$

The recursive feedback loop modifies the geometric evolution:

$$\frac{\partial \rho_k}{\partial t} = F(\rho_k, T_{ijk}, \Omega_B, \Omega_{FS}). \quad (4)$$

5 Applications of Multiplicity-Enhanced Lagrangian Submanifolds

The integration of multiplicity with Lagrangian submanifolds enables advanced modeling and problem-solving in various fields. Below, we provide mathematical formulations for key applications.

5.1 Quantum Gravity

The quantum-corrected Einstein Field Equations (EFE) incorporating multiplicity are:

$$R_{\mu\nu} - \frac{1}{2} g_{\mu\nu} R + \Lambda g_{\mu\nu} + \epsilon \cdot \nabla^2 Q_{\mu\nu} = 8\pi G \langle T_{\mu\nu} \rangle_{\text{quantum}}, \quad (5)$$

where:

- $Q_{\mu\nu}$: Quantum correction tensor,
- ϵ : Small coupling parameter for quantum fluctuations,
- $\langle T_{\mu\nu} \rangle_{\text{quantum}}$: Quantum-averaged stress-energy tensor.

This form enhances classical GR by integrating quantum corrections and is particularly effective in:

- Modeling black hole evaporation and Hawking radiation.

- Describing quantum fluctuations in spacetime during inflation.
- Capturing the interplay of classical and quantum gravitational phenomena.

5.2 Machine Learning

Tensor-based representations of data with multiplicity can be expressed as:

$$T_{ijk} = \sum_{p,q} \psi_p \psi_q \otimes \rho_k, \quad (6)$$

where:

- T_{ijk} : Tensor capturing multi-modal interactions,
- ψ_p, ψ_q : Feature vectors from training data,
- ρ_k : Multiplicity-enhanced state vector.

The recursive feedback for optimizing the model is given as:

$$\frac{\partial \rho_k}{\partial t} = \alpha_k \rho_k + \beta_k T_{ijk} \psi_i \psi_j, \quad (7)$$

where α_k and β_k are learning rates dependent on multiplicity and dataset complexity.

5.3 Material Science

The interaction potential between particles, incorporating multiplicity and tensor dynamics, is described by:

$$V(\mathbf{r}) = \sum_{i,j} T_{ij} \phi_i(\mathbf{r}) \phi_j(\mathbf{r}), \quad (8)$$

where:

- $V(\mathbf{r})$: Potential energy as a function of position $\mathbf{r}$,
- T_{ij} : Tensor capturing interactions between particles i and j ,
- $\phi_i(\mathbf{r})$: Wavefunction of particle i .

The dynamic evolution of the material's state is expressed as:

$$\frac{\partial \phi_i(\mathbf{r}, t)}{\partial t} = \frac{-i}{\hbar} \left[H \phi_i(\mathbf{r}, t) + \lambda \sum_k T_{ik} \phi_k(\mathbf{r}, t) \right], \quad (9)$$

where H is the Hamiltonian operator.

5.4 Black Hole Dynamics

Using tensor networks to describe entanglement across horizons, the quantum state of a black hole evolves as:

$$\Psi_{\text{BH}}(t) = \sum_{i,j} T_{ij} \Psi_i \otimes \Psi_j e^{i(\theta_i + \theta_j)}, \quad (10)$$

where:

- T_{ij} : Tensor representing entanglement structure,
- Ψ_i, Ψ_j : Quantum states near the horizon,
- θ_i : Phase of the quantum state.

This formulation integrates multiplicity to model quantum coherence and Hawking radiation effects.

Summary of Applications

The multiplicity-enhanced Lagrangian framework offers novel tools for addressing challenges in physics, computational modeling, and material science. By leveraging tensor dynamics, recursive feedback, and eigenvalue multiplicity, it provides scalable and robust solutions for analyzing complex systems.

6 Conclusion

This framework integrates Lagrangian submanifolds with the principles of multiplicity, paving the way for innovative applications across disciplines.

References

1. Michael A Nielsen and Isaac L Chuang. *Quantum Computation and Quantum Information*. Cambridge University Press, 2010.
2. Michael V Berry. Quantal phase factors accompanying adiabatic changes. *Proceedings of the Royal Society of London. A. Mathematical and Physical Sciences*, 392(1802):45–57, 1984.
3. Erwin Schrödinger. Discussion of probability relations between separated systems. *Mathematical Proceedings of the Cambridge Philosophical Society*, 31(4):555–563, 1935.
4. Albert-László Barabási. *Network Science*. Cambridge University Press, 2016.
5. Jutho Haegeman, J Ignacio Cirac, Tobias J Osborne, Henri Verschelde, and Frank Verstraete. Time-dependent variational principle for quantum lattices. *Physical Review Letters*, 107(7):070601, 2011.
6. Guido Fubini. Sugli spazi a curvatura costante. *Rendiconti del Circolo Matematico di Palermo (1884-1940)*, 21:1–13, 1906.

7. David Bohm. A suggested interpretation of the quantum theory in terms of “hidden” variables. i. *Physical Review*, 85(2):166–179, 1952.
8. Fritjof Capra and Pier Luigi Luisi. *The Systems View of Life: A Unifying Vision*. Cambridge University Press, 2014.
9. Melanie Mitchell. *Complexity: A Guided Tour*. Oxford University Press, 2009.
10. Yoshua Bengio. Learning deep architectures for ai. *Foundations and Trends in Machine Learning*, 2(1):1–127, 2009.
11. Nicholas Galioto and Ryan O. Van Gelder. Dynamic multiplicity equation.
https://github.com/CitizenGardens-org/Multiplicity/blob/Multiplicity/Mathematics/Dynamic_Multiplicity_Equation.pdf, 2024. Accessed: 2025-01-13.