

Change Detection For High-Resolution Satellite Images By Using Gaussian Mixture Model


1. Introduction

In modern times, human civilization's settlement landscapes and patterns are changing rapidly because of the growth of human population, acceleration of urbanization and the application of varied technologies. Keep detecting and assessing these changes and mastering the accurate information are quite important for civilized human development. High resolution images provided by satellites are good resource to use to identify and quantify landscapes changes. We explore a detection method to identify landscape changes using high resolution satellite images. This grid based method is helpful in Bi-temporal change detection. By given two satellite images from the same area, it can identify changes accurately.

2. Data description


The original data are two high resolution satellite images about the same area with some artificial and natural objects from different time. Inside the images, there are some changes between older and newer ones.

Before implementing our method to the data, we need to do a pre-processing procedure. After executing some rotation and crop steps, these two images will be ready to use.



Shijie Li, Tianpei Xia, Jingjuan Deng, Zhiren Lu, Zifan Nan, Rui Liu
North Carolina State University

3. Methodology




After dividing images into the grids properly, we convert all of these grids into their relative Gaussian distributions.

$$P(x|y_i) = \frac{1}{\sqrt{(2\pi)^{-N} |\Sigma_j|}} e^{-\frac{1}{2}(x-\mu_j)^t |\Sigma_j|^{-1} (x-\mu_j)}$$

We compute the Kullback-Leibler(KL) divergence between each pair of Gaussian distributions, then we can create a matrix map of KL divergence distance:


We model the image data into grids. Given two satellite images (one old and one new) which describe exactly the same land area, we divide these two images into two square grid clusters. The grids have the same size and each image has the same number of grids.




Using this KL divergence matrix, we model the D_{KL} map into Gaussian Mixture Model, then we can apply this clustering model to categorize the changes on KL divergence map, plot the change map to detect changes.

4. Results

To compare the changes between two images, we overlaid this GMM change cluster map onto the original satellite maps. the red and green rectangle grids suggest the areas has relatively greater changes, all other parts are the areas that basically remain the same.



5. Parameter choices



Divide each image into square grids of RGB pixel blocks, which affects the quality of object recognition. Grid size is tuned based on final cluster pattern with expert knowledge.

6. Conclusions

In our project, we presented a GMM based change detection method for satellite images. Our method can detect a quite part of artificial changes between different high resolution images data, although to some extent, it still has some inaccuracy issues about detecting changing details.

It's always a very challenging task to keep improving the performance for some new technologies. However, we believe this method will have good prospects for further research.

7. References

- [1] Vatsavai, R. Graesser, J. 2012. "Probabilistic Change Detection Framework for Analyzing Settlement Dynamics Using Very High-resolution Satellite Imagery. ICCS: 907-916.
- [2] Hershey, John R., and Peder A. Olsen. "Approximating the Kullback Leibler divergence between Gaussian mixture models." 2007 IEEE International Conference on Acoustics, Speech and Signal Processing-ICASSP'07. Vol. 4. IEEE, 2007.

Github Link:
https://github.ncsu.edu/sli41/CSC522_Project