



[Home](#)

2. [Vibramycin](#)

3. [FDA PI](#)

Vibramycin: Package Insert / Prescribing Info

Package insert / product label

Generic name: [doxycycline](#)

Dosage form: tablet, capsule, syrup, powder for oral suspension

Drug classes: [Miscellaneous antimalarials](#), [Tetracyclines](#)

[Medically reviewed](#) by Drugs.com. Last updated on Jul 22, 2024.

On This Page

[Description](#)

[Clinical Pharmacology](#)

[Indications and Usage](#)

[Contraindications](#)

[Warnings](#)

[Precautions](#)

[Patient Counseling Information](#)

[Drug Interactions](#)

[Adverse Reactions/Side Effects](#)

[Overdosage](#)

[Dosage and Administration](#)

[How Supplied/Storage and Handling](#)

[Storage and Handling](#)


[References](#)

To reduce the development of drug-resistant bacteria and maintain the effectiveness of Vibramycin® and other antibacterial drugs, Vibramycin should be used only to treat or prevent infections that are proven or strongly suspected to be caused by bacteria.

Vibramycin Description

Vibramycin is an antibacterial drug synthetically derived from oxytetracycline, and is available as Vibramycin Monohydrate (doxycycline monohydrate); Vibramycin Hyclate and Vibra-Tabs (doxycycline hydrochloride hemiethanolate hemihydrate); and Vibramycin Calcium (doxycycline calcium) for oral administration.

The structural formula of doxycycline monohydrate is



with a molecular formula of $C_{22}H_{24}N_2O_8 \cdot H_2O$ and a molecular weight of 462.46. The chemical designation for doxycycline is 4-(Dimethylamino)-1,4,4a,5,5a,6,11,12a-octahydro-3,5,10,12,12a-pentahydroxy-6-methyl-1,11-dioxo-2-naphthacenecarboxamide monohydrate. The molecular formula for doxycycline hydrochloride hemiethanolate hemihydrate is $(C_{22}H_{24}N_2O_8 \cdot HCl)_2 \cdot C_2H_6O \cdot H_2O$ and the molecular weight is 1025.89. Doxycycline is a light-yellow crystalline powder. Doxycycline hyclate is soluble in water, while doxycycline monohydrate is very slightly soluble in water.

Doxycycline has a high degree of lipid solubility and a low affinity for calcium binding. It is highly stable in normal human serum. Doxycycline will not degrade into an epianhydro form.

Inert ingredients in the syrup formulation are: apple flavor; butylparaben; calcium chloride; carmine; glycerin; hydrochloric acid; magnesium aluminum silicate; povidone; propylene glycol; propylparaben; raspberry flavor; simethicone emulsion; sodium hydroxide; sodium metabisulfite; sorbitol solution; water.

Inert ingredients in the capsule formulations are: hard gelatin capsules (which may contain Blue 1 and other inert ingredients); magnesium stearate; microcrystalline cellulose; sodium lauryl sulfate.

Inert ingredients for the oral suspension formulation are: carboxymethylcellulose sodium; Blue 1; methylparaben; microcrystalline cellulose; propylparaben; raspberry flavor; Red 28; simethicone emulsion; sucrose.

Inert ingredients for the tablet formulation are: ethylcellulose; hypromellose; magnesium stearate; microcrystalline cellulose; propylene glycol; sodium lauryl sulfate; talc; titanium dioxide; Yellow 6 Lake.

Vibramycin - Clinical Pharmacology

Tetracyclines are readily absorbed and are bound to plasma proteins in varying degree. They are concentrated by the liver in the bile, and excreted in the urine and feces at high concentrations and in a biologically active form. Doxycycline is virtually completely absorbed after oral administration.

Following a 200 mg dose, normal adult volunteers averaged peak serum levels of 2.6 mcg/mL of doxycycline at 2 hours, decreasing to 1.45 mcg/mL at 24 hours. Excretion of doxycycline by the kidney is about 40%/72 hours in individuals with normal function (creatinine clearance about 75 mL/min.). This percentage excretion may fall as low as 1–5%/72 hours in individuals with severe renal insufficiency (creatinine clearance below 10 mL/min.). Studies have shown no significant difference in serum half-life of doxycycline (range 18–22 hours) in individuals with normal and severely impaired renal function.

Hemodialysis does not alter serum half-life.

Results of animal studies indicate that tetracyclines cross the placenta and are found in fetal tissues.

Population pharmacokinetic analysis of sparse concentration-time data of doxycycline following standard of care intravenous and oral dosing in 44 pediatric patients (2-18 years of age) showed that allometrically -scaled clearance (CL) of doxycycline in pediatric patients ≥ 2 to ≤ 8 years of age (median [range] 3.58 [2.27-10.82] L/h/70 kg, N =11) did not

differ significantly from pediatric patients >8 to 18 years of age (3 .27 [1.11-8.12] L/h/70 kg, N=33). For pediatric patients weighing ≤45 kg, body weight normalized doxycycline CL in those ≥2 to ≤8 years of age (median [range] 0.071 [0 .041-0.202] L/kg/h, N=10) did not differ significantly from those >8 to 18 years of age (0.081 [0.035-0.126] L/kg/h, N=8). In pediatric patients weighing >45 kg, no clinically significant differences in body weight normalized doxycycline CL were observed between those ≥2 to ≤8 years (0.050 L/kg/h, N=1) and those >8 to 18 years of age (0.044 [0.014-0.121] L/kg/h, N=25). No clinically significant difference in CL between oral and IV dosing was observed in the small cohort of pediatric patients who received the oral (N=19) or IV (N=21) formulation alone.

Microbiology

Mechanism of Action

Doxycycline inhibits bacterial protein synthesis by binding to the 30S ribosomal subunit. Doxycycline has bacteriostatic activity against a broad range of Gram-positive and Gram-negative bacteria.

Resistance

Cross resistance with other tetracyclines is common.

Antimicrobial Activity

Doxycycline has been shown to be active against most isolates of the following microorganisms, both in vitro and in clinical infections as described in the INDICATIONS AND USAGE section of the package insert for Vibramycin.

Gram-Negative Bacteria

Acinetobacter species
Bartonella bacilliformis
Brucella species
Klebsiella species
Klebsiella granulomatis
Campylobacter fetus
Enterobacter aerogenes
Escherichia coli
Francisella tularensis
Haemophilus ducreyi
Haemophilus influenzae
Neisseria gonorrhoeae
Shigella species
Vibrio cholerae
Yersinia pestis

Gram-Positive Bacteria

Bacillus anthracis
Listeria monocytogenes
Streptococcus pneumoniae

Anaerobic Bacteria

Clostridium species
Fusobacterium fusiforme
Propionibacterium acnes

Other Bacteria

Nocardiae and other aerobic Actinomyces species

Borrelia recurrentis

Chlamydophila psittaci

Chlamydia trachomatis

Mycoplasma pneumoniae

Rickettsiae

Treponema pallidum

Treponema pallidum subspecies pertenue

Ureaplasma urealyticum

Parasites

Balantidium coli

Entamoeba species

Plasmodium falciparum¹

- ¹ Doxycycline has been found to be active against the asexual erythrocytic forms of Plasmodium falciparum, but not against the gametocytes of P. falciparum. The precise mechanism of action of the drug is not known.

Susceptibility Testing

For specific information regarding susceptibility test interpretive criteria and associated test methods and quality control standards recognized by FDA for this drug, please see: <https://www.fda.gov/STIC>.

Indications and Usage for Vibramycin

To reduce the development of drug-resistant bacteria and maintain effectiveness of Vibramycin and other antibacterial drugs, Vibramycin should be used only to treat or prevent infections that are proven or strongly suspected to be caused by susceptible bacteria. When culture and susceptibility information are available, they should be considered in selecting or modifying antibacterial therapy. In the absence of such data, local epidemiology and susceptibility patterns may contribute to the empiric selection of therapy.

Treatment

Doxycycline is indicated for the treatment of the following infections:

- Rocky Mountain spotted fever, typhus fever and the typhus group, Q fever, rickettsialpox, and tick fevers caused by Rickettsiae.
- Respiratory tract infections caused by Mycoplasma pneumoniae.
- Lymphogranuloma venereum caused by Chlamydia trachomatis.
- Psittacosis (ornithosis) caused by Chlamydophila psittaci.
- Trachoma caused by Chlamydia trachomatis, although the infectious agent is not always eliminated, as judged by immunofluorescence.
- Inclusion conjunctivitis caused by Chlamydia trachomatis.
- Uncomplicated urethral, endocervical, or rectal infections in adults caused by Chlamydia trachomatis.
- Nongonococcal urethritis caused by Ureaplasma urealyticum.

- Relapsing fever due to *Borrelia recurrentis*.

Doxycycline is also indicated for the treatment of infections caused by the following gram-negative microorganisms:

- Chancroid caused by *Haemophilus ducreyi*.
- Plague due to *Yersinia pestis*.
- Tularemia due to *Francisella tularensis*.
- Cholera caused by *Vibrio cholerae*.
- *Campylobacter fetus* infections caused by *Campylobacter fetus*.
- Brucellosis due to *Brucella* species (in conjunction with streptomycin).
- Bartonellosis due to *Bartonella bacilliformis*.
- Granuloma inguinale caused by *Klebsiella granulomatis*.

Because many strains of the following groups of microorganisms have been shown to be resistant to doxycycline, culture and susceptibility testing are recommended.

Doxycycline is indicated for treatment of infections caused by the following gram-negative bacteria, when bacteriologic testing indicates appropriate susceptibility to the drug:

- *Escherichia coli*.
- *Enterobacter aerogenes*.
- *Shigella* species.
- *Acinetobacter* species.
- Respiratory tract infections caused by *Haemophilus influenzae*.
- Respiratory tract and urinary tract infections caused by *Klebsiella* species.

Doxycycline is indicated for treatment of infections caused by the following gram-positive microorganisms when bacteriologic testing indicates appropriate susceptibility to the drug:

- Upper respiratory infections caused by *Streptococcus pneumoniae*.
- Anthrax due to *Bacillus anthracis*, including inhalational anthrax (post-exposure): to reduce the incidence or progression of disease following exposure to aerosolized *Bacillus anthracis*.

When penicillin is contraindicated, doxycycline is an alternative drug in the treatment of the following infections:

- Uncomplicated gonorrhea caused by *Neisseria gonorrhoeae*.
- Syphilis caused by *Treponema pallidum*.
- Yaws caused by *Treponema pallidum* subspecies *pertenue*.
- Listeriosis due to *Listeria monocytogenes*.
- Vincent's infection caused by *Fusobacterium fusiforme*.
- Actinomycosis caused by *Actinomyces israelii*.
- Infections caused by *Clostridium* species.

In acute intestinal amebiasis, doxycycline may be a useful adjunct to amebicides.

In severe acne, doxycycline may be useful adjunctive therapy.

Prophylaxis

Doxycycline is indicated for the prophylaxis of malaria due to *Plasmodium falciparum* in short-term travelers (<4 months) to areas with chloroquine and/or pyrimethamine-sulfadoxine resistant strains. (See [DOSAGE AND ADMINISTRATION](#) section and [Information for Patients](#) subsection of the PRECAUTIONS section.)

Contraindications

This drug is contraindicated in persons who have shown hypersensitivity to any of the tetracyclines.

Warnings

The use of drugs of the tetracycline class during tooth development (last half of pregnancy, infancy and childhood to the age of 8 years) may cause permanent discoloration of the teeth (yellow-gray-brown). This adverse reaction is more common during long-term use of the drugs, but it has been observed following repeated short-term courses. Enamel hypoplasia has also been reported. Use doxycycline in pediatric patients 8 years of age or less only when the potential benefits are expected to outweigh the risks in severe or life-threatening conditions (e.g., anthrax, Rocky Mountain spotted fever), particularly when there are no alternative therapies.

Clostridioides difficile associated diarrhea (CDAD) has been reported with use of nearly all antibacterial agents, including Vibramycin, and may range in severity from mild diarrhea to fatal colitis. Treatment with antibacterial agents alters the normal flora of the colon leading to overgrowth of *C. difficile*.

C. difficile produces toxins A and B which contribute to the development of CDAD. Hypertoxin producing strains of *C. difficile* cause increased morbidity and mortality, as these infections can be refractory to antimicrobial therapy and may require colectomy. CDAD must be considered in all patients who present with diarrhea following the use of antibacterial drugs. Careful medical history is necessary since CDAD has been reported to occur over two months after the administration of antibacterial agents.

If CDAD is suspected or confirmed, ongoing use of antibacterial drugs not directed against *C. difficile* may need to be discontinued. Appropriate fluid and electrolyte management, protein supplementation, antibacterial treatment of *C. difficile*, and surgical evaluation should be instituted as clinically indicated.

Severe skin reactions, such as exfoliative dermatitis, erythema multiforme, Stevens-Johnson syndrome, toxic epidermal necrolysis, and drug reaction with eosinophilia and systemic symptoms (DRESS) have been reported in patients receiving doxycycline. (See [ADVERSE REACTIONS](#).) If severe skin reactions occur, doxycycline should be discontinued immediately and appropriate therapy should be instituted.

Intracranial hypertension (IH, pseudotumor cerebri) has been associated with the use of tetracyclines including Vibramycin. Clinical manifestations of IH include headache, blurred vision, diplopia, and vision loss; papilledema can be found on fundoscopy. Women of childbearing age who are overweight or have a history of IH are at greater risk for developing tetracycline associated IH. Concomitant use of isotretinoin and Vibramycin should be avoided because isotretinoin is also known to cause pseudotumor cerebri.

Although IH typically resolves after discontinuation of treatment, the possibility for permanent visual loss exists. If visual disturbance occurs during treatment, prompt ophthalmologic evaluation is warranted. Since intracranial pressure can

remain elevated for weeks after drug cessation patients should be monitored until they stabilize.

All tetracyclines form a stable calcium complex in any bone-forming tissue. A decrease in fibula growth rate has been observed in prematures given oral tetracycline in doses of 25 mg/kg every 6 hours. This reaction was shown to be reversible when the drug was discontinued.

Results of animal studies indicate that tetracyclines cross the placenta, are found in fetal tissues, and can have toxic effects on the developing fetus (often related to retardation of skeletal development). Evidence of embryotoxicity has also been noted in animals treated early in pregnancy. If any tetracycline is used during pregnancy or if the patient becomes pregnant while taking this drug, the patient should be apprised of the potential hazard to the fetus.

The antianabolic action of the tetracyclines may cause an increase in BUN. Studies to date indicate that this does not occur with the use of doxycycline in patients with impaired renal function.

Photosensitivity manifested by an exaggerated sunburn reaction has been observed in some individuals taking tetracyclines. Patients apt to be exposed to direct sunlight or ultraviolet light should be advised that this reaction can occur with tetracycline drugs, and treatment should be discontinued at the first evidence of skin erythema.

Vibramycin Syrup contains sodium metabisulfite, a sulfite that may cause allergic-type reactions, including anaphylactic symptoms and life-threatening or less severe asthmatic episodes in certain susceptible people. The overall prevalence of sulfite sensitivity in the general population is unknown and probably low. Sulfite sensitivity is seen more frequently in asthmatic than in non-asthmatic people.

Precautions

General

As with other antibacterial drugs, use of Vibramycin may result in overgrowth of nonsusceptible organisms, including fungi. If superinfection occurs, Vibramycin should be discontinued and appropriate therapy instituted.

Incision and drainage or other surgical procedures should be performed in conjunction with antibacterial therapy, when indicated.

Doxycycline offers substantial but not complete suppression of the asexual blood stages of Plasmodium strains.

Doxycycline does not suppress *P. falciparum*'s sexual blood stage gametocytes. Subjects completing this prophylactic regimen may still transmit the infection to mosquitoes outside endemic areas.

Prescribing Vibramycin in the absence of proven or strongly suspected bacterial infection or a prophylactic indication is unlikely to provide benefit to the patient and increases the risk of the development of drug-resistant bacteria.

Information For Patients

Patients taking doxycycline for malaria prophylaxis should be advised:

- that no present-day antimalarial agent, including doxycycline, guarantees protection against malaria.
- to avoid being bitten by mosquitoes by using personal protective measures that help avoid contact with mosquitoes, especially from dusk to dawn (e.g., staying in well-screened areas, using mosquito nets, covering the body with clothing, and using an effective insect repellent).
- that doxycycline prophylaxis:

- should begin 1–2 days before travel to the malarious area,
- should be continued daily while in the malarious area and after leaving the malarious area,
- should be continued for 4 further weeks to avoid development of malaria after returning from an endemic area,
- should not exceed 4 months.

All patients taking doxycycline should be advised:

- to avoid excessive sunlight or artificial ultraviolet light while receiving doxycycline and to discontinue therapy if phototoxicity (e.g., skin eruption, etc.) occurs. Sunscreen or sunblock should be considered. (See [WARNINGS](#).)
- to drink fluids liberally along with doxycycline to reduce the risk of esophageal irritation and ulceration. (See [ADVERSE REACTIONS](#).)
- that the absorption of tetracyclines is reduced when taken with foods, especially those which contain calcium. However, the absorption of doxycycline is not markedly influenced by simultaneous ingestion of food or milk. (See [DRUG INTERACTIONS](#).)
- that the absorption of tetracyclines is reduced when taking bismuth subsalicylate. (See [DRUG INTERACTIONS](#).)
- that the use of doxycycline might increase the incidence of vaginal candidiasis.

Patients should be counseled that antibacterial drugs, including Vibramycin should only be used to treat bacterial infections. They do not treat viral infections (e.g., the common cold). When Vibramycin is prescribed to treat a bacterial infection, patients should be told that although it is common to feel better early in the course of therapy, the medication should be taken exactly as directed. Skipping doses or not completing the full course of therapy may (1) decrease the effectiveness of the immediate treatment and (2) increase the likelihood that bacteria will develop resistance and will not be treatable by Vibramycin or other antibacterial drugs in the future.

Diarrhea is a common problem caused by antibacterial drugs, which usually ends when the antibacterials are discontinued. Sometimes after starting treatment with antibacterial drugs, patients can develop watery and bloody stools (with or without stomach cramps and fever) even as late as two or more months after having taken the last dose of the antibacterial drug. If this occurs, patients should contact their physician as soon as possible.

Laboratory Tests

In venereal disease, when co-existent syphilis is suspected, dark field examinations should be done before treatment is started and the blood serology repeated monthly for at least 4 months.

In long-term therapy, periodic laboratory evaluation of organ systems, including hematopoietic, renal, and hepatic studies, should be performed.

Drug Interactions

Because tetracyclines have been shown to depress plasma prothrombin activity, patients who are on anticoagulant therapy may require downward adjustment of their anticoagulant dosage.

Since bacteriostatic drugs may interfere with the bactericidal action of penicillin, it is advisable to avoid giving tetracyclines in conjunction with penicillin.

Absorption of tetracyclines is impaired by antacids containing aluminum, calcium, or magnesium, and iron-containing preparations.

Absorption of tetracyclines is impaired by bismuth subsalicylate.

Barbiturates, carbamazepine, and phenytoin decrease the half-life of doxycycline.

The concurrent use of tetracycline and Penthrane[®] (methoxyflurane) has been reported to result in fatal renal toxicity.

Concurrent use of tetracycline may render oral contraceptives less effective.

Drug/Laboratory Test Interactions

False elevations of urinary catecholamine levels may occur due to interference with the fluorescence test.

Carcinogenesis, Mutagenesis, Impairment Of Fertility

Long-term studies in animals to evaluate carcinogenic potential of doxycycline have not been conducted. However, there has been evidence of oncogenic activity in rats in studies with the related antibacterial drugs, oxytetracycline (adrenal and pituitary tumors), and minocycline (thyroid tumors).

Likewise, although mutagenicity studies of doxycycline have not been conducted, positive results in in vitro mammalian cell assays have been reported for related antibacterial drugs (tetracycline, oxytetracycline).

Doxycycline administered orally at dosage levels as high as 250 mg/kg/day had no apparent effect on the fertility of female rats. Effect on male fertility has not been studied.

Pregnancy

Teratogenic Effects

There are no adequate and well-controlled studies on the use of doxycycline in pregnant women. The vast majority of reported experience with doxycycline during human pregnancy is short-term, first trimester exposure. There are no human data available to assess the effects of long-term therapy of doxycycline in pregnant women, such as that proposed for treatment of anthrax exposure. An expert review of published data on experiences with doxycycline use during pregnancy by TERIS – the Teratogen Information System – concluded that therapeutic doses during pregnancy are unlikely to pose a substantial teratogenic risk (the quantity and quality of data were assessed as limited to fair), but the data are insufficient to state that there is no risk.¹ A case-control study (18,515 mothers of infants with congenital anomalies and 32,804 mothers of infants with no congenital anomalies) shows a weak but marginally statistically significant association with total malformations and use of doxycycline anytime during pregnancy. Sixty-three (0.19%) of the controls and fifty-six (0.30%) of the cases were treated with doxycycline. This association was not seen when the analysis was confined to maternal treatment during the period of organogenesis (i.e., in the second and third months of gestation) with the exception of a marginal relationship with neural tube defect based on only two exposed cases.²

A small prospective study of 81 pregnancies describes 43 pregnant women treated for 10 days with doxycycline during early first trimester. All mothers reported their exposed infants were normal at 1 year of age.³

Nonteratogenic Effects

(See [WARNINGS](#).)

Labor and Delivery

The effect of tetracyclines on labor and delivery is unknown.

Nursing Mothers

Tetracyclines are excreted in human milk; however, the extent of absorption of tetracyclines, including doxycycline, by the breastfed infant is not known. Short-term use by lactating women is not necessarily contraindicated; however, the effects of prolonged exposure to doxycycline in breast milk are unknown.⁴ Because of the potential for serious adverse reactions in nursing infants from doxycycline, a decision should be made whether to discontinue nursing or to discontinue the drug, taking into account the importance of the drug to the mother. (See [WARNINGS](#).)

Pediatric Use

Because of the effects of drugs of the tetracycline-class on tooth development and growth, use doxycycline in pediatric patients 8 years of age or less only when the potential benefits are expected to outweigh the risks in severe or life-threatening conditions (e.g., anthrax, Rocky Mountain spotted fever), particularly when there are no alternative therapies. (See [WARNINGS](#) and [DOSAGE AND ADMINISTRATION](#).)

Adverse Reactions/Side Effects

Due to oral doxycycline's virtually complete absorption, side effects of the lower bowel, particularly diarrhea, have been infrequent. The following adverse reactions have been observed in patients receiving tetracyclines:

Gastrointestinal: anorexia, nausea, vomiting, diarrhea, glossitis, dysphagia, enterocolitis, inflammatory lesions (with monilial overgrowth) in the anogenital region, and pancreatitis. Hepatotoxicity has been reported rarely. These reactions have been caused by both the oral and parenteral administration of tetracyclines. Superficial discoloration of the adult permanent dentition, reversible upon drug discontinuation and professional dental cleaning has been reported. Permanent tooth discoloration and enamel hypoplasia may occur with drugs of the tetracycline class when used during tooth development. (See [WARNINGS](#).) Rare instances of esophagitis and esophageal ulcerations have been reported in patients receiving capsule and tablet forms of the drugs in the tetracycline class. Most of these patients took medications immediately before going to bed. (See [DOSAGE AND ADMINISTRATION](#).)

Skin: toxic epidermal necrolysis, Stevens-Johnson syndrome, erythema multiforme, fixed eruption, skin hyperpigmentation, maculopapular and erythematous rashes. Exfoliative dermatitis has been reported but is uncommon. Photosensitivity is discussed above. (See [WARNINGS](#).)

Renal toxicity: Rise in BUN has been reported and is apparently dose related. (See [WARNINGS](#).)

Immune: Hypersensitivity reactions including urticaria, angioneurotic edema, anaphylaxis, anaphylactoid purpura, serum sickness, pericarditis, exacerbation of systemic lupus erythematosus, drug reaction with eosinophilia and systemic symptoms (DRESS), and Jarisch-Herxheimer reaction has been reported in the setting of spirochete infections treated with doxycycline.

Blood: Hemolytic anemia, thrombocytopenia, neutropenia, and eosinophilia have been reported.

Other: Bulging fontanel in infants and intracranial hypertension in adults. (See [WARNINGS](#).)

When given over prolonged periods, tetracyclines have been reported to produce brown-black microscopic discoloration of the thyroid gland. No abnormalities of thyroid function studies are known to occur.

Overdosage

In case of overdosage, discontinue medication, treat symptomatically and institute supportive measures. Dialysis does

not alter serum half-life and thus would not be of benefit in treating cases of overdosage.

Vibramycin Dosage and Administration

The usual dosage and frequency of administration of doxycycline differs from that of the other tetracyclines. Exceeding the recommended dosage may result in an increased incidence of side effects.

Adults

The usual dose of oral doxycycline is 200 mg on the first day of treatment (administered 100 mg every 12 hours) followed by a maintenance dose of 100 mg/day. In the management of more severe infections (particularly chronic infections of the urinary tract), 100 mg every 12 hours is recommended.

Pediatric Patients

For all pediatric patients weighing less than 45 kg with severe or life-threatening infections (e.g., anthrax, Rocky Mountain spotted fever), the recommended dosage is 2.2 mg/kg of body weight administered every 12 hours. Children weighing 45 kg or more should receive the adult dose. (See [WARNINGS](#) and [PRECAUTIONS](#).)

For pediatric patients with less severe disease (greater than 8 years of age and weighing less than 45 kg), the recommended dosage schedule is 4.4 mg/kg of body weight divided into two doses on the first day of treatment, followed by a maintenance dose of 2.2 mg/kg of body weight (given as a single daily dose or divided into twice daily doses). For pediatric patients weighing over 45 kg, the usual adult dose should be used.

The therapeutic antibacterial serum activity will usually persist for 24 hours following recommended dosage.

When used in streptococcal infections, therapy should be continued for 10 days.

Administration of adequate amounts of fluid along with capsule and tablet forms of drugs in the tetracycline class is recommended to wash down the drugs and reduce the risk of esophageal irritation and ulceration. (See [ADVERSE REACTIONS](#).)

If gastric irritation occurs, it is recommended that doxycycline be given with food or milk. The absorption of doxycycline is not markedly influenced by simultaneous ingestion of food or milk.

Studies to date have indicated that administration of doxycycline at the usual recommended doses does not lead to excessive accumulation of doxycycline in patients with renal impairment.

Uncomplicated gonococcal infections in adults (except anorectal infections in men): 100 mg, by mouth, twice a day for 7 days. As an alternate single visit dose, administer 300 mg stat followed in one hour by a second 300 mg dose. The dose may be administered with food, including milk or carbonated beverage, as required.

Uncomplicated urethral, endocervical, or rectal infection in adults caused by *Chlamydia trachomatis*: 100 mg, by mouth twice a day for 7 days.

Nongonococcal urethritis (NGU) caused by *C. trachomatis* or *U. urealyticum*: 100 mg by mouth, twice a day for 7 days.

Syphilis – early: Patients who are allergic to penicillin should be treated with doxycycline 100 mg, by mouth, twice a day for 2 weeks.

Syphilis of more than one year's duration: Patients who are allergic to penicillin should be treated with doxycycline 100 mg, by mouth, twice a day for 4 weeks.

Acute epididymo-orchitis caused by *N. gonorrhoeae*: 100 mg, by mouth, twice a day for at least 10 days.

Acute epididymo-orchitis caused by *C. trachomatis*: 100 mg, by mouth, twice a day for at least 10 days.

For prophylaxis of malaria: For adults, the recommended dose is 100 mg daily. For children over 8 years of age, the recommended dose is 2 mg/kg given once daily up to the adult dose. Prophylaxis should begin 1–2 days before travel to the malarious area. Prophylaxis should be continued daily during travel in the malarious area and for 4 weeks after the traveler leaves the malarious area.

Inhalational anthrax (post-exposure):

ADULTS: 100 mg of doxycycline, by mouth, twice a day for 60 days.

CHILDREN: weighing less than 45 kg; 2.2 mg/kg of body weight by mouth, twice a day for 60 days. Children weighing 45 kg or more should receive the adult dose.

How is Vibramycin supplied

Product Name	Dosage Form	Product Description	NDC Number and Pack Size
Vibramycin Hyclate (doxycycline hyclate)	Capsules	Aqua blue capsule imprinted with "VIBRA" on one half and "PFIZER 095" on the other half.	NDC 0069-0950-50 100 mg doxycycline bottles of 50
Vibra-Tabs (doxycycline hyclate)	Tablets	Salmon colored film-coated tablets, imprinted on one side with "VIBRA-TABS" and "PFIZER 099" on the other side.	NDC 0069-0990-50 100 mg doxycycline bottles of 50 PRODUCT DISCONTINUED
Vibramycin Calcium (doxycycline calcium)	Syrup-oral suspension	Raspberry-apple flavored oral suspension.	NDC 0069-0971-93 50 mg/5mL: 1 pint (473 mL) bottle or NDC 0069-0971-74 NDC 0069-0971-95 50 mg/5mL: 1 pint (473 mL) bottle (with carton)
Vibramycin Monohydrate (doxycycline monohydrate)	Oral suspension	Raspberry-flavored, dry powder for oral suspension.	NDC 0069-0970-65 25 mg/5mL, 2 oz (60 mL) bottles

All products are to be stored below 86°F (30°C) and dispensed in tight, light-resistant containers (USP).

ANIMAL PHARMACOLOGY AND ANIMAL TOXICOLOGY

Hyperpigmentation of the thyroid has been produced by members of the tetracycline class in the following species: in rats by oxytetracycline, doxycycline, tetracycline PO₄, and methacycline; in minipigs by doxycycline, minocycline, tetracycline PO₄, and methacycline; in dogs by doxycycline and minocycline; in monkeys by minocycline.

Minocycline, tetracycline PO₄, methacycline, doxycycline, tetracycline base, oxytetracycline HCl, and tetracycline HCl were goitrogenic in rats fed a low iodine diet. This goitrogenic effect was accompanied by high radioactive iodine uptake. Administration of minocycline also produced a large goiter with high radioiodine uptake in rats fed a relatively high iodine diet.

Treatment of various animal species with this class of drugs has also resulted in the induction of thyroid hyperplasia in the following: in rats and dogs (minocycline); in chickens (chlortetracycline); and in rats and mice (oxytetracycline). Adrenal gland hyperplasia has been observed in goats and rats treated with oxytetracycline.

References

1. Friedman JM and Polifka JE. Teratogenic Effects of Drugs. A Resource for Clinicians (TERIS). Baltimore, MD: The Johns Hopkins University Press, 2000: 149–195.
2. Cziezel AE and Rockenbauer M. Teratogenic study of doxycycline. Obstet Gynecol 1997; 89: 524–528.
3. Horne HW Jr and Kundsinn RB. The role of mycoplasma among 81 consecutive pregnancies: a prospective study. Int J Fertil 1980; 25: 315–317.
4. Hale T. Medications and Mothers Milk. 9th edition. Amarillo, TX: Pharmasoft Publishing, 2000: 225–226.

This product's labeling may have been updated. For the most recent prescribing information, please visit www.pfizer.com.



LAB-0074-25.0

Revised: 01/2024

PRINCIPAL DISPLAY PANEL - 25 mg Oral Suspension Bottle Label

NDC 0069-0970-65

60 mL when reconstituted

Vibramycin®

(doxycycline monohydrate)

FOR ORAL SUSPENSION

25 mg/5 mL*

RASPBERRY FLAVORED

Pfizer

Pfizer Labs

Division of Pfizer Inc, NY, NY 10017

FOR ORAL USE ONLY.
Store dry powder below 86°F (30°C).
SHAKE WELL BEFORE EACH USE.
MIXING DIRECTIONS
Tap bottle lightly to loosen powder. Add 47.6 mL of water to the bottle to make a total volume of 60 mL. Shake well.

This prescription, when in suspension, will maintain its potency for two weeks when kept at room temperature.

DISCARD UNUSED PORTION AFTER TWO WEEKS.

DOSAGE AND USE
See accompanying prescribing information.

* When reconstituted as directed, each teaspoonful (5 mL) contains doxycycline monohydrate equivalent to 25 mg of doxycycline.

Each bottle contains doxycycline monohydrate equivalent to 300 mg of doxycycline.

6505-00-783-6699

NDC 0069-0970-65


60 mL when reconstituted

Vibramycin®
(doxycycline monohydrate)


FOR ORAL SUSPENSION

25 mg/5 mL *

RASPBERRY FLAVORED



Pfizer Labs
Division of Pfizer Inc, NY, NY 10017




FOR POSITION ONLY
Specs: UPC 100% x 8mm
3 N 0069-0970-65 3

6277

Rx only

MADE IN USA

62103500



PRINCIPAL DISPLAY PANEL - 25 mg/5 mL Bottle Carton

NDC 0069-0970-65

60 mL when reconstituted

Vibramycin®

(doxycycline monohydrate)

FOR ORAL SUSPENSION


25 mg/5 mL*

RASPBERRY FLAVORED

Pfizer

Pfizer Labs

Division of Pfizer Inc, NY, NY 10017



PRINCIPAL DISPLAY PANEL - 473 mL Bottle Label

NDC 0069-0971-95

Pfizer

Vibramycin®

calcium

(doxycycline calcium)

syrup

50 mg/5 mL*

ORAL SUSPENSION

FOR ORAL USE ONLY

RASPBERRY/APPLE FLAVORED

SHAKE WELL BEFORE USING

1 Pint (473 mL)

Rx only



NDC 0069-0971-95

Vibramycin[®]
calcium
(doxycycline calcium)
syrup

50 mg/5 mL*

**ORAL SUSPENSION
FOR ORAL USE ONLY**

RASPBERRY/APPLE FLAVORED

SHAKE WELL BEFORE USING

1 Pint (473 mL)

Rx only

Store below 86°F (30°C).
Dispense in tight (USP), light-resistant containers.

DOSAGE AND USE
See accompanying prescribing information.

USUAL DOSAGE
Adults: 200 mg on the first day (100 mg every 12 hours) followed by a maintenance dose of 100 mg a day.
Children above eight years of age:
Under 100 lbs. — 2 mg/lb. of body weight daily divided in two doses on the first day, followed by 1 mg/lb. of body weight on subsequent days in one or two doses.
Over 100 lbs. — See adult dosage.
doxycycline U.S. Pat. No. 3,200,149
* Each teaspoonful (5 mL) contains doxycycline calcium equivalent to 50 mg of doxycycline.

Distributed by
Pfizer Labs
Division of Pfizer Inc
NY, NY 10017

FPO GS1 Data Bar Limited - 10 mil

(01) 10300690971954

PA A 1 3 8 5 3 4
LOT/EXP:

Imprint Area
reads this way
NO VARNISH
NO COPY
NO INK

PRINCIPAL DISPLAY PANEL - 473 mL Bottle Carton

Pfizer

NDC 0069-0971-95

Vibramycin®

calcium

(doxycycline calcium)

syrup

50 mg/5 mL*

ORAL SUSPENSION

FOR ORAL USE ONLY

IMPORTANT

THIS CLOSURE IS NOT

CHILD-RESISTANT.

RASPBERRY/APPLE FLAVORED

SHAKE WELL BEFORE USING

1 Pint (473 mL)

Rx only



PRINCIPAL DISPLAY PANEL - 473 mL Bottle Label - NDC 0069-0971-74

NDC 0069-0971-74

Pfizer

Vibramycin®
calcium
(doxycycline calcium)

syrup

50 mg/5 mL*

ORAL SUSPENSION
FOR ORAL USE ONLY

RASPBERRY/APPLE FLAVORED

SHAKE WELL BEFORE USING

1 Pint (473 mL)

Rx only



NDC 0069-0971-74

Vibramycin®
calcium
(doxycycline calcium)
syrup

50 mg/5 mL*

ORAL SUSPENSION
FOR ORAL USE ONLY

RASPBERRY/APPLE FLAVORED

SHAKE WELL BEFORE USING

1 Pint (473 mL)

Rx only

Store below 86°F (30°C).
Dispense in tight (USP), light-resistant containers.
DOSAGE AND USE
See accompanying prescribing information.
USUAL DOSAGE
Adults: 200 mg on the first day (100 mg every 12 hours) followed by a maintenance dose of 100 mg a day.
Children above eight years of age:
Under 100 lbs. — 2 mg/lb. of body weight daily divided in two doses on the first day, followed by 1 mg/lb. of body weight on subsequent days in one or two doses.
Over 100 lbs. — See adult dosage.
doxycycline U.S. Pat. No. 3,200,149
* Each teaspoonful (5 mL) contains doxycycline calcium equivalent to 50 mg of doxycycline.

Distributed by
Pfizer Labs
Division of Pfizer Inc
NY, NY 10017

FPO GST Data Bar Limited - 10 mil
(01)10300690971749

PAA165104
LOT/EXP:

Imprint Area
reads this way
NO VARNISH
NO COPY
NO INK

PRINCIPAL DISPLAY PANEL - 473 mL Bottle Carton - NDC 0069-0971-74

Pfizer

NDC 0069-0971-74

Vibramycin®
calcium
(doxycycline calcium)
syrup

50 mg/5 mL*

**ORAL SUSPENSION
FOR ORAL USE ONLY**

IMPORTANT

THIS CLOSURE IS NOT
CHILD-RESISTANT.

RASPBERRY/APPLE FLAVORED

SHAKE WELL BEFORE USING

1 Pint (473 mL)

Rx only



PRINCIPAL DISPLAY PANEL - 100 mg Capsule Bottle Label

Pfizer

NDC 0069-0950-50

Vibramycin®
 (doxycycline hyclate)

100 mg*

50 Capsules


Rx only

Store below 86°F (30°C).
Dispense in tight, light-resistant containers (USP).

DOSAGE AND USE
See accompanying prescribing information.

*Each capsule contains doxycycline hyclate equivalent to 100 mg of doxycycline.


Distributed by
Pfizer Labs
Division of Pfizer Inc, NY, NY 10017

 NDC 0069-0950-50

Vibramycin®
(doxycycline hyclate)

100 mg*

50 Capsules Rx only


FPO: UPC @ 80%
3 0069-0950-50 1

14467001

GTIN: 00300690950501
LOT: /EXP:

PRINCIPAL DISPLAY PANEL - 100 mg Tablet Bottle Label

NDC 0069-0990-50

50 Tablets

Vibra-Tabs®

(doxycycline hyclate)

100 mg*

Pfizer

Distributed by

Pfizer Labs

Division of Pfizer Inc, NY, NY 10017

Store below 86°F (30°C).
Dispense in tight, light-resistant containers (USP).


DOSAGE AND USE
See accompanying prescribing information.

*Each tablet contains doxycycline hyclate equivalent to 100 mg doxycycline.


MADE IN USA


Rx only

50 Tablets

Vibra-Tabs® 
(doxycycline hyclate)

100 mg*

Distributed by
 **Pfizer Labs**
Division of Pfizer Inc, NY, NY 10017


FPO UPC 80% x 4 mm
N 3 0069-0990-50 7

05-3193-31-5

6081

VIBRAMYCIN MONOHYDRATE
doxycycline powder, for suspension

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0069-0970
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DOXYCYCLINE (UNII: N12000U13O) (DOXYCYCLINE ANHYDROUS - UNII:334895S862)	DOXYCYCLINE ANHYDROUS	25 mg in 5 mL

Inactive Ingredients

Ingredient Name	Strength
CARBOXYMETHYLCELLULOSE SODIUM, UNSPECIFIED (UNII: K679OBS311)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
METHYLPARABEN (UNII: A2I8C7HI9T)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
PROPYLPARABEN (UNII: Z8IX2SC1OH)	
D&C RED NO. 28 (UNII: 767IP0Y5NH)	
SUCROSE (UNII: C151H8M554)	

Product Characteristics

Color	Score
Shape	Size
Flavor	Imprint Code
Contains	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0069-0970-65	1 in 1 CARTON	12/06/1967	06/24/2017
1		60 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA050006	12/06/1967	06/24/2017

VIBRAMYCIN CALCIUM
doxycycline calcium syrup

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0069-0971
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DOXYCYCLINE CALCIUM (UNII: 8ZL07I20SB) (DOXYCYCLINE ANHYDROUS - UNII:334895S862)	DOXYCYCLINE ANHYDROUS	50 mg in 5 mL

Inactive Ingredients

Ingredient Name	Strength
CARBOXYMETHYLCELLULOSE SODIUM, UNSPECIFIED (UNII: K679OBS311)	
BUTYLPARABEN (UNII: 3QPI1U3FV8)	
CALCIUM CHLORIDE (UNII: M4I0D6VV5M)	
GLYCERIN (UNII: PDC6A3C0OX)	
HYDROCHLORIC ACID (UNII: QTT17582CB)	
MAGNESIUM ALUMINUM SILICATE (UNII: 6M3P64V0NC)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
PROPYLPARABEN (UNII: Z8IX2SC1OH)	
SODIUM HYDROXIDE (UNII: 55X04QC32I)	
SODIUM METABISULFITE (UNII: 4VON5FNS3C)	
SORBITOL (UNII: 506T60A25R)	
WATER (UNII: 059QF0KO0R)	

Product Characteristics

Color	Score
Shape	Size
Flavor	Imprint Code
Contains	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0069-0971-93	473 mL in 1 BOTTLE; Type 0: Not a Combination Product	09/23/1974	11/30/2021
2	NDC:0069-0971-95	1 in 1 CARTON	11/01/2019	08/31/2023
2		473 mL in 1 BOTTLE; Type 0: Not a Combination Product		
3	NDC:0069-0971-74	1 in 1 CARTON	08/01/2020	08/01/2020

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA050480	09/23/1974	08/31/2023

VIBRAMYCIN HYCLATE
doxycycline hyclate capsule

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0069-0950
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DOXYCYCLINE HYCLATE (UNII: 19XTS3T51U) (DOXYCYCLINE ANHYDROUS - UNII:334895S862)	DOXYCYCLINE ANHYDROUS	100 mg

Inactive Ingredients

Ingredient Name	Strength
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	

Product Characteristics

Color	BLUE (Aqua Blue)	Score	no score
Shape	CAPSULE	Size	22mm
Flavor		Imprint Code	Vibra;Pfizer;095

Contains

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0069-0950-50	50 in 1 BOTTLE; Type 0: Not a Combination Product	12/05/1967	11/30/2024

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA050007	12/05/1967	11/30/2024

VIBRA-TABS
doxycycline hyclate tablet, film coated

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0069-0990
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DOXYCYCLINE HYCLATE (UNII: 19XTS3T51U) (DOXYCYCLINE ANHYDROUS - UNII:334895S862)	DOXYCYCLINE ANHYDROUS	100 mg

Inactive Ingredients

Ingredient Name	Strength
ETHYLCELLULOSE, UNSPECIFIED (UNII: 7Z8S9VYZ4B)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
ALUMINUM OXIDE (UNII: LMI26O6933)	

Product Characteristics

Color	ORANGE	Score	no score
Shape	ROUND	Size	9mm
Flavor		Imprint Code	Vibra;Pfizer;099

Contains

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0069-0990-50	50 in 1 BOTTLE; Type 0: Not a Combination Product	01/15/1980	10/01/2010

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA050533	01/15/1980	10/01/2010

Labeler - Pfizer Laboratories Div Pfizer Inc (134489525)

Establishment

Name	Address	ID/FEI	Business Operations
Pharmacia & Upjohn Company LLC		618054084	ANALYSIS(0069-0971) , MANUFACTURE(0069-0971) , PACK(0069-0971) , LABEL(0069-0971)

Establishment

Name	Address	ID/FEI	Business Operations
Pfizer Ireland Pharmaceuticals		985052076	ANALYSIS(0069-0950, 0069-0970, 0069-0971, 0069-0990) , API MANUFACTURE(0069-0950, 0069-0970, 0069-0971, 0069-0990)

Establishment

Name	Address	ID/FEI	Business Operations
Pfizer Laboratories Div Pfizer Inc		001147495	ANALYSIS(0069-0970, 0069-0990)

Establishment

Name	Address	ID/FEI	Business Operations
Pfizer Inc		943955690	ANALYSIS(0069-0970, 0069-0990, 0069-0971)

Frequently asked questions

- [Does doxycycline work for acne?](#)
- [What are the most common skin conditions? \(with photos\)](#)
- [Is Doxycycline safe for dogs? Uses, dosage, side effects](#)
- [Should doxycycline be taken with food?](#)
- [Doxycycline for STDs: Effectiveness, Dosage and Side Effects](#)
- [What are the best antibiotics for pneumonia?](#)
- [What is the best antibiotic to treat a sinus infection?](#)
- [Does Doxycycline make you tired?](#)

More about Vibramycin (doxycycline)

- [Check interactions](#)
- [Compare alternatives](#)
- [Reviews \(9\)](#)
- [Drug images](#)
- [Side effects](#)

- [Dosage information](#)
- [During pregnancy](#)
- [Drug class: miscellaneous antimalarials](#)
- [Breastfeeding](#)

Patient resources

- [Vibramycin Calcium advanced reading](#)
- [Vibramycin \(Doxycycline Syrup\)](#)
- [Vibramycin \(Doxycycline Tablets and Capsules\)](#)

Professional resources

- [Doxycycline Hyclate \(EENT\) monograph](#)
- [Doxycycline Suspension \(FDA\)](#)

Other brands

[Monodox](#), [Doryx](#), [Oracea](#), [Doxy 100](#), ... +6 more

Related treatment guides

- [Gonococcal Infection, Uncomplicated](#)
- [Acne](#)
- [Actinomycosis](#)
- [Periodontitis](#)

[Medical Disclaimer](#)

Antibiotics 101

[List of Common Antibiotics & Types](#)

[Antibiotics and Alcohol](#)

[Probiotics and Antibiotics](#)

[Antibiotics for Strep Throat](#)

[Antibiotics for a Tooth Infection](#)

[Antibiotics for UTI's](#)

DRUG STATUS

Availability

Rx Prescription only

Pregnancy & Lactation

no Risk data available



CSA Schedule*

N/A Not a controlled drug



Approval History



Drug history at FDA



User Reviews & Ratings

7.0 / 10

[9 Reviews](#)

Related News

[Rare Hair Loss Condition Can Be Treated With Low Doses Of Antibiotic Doxycycline](#)

[DoxyPEP Lowers Rate of Sexually Transmitted Infections \(STIs\), Data Suggests](#)

Images

[Vibramycin 100 mg \(VIBRA PFIZER 095\)](#)



[View larger images](#)

Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.



About

[About Drugs.com](#)

[Advertising policy](#)

[Attribution & citations](#)

Terms & privacy

[Terms of use](#)

[Editorial policy](#)

[Privacy policy](#)

Support

[Help center](#)

[Sitemap](#)

Contact us



[Subscribe to our newsletter](#) for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.



Ad Choices
ADVERTISEMENT



Copyright © 2000-2025 Drugs.com. All rights reserved.