

[Home](#)

2. Xopenex HFA

Xopenex HFA

Generic name: [levalbuterol inhalation](#) [*leh-val-BYOO-ter-all*]**Brand names:** [Xopenex](#), Xopenex HFA, [Xopenex Concentrate](#)**Drug class:** [Adrenergic bronchodilators](#)[Medically reviewed](#) by Drugs.com on Aug 11, 2023. Written by [Cerner Multum](#).[Uses](#) [Side effects](#) [Warnings](#) [Before taking](#) [Dosage](#) [Interactions](#) [FAQ](#)

What is Xopenex HFA?

[Levalbuterol](#) is a short-acting bronchodilator that relaxes muscles in the airways and increases air flow to the lungs.

Xopenex HFA is used to treat or prevent asthma attacks in adults and children who are at least 4 years old.

Xopenex HFA may also be used for purposes not listed in this medication guide.

Xopenex HFA side effects

Get emergency medical help if you have **signs of an allergic reaction**: hives; difficult breathing; swelling of your face, lips, tongue, or throat.

Xopenex HFA may cause serious side effects. Call your doctor at once if you have:

- [wheezing](#), choking, or other breathing problems after using Xopenex HFA;
- pounding heartbeats or fluttering in your chest;
- worsening asthma symptoms; or
- **low potassium**--leg cramps, [constipation](#), irregular heartbeats, fluttering in your chest, extreme thirst, increased urination, numbness or tingling, muscle weakness or limp feeling.

Common side effects of Xopenex HFA may include:

- dizziness, nervousness, tremors;
- runny nose, [sore throat](#);
- chest pain or tightness, irregular heartbeats;
- pain; or
- vomiting.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

 [Xopenex HFA side effects](#) (more detail)

Related/similar drugs

Breztri Aerosphere

Breztri

(budesonide/glycopyrrolate/formoterol fumarate) is a combination inhaler that may be used ...

Reviews & ratings

6.9 / 10

[119 Reviews](#)

[View more](#)

Xolair

Xolair injection (omalizumab) is used to reduce the risk of severe food allergy reactions ...

Reviews & ratings

7.5 / 10

[229 Reviews](#)

[View more](#)

FEATURED

Dupixent

Dupixent may be used to treat eczema, eosinophilic or oral-corticosteroid-dependent asthma, chronic ...

Reviews & ratings

7.2 / 10

[499 Reviews](#)

[View more](#)

Symbicort

Symbicort (budesonide and formoterol) is used to prevent bronchospasm in people with asthma or

or ...

Reviews & ratings

6.6 / 10

[226 Reviews](#)

[View more](#)

Ventolin HFA

Ventolin HFA (albuterol) is used to treat or prevent breathing problems in patients who have asthma ...

Reviews & ratings

7.5 / 10

[18 Reviews](#)

[View more](#)

Epinephrine

Epinephrine (adrenaline) Injection is used to treat severe allergic reactions (anaphylaxis) to ...

Reviews & ratings

7.4 / 10

[59 Reviews](#)

[View more](#)

Dexamethasone

Dexamethasone is used to treat inflammatory conditions such as allergies, skin conditions ...

Reviews & ratings

5.9 / 10

[162 Reviews](#)

[View more](#)

Warnings

Follow all directions on your medicine label and package. Tell each of your healthcare providers about all your medical conditions, allergies, and all medicines you use.

Seek medical attention if you think your asthma medications are not working as well.

Before taking this medicine

You should not use Xopenex HFA if you are allergic to levalbuterol or [albuterol](#) (Accuneb, ProAir, [Proventil](#), [Ventolin](#)).

To make sure levalbuterol is safe for you, tell your doctor if you have:

- [heart disease](#), [high blood pressure](#), or congestive [heart failure](#);
- a seizure disorder;
- diabetes; or
- a thyroid disorder.

It is not known whether this medicine will harm an unborn baby. Tell your doctor if you are pregnant or plan to become pregnant.

It is not known whether levalbuterol inhalation passes into breast milk or if it could affect the nursing baby. Tell your doctor if you are breast-feeding.

A levalbuterol **inhaler** should not be given to a child younger than 4 years old. Levalbuterol solution in a **nebulizer** should not be given to a child younger than 6 years old.

How should I use Xopenex HFA?

Follow all directions on your prescription label. Do not use this medicine in larger or smaller amounts or for longer than recommended.

Read all patient information, medication guides, and instruction sheets provided to you. Ask your doctor or pharmacist if you have any questions.

Any child using Xopenex HFA should be supervised by an adult while using this medicine.

Prime the **inhaler** device before your first use. Pump 4 test sprays into the air, away from your face. Shake the inhaler for at least 5 seconds before each spray. Prime again whenever the inhaler has not been used in longer than 3 days.

To use the **inhaler**:

- **Shake the canister well just before each spray.**
- Uncap the mouthpiece of the inhaler. Breathe out fully. Put the mouthpiece into your mouth and close your lips. Breathe in slowly while pushing down on the canister. Hold your breath for 10 seconds, then breathe out slowly.
- If you use more than one inhalation at a time, wait at least 1 minute before using the second inhalation and shake the inhaler again.
- Keep your inhaler clean and dry, and store it with the cap on the mouthpiece. Clean your inhaler once a week by removing the canister and placing the mouthpiece under warm running water for at least 30 seconds. Shake out the excess water and allow the parts to air dry completely before putting the inhaler back together.

Store the levalbuterol **inhaler** with the mouthpiece down. Keep the inhaler at room temperature away from heat, sunlight, or freezing temperatures. If your inhaler has a dose number indicator on it, throw away the inhaler when the number

displays "0" to show that all sprays have been used.

Keep the inhaler canister away from open flame or high heat, such as in a car on a hot day. The canister may explode if it gets too hot. Do not puncture or burn an empty inhaler canister.

Levalbuterol concentrate solution is given with a nebulizer. The concentrate must be mixed with a liquid (diluent) before using it. Ask your pharmacist about which diluent to use and where you can buy it. Be sure you understand how to properly mix the medicine and place it into the nebulizer.

To use the solution with a **nebulizer**:

- Open the foil pouch and empty the medicine into the chamber of the nebulizer. Add the correct amount of diluent as directed by your doctor.
- Attach the mouthpiece or face mask, then attach the drug chamber to the compressor.
- Sit upright in a comfortable position. Place the mouthpiece into your mouth or put on the face mask, covering your nose and mouth. Turn on the compressor.
- Breathe in slowly and evenly until no more mist is formed by the nebulizer and the drug chamber is empty.
- Clean the nebulizer after each use. Follow the cleaning directions that came with your nebulizer.

Do not use the nebulizer solution if it does not appear clear and colorless.

Store the levalbuterol **concentrate** vials in the protective foil pouch at room temperature, away from moisture, heat, and light. Keep each vial in the foil pouch until you are ready to prepare a dose in the nebulizer.

Each single-use plastic vial of levalbuterol concentrate is for one use only. Throw away after one use, even if there is still some medicine left in it after inhaling your dose.

Asthma is often treated with a combination of drugs. Use all medications as directed by your doctor. **Do not change your doses or medication schedule without your doctor's advice.**

Using too much levalbuterol or using it too often can cause life-threatening side effects. **Seek medical attention if you think your asthma medications are not working as well.** An increased need for medication could be an early sign of a serious asthma attack.

 [Detailed Xopenex HFA dosage information](#)

What happens if I miss a dose?

Use the missed dose as soon as you remember. Skip the missed dose if it is almost time for your next scheduled dose.

Do not use extra medicine to make up the missed dose.

Use Xopenex HFA regularly to get the most benefit. Get your prescription refilled before you run out of medicine completely.

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222. **An overdose of levalbuterol can be fatal.**

What should I avoid while using Xopenex HFA?

Avoid situations that may make your condition worse such as exercising in cold, dry air; smoking; breathing in dust; and exposure to allergens such as pet fur.

What other drugs will affect Xopenex HFA?

Tell your doctor if you have used an MAO inhibitor in the past 14 days. MAO inhibitors include [isocarboxazid](#), [linezolid](#), [methylene blue](#) injection, [phenelzine](#), [rasagiline](#), [selegiline](#), [tranylcypromine](#), and others.

Tell your doctor about all your current medicines and any you start or stop using, especially:

- any other inhaled medicines to treat asthma or COPD (chronic obstructive pulmonary disease);
- an antidepressant;
- a diuretic or "water pill"; or
- medicine to treat a heart condition.

This list is not complete. Other drugs may interact with levalbuterol, including prescription and over-the-counter medicines, vitamins, and [herbal products](#). Not all possible interactions are listed in this medication guide.

 [Xopenex HFA drug interactions](#) (more detail)

Does Xopenex HFA interact with my other drugs?

Enter medications to view a detailed interaction report using our [Drug Interaction Checker](#).

Xopenex HFA

+

Enter a drug name

Add

Popular FAQ

Can asthma inhalers cause a sore throat?



More about Xopenex HFA (levalbuterol)

- [Check interactions](#)
- [Compare alternatives](#)
- [Pricing & coupons](#)
- [Reviews \(5\)](#)
- [Side effects](#)

- [Dosage information](#)
- [During pregnancy](#)
- [Drug class: adrenergic bronchodilators](#)
- [Breastfeeding](#)
- [En español](#)

Patient resources

- [Xopenex HFA advanced reading](#)

Professional resources

- [Xopenex HFA prescribing information](#)

Other formulations

- [Xopenex](#)
- [Xopenex Concentrate](#)

Related treatment guides

- [COPD, Maintenance](#)
- [COPD](#)
- [Asthma](#)
- [Asthma, Maintenance](#)
- [Asthma, acute](#)
- [COPD, Acute](#)

Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use this medication only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

[Medical Disclaimer](#)

Copyright 1996-2025 Cerner Multum, Inc. Version: 10.01.

DRUG STATUS

Availability

Rx Prescription only

Pregnancy & Lactation



Risk data available

CSA Schedule*

N/A Not a controlled drug

Approval History



Drug history at FDA



User Reviews & Ratings

[5 Reviews](#)

Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.



About

[About Drugs.com](#)

[Advertising policy](#)

[Attribution & citations](#)

Terms & privacy

[Terms of use](#)

[Editorial policy](#)

[Privacy policy](#)

Support

[Help center](#)

[Sitemap](#)

[Contact us](#)



[Subscribe to our newsletter](#) for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.



Ad Choices
ADVERTISEMENT



Copyright © 2000-2025 Drugs.com. All rights reserved.