



[Home](#)

2. [FDA PI](#)

3. [Vicodin ES](#)

Vicodin ES: Package Insert / Prescribing Info

Package insert / product label

Generic name: [acetaminophen and hydrocodone bitartrate](#)

Dosage form: tablet

Drug class: [Narcotic analgesic combinations](#)

[Medically reviewed](#) by Drugs.com. Last updated on Mar 25, 2025.



The Vicodin ES brand name has been **discontinued** in the U.S. If generic versions of this product have been approved by the FDA, there may be [generic equivalents available](#).

On This Page

[Warnings](#)

[Description](#)

[Clinical Pharmacology](#)

[Indications and Usage](#)

[Contraindications](#)

[Precautions](#)

[Patient Counseling Information](#)

[Drug Interactions](#)

[Adverse Reactions/Side Effects](#)

[Drug Abuse and Dependence](#)

[Overdosage](#)

[Dosage and Administration](#)

[How Supplied/Storage and Handling](#)

HEPATOTOXICITY

ACETAMINOPHEN HAS BEEN ASSOCIATED WITH CASES OF ACUTE LIVER FAILURE, AT TIMES RESULTING IN LIVER TRANSPLANT AND DEATH. MOST OF THE CASES OF LIVER INJURY ARE ASSOCIATED WITH THE USE OF ACETAMINOPHEN AT DOSES THAT EXCEED 4000 MILLIGRAMS PER DAY, AND OFTEN INVOLVE MORE THAN ONE ACETAMINOPHEN-CONTAINING PRODUCT.


Warnings

Vicodin ES Description


Hydrocodone bitartrate and acetaminophen is supplied in tablet form for oral administration.

WARNING: May be habit-forming (see **PRECAUTIONS**, [Information for Patients/Caregivers](#), and [DRUG ABUSE AND DEPENDENCE](#)).

Hydrocodone bitartrate is an opioid analgesic and antitussive and occurs as fine, white crystals or as a crystalline powder. It is affected by light. The chemical name is 4,5 α -epoxy-3-methoxy-17-methylmorphinan-6-one tartrate (1:1) hydrate (2:5). It has the following structural formula:



Acetaminophen, 4'-hydroxyacetanilide, a slightly bitter, white, odorless, crystalline powder, is a non-opiate, non-salicylate analgesic and antipyretic. It has the following structural formula:



Hydrocodone Bitartrate and Acetaminophen Tablets, USP is available in the following strengths:

VICODIN[®]: Hydrocodone Bitartrate..... 5 mg

WARNING: May be habit-forming.

Acetaminophen..... 300 mg

VICODIN ES[®]: Hydrocodone Bitartrate..... 7.5 mg

WARNING: May be habit-forming.

Acetaminophen..... 300 mg

VICODIN HP[®]: Hydrocodone Bitartrate..... 10 mg

WARNING: May be habit-forming.

Acetaminophen..... 300 mg

In addition each tablet contains the following inactive ingredients: colloidal silicon dioxide, crospovidone, magnesium stearate, microcrystalline cellulose, povidone, pregelatinized starch, and stearic acid.

This product complies with USP dissolution test 2.

Vicodin ES - Clinical Pharmacology

Hydrocodone is a semisynthetic narcotic analgesic and antitussive with multiple actions qualitatively similar to those of codeine. Most of these involve the central nervous system and smooth muscle. The precise mechanism of action of hydrocodone and other opiates is not known, although it is believed to relate to the existence of opiate receptors in the central nervous system. In addition to analgesia, narcotics may produce drowsiness, changes in mood and mental clouding.

The analgesic action of acetaminophen involves peripheral influences, but the specific mechanism is as yet undetermined. Antipyretic activity is mediated through hypothalamic heat regulating centers. Acetaminophen inhibits prostaglandin synthetase. Therapeutic doses of acetaminophen have negligible effects on the cardiovascular or respiratory systems; however, toxic doses may cause circulatory failure and rapid, shallow breathing.

Pharmacokinetics

The behavior of the individual components is described below.

Hydrocodone

Following a 10 mg oral dose of hydrocodone administered to five adult male subjects, the mean peak concentration was 23.6 ± 5.2 ng/mL. Maximum serum levels were achieved at 1.3 ± 0.3 hours and the half-life was determined to be 3.8 ± 0.3 hours. Hydrocodone exhibits a complex pattern of metabolism including O-demethylation, N-demethylation and 6-keto reduction to the corresponding 6- α - and 6- β -hydroxy- metabolites. See **OVERDOSAGE** for toxicity information.

Acetaminophen

Acetaminophen is rapidly absorbed from the gastrointestinal tract and is distributed throughout most body tissues. The plasma half-life is 1.25 to 3 hours, but may be increased by liver damage and following overdosage. Elimination of acetaminophen is principally by liver metabolism (conjugation) and subsequent renal excretion of metabolites. Approximately 85% of an oral dose appears in the urine within 24 hours of administration, most as the glucuronide conjugate, with small amounts of other conjugates and unchanged drug. See **OVERDOSAGE** for toxicity information.

Indications and Usage for Vicodin ES

Hydrocodone bitartrate and acetaminophen tablets are indicated for the relief of moderate to moderately severe pain.

Contraindications

This product should not be administered to patients who have previously exhibited hypersensitivity to hydrocodone or acetaminophen.

Patients known to be hypersensitive to other opioids may exhibit cross sensitivity to hydrocodone.

Warnings

Hepatotoxicity

Acetaminophen has been associated with cases of acute liver failure, at times resulting in liver transplant and death. Most of the cases of liver injury are associated with the use of acetaminophen at doses that exceed 4000 milligrams per day,

and often involve more than one acetaminophen-containing product. The excessive intake of acetaminophen may be intentional to cause self-harm or unintentional as patients attempt to obtain more pain relief or unknowingly take other acetaminophen-containing products.

The risk of acute liver failure is higher in individuals with underlying liver disease and in individuals who ingest alcohol while taking acetaminophen.

Instruct patients to look for acetaminophen or APAP on package labels and not to use more than one product that contains acetaminophen. Instruct patients to seek medical attention immediately upon ingestion of more than 4000 milligrams of acetaminophen per day, even if they feel well.

Serious skin reactions

Rarely, acetaminophen may cause serious skin reactions such as acute generalized exanthematous pustulosis (AGEP), Stevens-Johnson Syndrome (SJS), and toxic epidermal necrolysis (TEN), which can be fatal. Patients should be informed about the signs of serious skin reactions, and use of the drug should be discontinued at the first appearance of skin rash or any other sign of hypersensitivity.

Hypersensitivity/anaphylaxis

There have been post-marketing reports of hypersensitivity and anaphylaxis associated with use of acetaminophen. Clinical signs included swelling of the face, mouth and throat, respiratory distress, urticaria, rash, pruritus, and vomiting. There were infrequent reports of life-threatening anaphylaxis requiring emergency medical attention. Instruct patients to discontinue hydrocodone bitartrate and acetaminophen tablets immediately and seek medical care if they experience these symptoms. Do not prescribe hydrocodone bitartrate and acetaminophen tablets for patients with acetaminophen allergy.

Respiratory Depression

At high doses or in sensitive patients, hydrocodone may produce dose-related respiratory depression by acting directly on the brain stem respiratory center. Hydrocodone also affects the center that controls respiratory rhythm, and may produce irregular and periodic breathing.

Head Injury and Increased Intracranial Pressure

The respiratory depressant effects of narcotics and their capacity to elevate cerebrospinal fluid pressure may be markedly exaggerated in the presence of head injury, other intracranial lesions or a preexisting increase in intracranial pressure. Furthermore, narcotics produce adverse reactions which may obscure the clinical course of patients with head injuries.

Acute Abdominal Conditions

The administration of narcotics may obscure the diagnosis or clinical course of patients with acute abdominal conditions.

Precautions

General

Special Risk Patients

As with any narcotic analgesic agent, hydrocodone bitartrate and acetaminophen tablets should be used with caution in elderly or debilitated patients and those with severe impairment of hepatic or renal function, hypothyroidism, Addison's disease, prostatic hypertrophy or urethral stricture. The usual precautions should be observed and the possibility of respiratory depression should be kept in mind.

Cough Reflex

Hydrocodone suppresses the cough reflex; as with all narcotics, caution should be exercised when hydrocodone bitartrate and acetaminophen tablets are used postoperatively and in patients with pulmonary disease.

Information for Patients/Caregivers

- Do not take hydrocodone bitartrate and acetaminophen tablets if you are allergic to any of its ingredients.
- If you develop signs of allergy such as a rash or difficulty breathing stop taking hydrocodone bitartrate and acetaminophen tablets and contact your healthcare provider immediately.
- Do not take more than 4000 milligrams of acetaminophen per day. Call your doctor if you took more than the recommended dose.

Hydrocodone, like all narcotics, may impair the mental and/or physical abilities required for the performance of potentially hazardous tasks such as driving a car or operating machinery; patients should be cautioned accordingly.

Alcohol and other CNS depressants may produce an additive CNS depression, when taken with this combination product, and should be avoided.

Hydrocodone may be habit forming. Patients should take the drug only for as long as it is prescribed, in the amounts prescribed, and no more frequently than prescribed.

Laboratory Tests

In patients with severe hepatic or renal disease, effects of therapy should be monitored with serial liver and/or renal function tests.

Drug Interactions

Patients receiving other narcotics, antihistamines, antipsychotics, antianxiety agents, or other CNS depressants (including alcohol) concomitantly with hydrocodone bitartrate and acetaminophen tablets may exhibit an additive CNS depression. When combined therapy is contemplated, the dose of one or both agents should be reduced.

The use of MAO inhibitors or tricyclic antidepressants with hydrocodone preparations may increase the effect of either the antidepressant or hydrocodone.

Drug/Laboratory Test Interactions

Acetaminophen may produce false-positive test results for urinary 5-hydroxyindoleacetic acid.

Carcinogenesis, Mutagenesis, Impairment of Fertility

No adequate studies have been conducted in animals to determine whether hydrocodone or acetaminophen have a potential for carcinogenesis, mutagenesis, or impairment of fertility.

Pregnancy

Teratogenic Effects

Pregnancy Category C

There are no adequate and well-controlled studies in pregnant women. Hydrocodone bitartrate and acetaminophen tablets should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus.

Nonteratogenic Effects

Babies born to mothers who have been taking opioids regularly prior to delivery will be physically dependent. The withdrawal signs include irritability and excessive crying, tremors, hyperactive reflexes, increased respiratory rate, increased stools, sneezing, yawning, vomiting, and fever. The intensity of the syndrome does not always correlate with the duration of maternal opioid use or dose. There is no consensus on the best method of managing withdrawal.

Labor and Delivery

As with all narcotics, administration of this product to the mother shortly before delivery may result in some degree of respiratory depression in the newborn, especially if higher doses are used.

Nursing Mothers

Acetaminophen is excreted in breast milk in small amounts, but the significance of its effects on nursing infants is not known. It is not known whether hydrocodone is excreted in human milk. Because many drugs are excreted in human milk and because of the potential for serious adverse reactions in nursing infants from hydrocodone and acetaminophen, a decision should be made whether to discontinue nursing or to discontinue the drug, taking into account the importance of the drug to the mother.

Pediatric Use

Safety and effectiveness in pediatric patients have not been established.

Geriatric Use

Clinical studies of hydrocodone bitartrate and acetaminophen tablets did not include sufficient numbers of subjects aged 65 and over to determine whether they respond differently from younger subjects. Other reported clinical experience has not identified differences in responses between the elderly and younger patients. In general, dose selection for an elderly patient should be cautious, usually starting at the low end of the dosing range, reflecting the greater frequency of decreased hepatic, renal, or cardiac function, and of concomitant disease or other drug therapy.

Hydrocodone and the major metabolites of acetaminophen are known to be substantially excreted by the kidney. Thus the risk of toxic reactions may be greater in patients with impaired renal function due to accumulation of the parent compound and/or metabolites in the plasma. Because elderly patients are more likely to have decreased renal function, care should be taken in dose selection, and it may be useful to monitor renal function.

Hydrocodone may cause confusion and over-sedation in the elderly; elderly patients generally should be started on low doses of hydrocodone bitartrate and acetaminophen tablets and observed closely.

Adverse Reactions/Side Effects

The most frequently reported adverse reactions are lightheadedness, dizziness, sedation, nausea and vomiting. These effects seem to be more prominent in ambulatory than in nonambulatory patients, and some of these adverse reactions may be alleviated if the patient lies down.

Other adverse reactions include:

Central Nervous System

Drowsiness, mental clouding, lethargy, impairment of mental and physical performance, anxiety, fear, dysphoria, psychic dependence, mood changes.

Gastrointestinal System

Prolonged administration of hydrocodone bitartrate and acetaminophen tablets may produce constipation.

Genitourinary System

Ureteral spasm, spasm of vesical sphincters and urinary retention have been reported with opiates.

Respiratory Depression

Hydrocodone bitartrate may produce dose-related respiratory depression by acting directly on the brain stem respiratory centers (see [OVERDOSAGE](#)).

Special Senses

Cases of hearing impairment or permanent loss have been reported predominantly in patients with chronic overdose.

Dermatological

Skin rash, pruritus.

The following adverse drug events may be borne in mind as potential effects of acetaminophen: allergic reactions, rash, thrombocytopenia, agranulocytosis.

Potential effects of high dosage are listed in the [OVERDOSAGE](#) section.

Drug Abuse and Dependence

Controlled Substance

Hydrocone bitartrate and acetaminophen tablets is classified as a Schedule III controlled substance.

Abuse and Dependence

Psychic dependence, physical dependence, and tolerance may develop upon repeated administration of narcotics; therefore, this product should be prescribed and administered with caution. However, psychic dependence is unlikely to develop when hydrocodone bitartrate and acetaminophen tablets are used for a short time for the treatment of pain.

Physical dependence, the condition in which continued administration of the drug is required to prevent the appearance of a withdrawal syndrome, assumes clinically significant proportions only after several weeks of continued narcotic use,

although some mild degree of physical dependence may develop after a few days of narcotic therapy. Tolerance, in which increasingly large doses are required in order to produce the same degree of analgesia, is manifested initially by a shortened duration of analgesic effect, and subsequently by decreases in the intensity of analgesia. The rate of development of tolerance varies among patients.

Overdosage

Following an acute overdosage, toxicity may result from hydrocodone or acetaminophen.

Signs and Symptoms

Hydrocodone: Serious overdose with hydrocodone is characterized by respiratory depression (a decrease in respiratory rate and/or tidal volume, Cheyne-Stokes respiration, cyanosis), extreme somnolence progressing to stupor or coma, skeletal muscle flaccidity, cold and clammy skin, and sometimes bradycardia and hypotension. In severe overdosage, apnea, circulatory collapse, cardiac arrest and death may occur.

Acetaminophen: In acetaminophen overdosage: dose-dependent, potentially fatal hepatic necrosis is the most serious adverse effect. Renal tubular necrosis, hypoglycemic coma, and coagulation defects may also occur.

Early symptoms following a potentially hepatotoxic overdose may include: nausea, vomiting, diaphoresis and general malaise. Clinical and laboratory evidence of hepatic toxicity may not be apparent until 48 to 72 hours post-ingestion.

Treatment

A single or multiple drug overdose with hydrocodone and acetaminophen is a potentially lethal polydrug overdose, and consultation with a regional poison control center is recommended.

Immediate treatment includes support of cardiorespiratory function and measures to reduce drug absorption.

Oxygen, intravenous fluids, vasopressors, and other supportive measures should be employed as indicated. Assisted or controlled ventilation should also be considered.

For hydrocodone overdose, primary attention should be given to the reestablishment of adequate respiratory exchange through provision of a patent airway and the institution of assisted or controlled ventilation. The narcotic antagonist naloxone hydrochloride is a specific antidote against respiratory depression which may result from overdosage or unusual sensitivity to narcotics, including hydrocodone. Since the duration of action of hydrocodone may exceed that of the antagonist, the patient should be kept under continued surveillance, and repeated doses of the antagonist should be administered as needed to maintain adequate respiration. A narcotic antagonist should not be administered in the absence of clinically significant respiratory or cardiovascular depression.

Gastric decontamination with activated charcoal should be administered just prior to N-acetylcysteine (NAC) to decrease systemic absorption if acetaminophen ingestion is known or suspected to have occurred within a few hours of presentation. Serum acetaminophen levels should be obtained immediately if the patient presents 4 hours or more after ingestion to assess potential risk of hepatotoxicity; acetaminophen levels drawn less than 4 hours post-ingestion may be misleading. To obtain the best possible outcome, NAC should be administered as soon as possible where impending or evolving liver injury is suspected. Intravenous NAC may be administered when circumstances preclude oral administration.

Vigorous supportive therapy is required in severe intoxication. Procedures to limit the continuing absorption of the drug must be readily performed since the hepatic injury is dose dependent and occurs early in the course of intoxication.

Vicodin ES Dosage and Administration

Dosage should be adjusted according to the severity of the pain and the response of the patient. However, it should be kept in mind that tolerance to hydrocodone can develop with continued use and that the incidence of untoward effects is dose related.

VICODIN® (Hydrocodone Bitartrate and Acetaminophen Tablets, USP 5 mg/300 mg): The usual adult dosage is one or two tablets every four to six hours as needed for pain. The total daily dosage should not exceed 8 tablets.

VICODIN ES® (Hydrocodone Bitartrate and Acetaminophen Tablets, USP 7.5 mg/300 mg): The usual adult dosage is one tablet every four to six hours as needed for pain. The total daily dosage should not exceed 6 tablets.

VICODIN HP® (Hydrocodone Bitartrate and Acetaminophen Tablets, USP 10 mg/300 mg): The usual adult dosage is one tablet every four to six hours as needed for pain. The total daily dosage should not exceed 6 tablets.

How is Vicodin ES supplied

VICODIN®, VICODIN ES® and VICODIN HP® (Hydrocodone Bitartrate and Acetaminophen) Tablets, USP are supplied as follows:

VICODIN® 5 mg/300 mg

White, capsule-shaped, bisected tablets, debossed “5” score “300” on one side and “VICODIN” on the other side in bottles of 100 and 500 tablets:

Bottles of 100 - NDC 0074-3041-13

Bottles of 500 - NDC 0074-3041-53

VICODIN ES® 7.5 mg/300 mg

White, capsule-shaped, bisected tablets, debossed “7.5” score “300” on one side and “VICODIN ES” on the other side in bottles of 100 and 500 tablets:

Bottles of 100 - NDC 0074-3043-13

Bottles of 500 - NDC 0074-3043-53

VICODIN HP® 10 mg/300 mg

White, capsule-shaped, bisected tablets, debossed “10” score “300” on one side and “VICODIN HP” on the other side in bottles of 100 and 500 tablets:

Bottles of 100 - NDC 0074-3054-13

Bottles of 500 - NDC 0074-3054-53

How is Vicodin ES supplied

Store at 20° to 25°C (68° to 77°F). [See USP Controlled Room Temperature].

PHARMACIST: Dispense in a tight, light-resistant container with a child-resistant closure.

A Schedule III Narcotic

© AbbVie Inc. 2014

Manufactured for

AbbVie Inc.

North Chicago, IL 60064 U.S.A.

Manufactured by:

Mikart, Inc.

Atlanta, GA 30318

1122F00 Rev. 06/14 June, 2014

NDC 0074–3054–53

Vicodin HP® (Hydrocodone Bitartrate and Acetaminophen) Tablets, USP

CIII

10 mg / 300 mg

500 Tablets

Rx only abbvie

This package is not intended for household use.

Protect from light.

Warning: Keep this and all medications out of the reach of children.

Manufactured for
AbbVie Inc.
North Chicago, IL 60064 U.S.A.

Manufactured by
MIKART, INC.
Atlanta, GA 30318

©AbbVie

Code 1122F50 Rev. 06/16

NDC 0074-3054-53

Vicodin HP®
(Hydrocodone Bitartrate
and Acetaminophen)
Tablets, USP

10 mg / 300 mg

500 Tablets


Rx only **abbvie**

Pharmacist: Dispense in a tight, light-resistant container with a child-resistant closure.

Each tablet contains 10 mg hydrocodone bitartrate (Warning: May be habit-forming) and 300 mg acetaminophen.

Storage: Store at 20° to 25°C (68° to 77°F) [see USP Controlled Room Temperature].

Usual adult dosage: One tablet every four to six hours, as needed for pain. Total daily dosage should not exceed six tablets. See package insert for full prescribing information.

 3 00743 05453 8

NDC 0074–3043–13

Vicodin ES® (Hydrocodone Bitartrate and Acetaminophen) Tablets, USP

CIII

7.5 mg / 300 mg

100 Tablets

Rx only abbvie

This package is not intended for household use.
Protect from light.
Warning: Keep this and all medications out of the reach of children.
Manufactured for
AbbVie Inc.
North Chicago, IL 60064 U.S.A.
Manufactured by
MIKART, INC.
Atlanta, GA 30318
©AbbVie Code 1124D10 Rev. 06/16

NDC 0074-3043-13

Vicodin ES®
(Hydrocodone Bitartrate
and Acetaminophen)
Tablets, USP

7.5 mg / 300 mg

100 Tablets

Rx only

abbvie

Pharmacist: Dispense in a tight, light-resistant container with a child-resistant closure.
Each tablet contains 7.5 mg hydrocodone bitartrate (Warning: May be habit-forming) and 300 mg acetaminophen.
Storage: Store at 20° to 25°C (68° to 77°F) [see USP Controlled Room Temperature].
Usual adult dosage: One tablet every four to six hours, as needed for pain. Total daily dosage should not exceed six tablets. See package insert for full prescribing information.

3 00743 04313 6

NDC 0074-3041-13

Vicodin® (Hydrocodone Bitartrate and Acetaminophen) Tablets, USP

CIII

5 mg / 300 mg

100 Tablets

Rx only abbvie

This package is not intended for household use.
Protect from light.
Warning: Keep this and all medications out of the reach of children.
Manufactured for
AbbVie Inc.
North Chicago, IL 60064 U.S.A.
Manufactured by
MIKART, INC.
Atlanta, GA 30318
©AbbVie Code 1123C10 Rev. 06/16

NDC 0074-3041-13

Vicodin®
(Hydrocodone Bitartrate
and Acetaminophen)
Tablets, USP

5 mg / 300 mg

100 Tablets

Rx only

abbvie

Pharmacist: Dispense in a tight, light-resistant container with a child-resistant closure.
Each tablet contains 5 mg hydrocodone bitartrate (Warning: May be habit-forming) and 300 mg acetaminophen.
Storage: Store at 20° to 25°C (68° to 77°F) [See USP Controlled Room Temperature].
Usual adult dosage: One or two tablets every four to six hours, as needed for pain. See package insert for full prescribing information.

3 00743 04113 2

VICODIN HP
hydrocodone bitartrate and acetaminophen tablet

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0074-3054
Route of Administration	ORAL	DEA Schedule	CIII

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
HYDROCODONE BITARTRATE (UNII: NO70W886KK) (HYDROCODONE - UNII:6YKS4Y3WQ7)	HYDROCODONE BITARTRATE	10 mg
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	300 mg

Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
STARCH, CORN (UNII: O8232NY3SJ)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
POVIDONES (UNII: FZ989GH94E)	
CROSPVIDONE (UNII: 68401960MK)	

Product Characteristics

Color	WHITE	Score	2 pieces
Shape	OVAL	Size	14mm
Flavor		Imprint Code	VICODIN;HP;10;300

Contains

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0074-3054-13	100 in 1 BOTTLE		
2	NDC:0074-3054-53	500 in 1 BOTTLE		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA040556	09/10/2012	

VICODIN ES

hydrocodone bitartrate and acetaminophen tablet

Product Information					
Product Type		HUMAN PRESCRIPTION DRUG		Item Code (Source)	NDC:0074-3043
Route of Administration		ORAL		DEA Schedule	CIII
Active Ingredient/Active Moiety					
Ingredient Name				Basis of Strength	Strength
HYDROCODONE BITARTRATE (UNII: NO70W886KK) (HYDROCODONE - UNII:6YKS4Y3WQ7)				HYDROCODONE BITARTRATE	7.5 mg
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)				ACETAMINOPHEN	300 mg
Product Characteristics					
Color	WHITE	Score	2 pieces		
Shape	OVAL	Size	14mm		
Flavor		Imprint Code	VICODIN;ES;75;300		
Contains					
Packaging					
#	Item Code	Package Description	Marketing Start Date	Marketing End Date	
1	NDC:0074-3043-13	100 in 1 BOTTLE			
2	NDC:0074-3043-53	500 in 1 BOTTLE			
Marketing Information					
Marketing Category		Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
ANDA		ANDA040556	09/10/2012		

VICODIN
hydrocodone bitartrate and acetaminophen tablet

Product Information			
Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0074-3041
Route of Administration	ORAL	DEA Schedule	CIII
Active Ingredient/Active Moiety			
Ingredient Name		Basis of Strength	Strength
HYDROCODONE BITARTRATE (UNII: NO70W886KK) (HYDROCODONE - UNII:6YKS4Y3WQ7)		HYDROCODONE BITARTRATE	5 mg
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)		ACETAMINOPHEN	300 mg
Product Characteristics			
Color	WHITE	Score	2 pieces
Shape	OVAL	Size	14mm
Flavor		Imprint Code	VICODIN;5;300
Contains			
Packaging			
#	Item Code	Package Description	Marketing Start Date Marketing End Date
1	NDC:0074-3041-13	100 in 1 BOTTLE	
2	NDC:0074-3041-53	500 in 1 BOTTLE	
Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA040658	09/10/2012	

Labeler - AbbVie Inc. (078458370)

More about Vicodin ES (acetaminophen / hydrocodone)

- [Check interactions](#)
- [Compare alternatives](#)
- [Reviews \(13\)](#)
- [Drug images](#)
- [Latest FDA alerts \(11\)](#)
- [Side effects](#)
- [Dosage information](#)

- [During pregnancy](#)
- [Drug class: narcotic analgesic combinations](#)

Professional resources

- [Hydrocodone and Acetaminophen Oral Solution prescribing information](#)

Other brands

[Norco](#), [Vicodin](#), [Lortab](#), [Hycet](#), ... +7 more

Related treatment guides

- [Cough](#)
- [Back Pain](#)
- [Pain](#)
- [Rheumatoid Arthritis](#)

[Medical Disclaimer](#)

DRUG STATUS

Availability		
	Discontinued	
Pregnancy & Lactation		
	Risk data available	
CSA Schedule*		
2	High potential for abuse	▼
Approval History		
	Drug history at FDA	▼

User Reviews & Ratings

8.7 / 10

[13 Reviews](#)

Images

[Vicodin ES 300 mg / 7.5 mg \(VICODIN ES 7.5 300\)](#)



Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.



About

[About Drugs.com](#)

[Advertising policy](#)

[Attribution & citations](#)

Terms & privacy

[Terms of use](#)

[Editorial policy](#)

[Privacy policy](#)

Support

[Help center](#)


[Sitemap](#)

[Contact us](#)



[Subscribe to our newsletter](#) for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.



Ad Choices
ADVERTISEMENT



Copyright © 2000-2025 Drugs.com. All rights reserved.