



[Home](#)

2. [Alli](#)

# alli

**Generic name:** [orlistat](#) [ *OR-li-stat* ]

**Brand names:** alli, [Xenical](#)

**Drug class:** [Peripherally acting antiobesity agents](#)

Medically reviewed by [Melisa Puckey, BPharm](#). Last updated on Mar 1, 2024.

[Uses](#) [Warnings](#) [Before taking](#) [Dosage](#) [Side effects](#) [Interactions](#)

## What is alli?

alli blocks some of the fat that you eat, keeping it from being absorbed by your body.

alli is used to aid in [weight loss](#), or to help reduce the risk of regaining weight already lost. This medicine must be used together with a reduced-calorie diet. Orlistat is for use only in adults.

Xenical is the prescription-strength form of orlistat. The alli brand of orlistat is available without a prescription.

## Warnings

**Do not take alli if you are pregnant.** Weight loss is not recommended during pregnancy.

You should not use alli if you have a digestive disorder (problems absorbing food). Do not use **alli** if you have had an organ transplant, if you use [cyclosporine](#), or if you are not overweight.

alli is only part of a complete program of treatment that also includes diet, exercise, and weight control. Your daily intake of fat, protein, and carbohydrates should be evenly divided over all of your daily meals. Follow your diet, medication, and exercise routines very closely.

Avoid a diet that is high in fat. High-fat meals taken in combination with orlistat can increase your risk of unpleasant side effects on your stomach or intestines.

## Before taking this medicine

You should not use alli if you are allergic to orlistat, or if you have malabsorption syndrome (an inability to absorb food and nutrients properly).

You also should not use Xenical if you have:

- gallbladder problems; or

- if you are pregnant.

Do not use alli if:

- you are not overweight;
- you have had an organ transplant; or
- you use cyclosporine (to prevent organ transplant rejection).

To make sure alli is safe for you, tell your doctor if you have ever had:

- kidney stones;
- gallbladder disease;
- pancreatitis;
- underactive thyroid;
- liver disease;
- kidney disease; or
- an eating disorder (anorexia or bulimia).

**Do not use alli if you are pregnant.** Weight loss is not recommended during pregnancy, even if you are overweight. Stop taking orlistat and tell your doctor right away if you become pregnant.

Taking orlistat can make it harder for your body to absorb certain vitamins. These vitamins are important if you are nursing a baby. **Do not use this medicine without a doctor's advice if you are breast-feeding a baby.**

Do not give **alli** to anyone under 18 years old.

 [Alli pregnancy and breastfeeding warnings](#) (more detail)

## How should I take alli?

Use alli exactly as directed on the label, or as prescribed by your doctor. Do not use in larger or smaller amounts or for longer than recommended.

**Never share orlistat with another person, especially someone with a history of eating disorders.**

Read all patient information, medication guides, and instruction sheets provided to you. Ask your doctor or pharmacist if you have any questions.

alli is usually taken 3 times per day with each main meal that contains some fat (no more than 30% of the calories for that meal). You may take the medicine either with your meal or up to 1 hour after eating.

If you skip a meal or you eat a meal that does not contain any fat, skip your alli dose for that meal.

The fat content of your daily diet should not be greater than 30% of your total daily caloric intake. For example, if you eat 1200 calories per day, no more than 360 of those calories should be in the form of fat.

Read the label of all food items you consume, paying special attention to the number of servings per container. Your doctor, nutrition counselor, or dietitian can help you develop a healthy eating plan.

alli is only part of a complete program of treatment that also includes diet, exercise, and weight control. Your daily intake of fat, protein, and carbohydrates should be evenly divided over all of your daily meals. Follow your diet, medication, and exercise routines very closely.

Orlistat can make it harder for your body to absorb certain vitamins, and you may need to take a vitamin and mineral supplement while you are taking orlistat. Follow your doctor's instructions about the type of supplement to use. **Take the supplement at bedtime, or at least 2 hours before or after you take alli.**

Store at room temperature away from moisture, heat, and light. Keep the bottle tightly closed. Throw away any unused alli after the expiration date on the medicine label has passed.

Keep track of your medicine. Orlistat is a drug that may be misused as a weight-loss aid, and you should be aware if anyone is using your medicine improperly or without a prescription.

 [Detailed Alli dosage information](#)

## What happens if I miss a dose?

Take the missed dose as soon as you remember, but no more than 1 hour after eating a meal. If it has been more than an hour since your last meal, skip the missed dose and take the medicine at your next regularly scheduled time. Do not take extra medicine to make up the missed dose.

## What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222.

## What to avoid

Avoid eating high-fat meals or you could have unpleasant side effects on your stomach or intestines.

If you also take cyclosporine, do not take it within 3 hours before or 3 hours after you take orlistat.

If you also take levothyroxine (such as Synthroid), do not take it within 4 hours before or 4 hours after you take orlistat.

## alli side effects

Get emergency medical help if you have **signs of an allergic reaction to alli**: hives; difficult breathing; swelling of your face, lips, tongue, or throat.

Stop using alli and call your doctor at once if you have:

- severe stomach pain;
- severe pain in your lower back;
- blood in your urine, painful or difficult urination;
- **kidney problems** - little or no urinating; swelling in your feet or ankles; feeling tired or short of breath; or
- **liver problems** - nausea, upper stomach pain, itching, tired feeling, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes).

Common alli side effects caused by its fat-blocking action. These are signs that the medicine is working properly. These side effects are usually temporary and may lessen as you continue using alli:

- oily or fatty stools;
- oily spotting in your undergarments;
- orange or brown colored oil in your stool;
- gas and oily discharge;
- loose stools, or an urgent need to go to the bathroom, inability to control bowel movements;
- an increased number of bowel movements; or
- stomach pain, nausea, rectal pain.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

 [Alli side effects](#) (more detail)

## Related/similar drugs

### Wegovy

Wegovy is a once-weekly semaglutide injection used for chronic weight management and to reduce ...

#### Reviews & ratings

7.6 / 10

[478 Reviews](#)

[View more](#)

### Ozempic

Learn about Ozempic (semaglutide) for type 2 diabetes treatment, weight management, cardiovascular ...

#### Reviews & ratings

6.6 / 10

[1,504 Reviews](#)

[View more](#)

#### FEATURED

### Mounjaro

Mounjaro (tirzepatide) is FDA-approved to manage type 2 diabetes in adults. It lowers blood sugar ...

## Reviews & ratings

8.5 / 10

[1,278 Reviews](#)

[View more](#)

## Rybelsus

Rybelsus tablets are used to improve blood sugar control in adults with type 2 diabetes, and may ...

## Reviews & ratings

6.4 / 10

[282 Reviews](#)

[View more](#)

## Zepbound

Zepbound (tirzepatide) is an FDA-approved weekly injection for weight loss and obstructive sleep ...

## Reviews & ratings

8.9 / 10

[289 Reviews](#)

[View more](#)

## Victoza

Victoza helps control blood sugar levels and reduce the risk of serious heart problems in people ...

## Reviews & ratings

7.6 / 10

[766 Reviews](#)

[View more](#)

## Saxenda

Saxenda (liraglutide) injection is used for weight loss in obese or overweight patients. Includes ...

## Reviews & ratings

7.5 / 10

[1,448 Reviews](#)

[View more](#)

### Liraglutide

Liraglutide may be used to improve blood sugar control in adults and children 10 years of age and ...

#### Reviews & ratings

7.5 / 10

[2,328 Reviews](#)

[View more](#)

### Tirzepatide

Tirzepatide is a once-weekly injection used for weight loss, sleep apnea, and type 2 diabetes ...

#### Reviews & ratings

8.6 / 10

[1,819 Reviews](#)

[View more](#)

### Phentermine

Phentermine is an appetite suppressant used together with diet and exercise to treat obesity. Learn ...

#### Reviews & ratings

8.6 / 10

[3,014 Reviews](#)

[View more](#)

## What other drugs will affect alli?

Ask a doctor or pharmacist if it is safe for you to use alli if you are also using any of the following drugs:

- amiodarone;
- insulin or oral diabetes medicine;
- HIV or AIDS medications;
- seizure medicine (especially if your seizures get worse while taking orlistat);

- a vitamin or mineral supplement that contains beta-carotene or vitamin E; or
- **a blood thinner** - warfarin, Coumadin, Jantoven.

This list is not complete. Other drugs may interact with orlistat, including prescription and over-the-counter medicines, vitamins, and herbal products. Not all possible interactions are listed in this medication guide.

**i** [Alli drug interactions](#) (more detail)

## Does Alli interact with my other drugs?

Enter medications to view a detailed interaction report using our [Drug Interaction Checker](#).

Alli

+

Enter a drug name

Add

### References

1. [Alli Product Label](#)

## More about Alli (orlistat)

- [Check interactions](#)
- [Compare alternatives](#)
- [Pricing & coupons](#)
- [Reviews \(121\)](#)
- [Drug images](#)
- [Side effects](#)
- [Dosage information](#)
- [During pregnancy](#)
- [Support group](#)
- [FDA approval history](#)
- [Drug class: peripherally acting antiobesity agents](#)
- [Breastfeeding](#)
- [En español](#)

### Patient resources

#### Other brands

[Xenical](#)

## Professional resources

- [Alli prescribing information](#)
- [Orlistat \(AHFS Monograph\)](#)

## Other brands

[Xenical](#)

## Related treatment guides

- [Weight Loss \(Obesity/Overweight\)](#)

## Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use alli only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

[Medical Disclaimer](#)

Copyright 1996-2025 Cerner Multum, Inc. Version: 12.01.

## Guide to Weight Loss Drugs

Find out everything you need to know about weight loss drugs in our prescription [weight loss drug guide](#).

### DRUG STATUS

Availability	
OTC	Over the counter
Pregnancy & Lactation	
	Risk data available
CSA Schedule*	
N/A	Not a controlled drug
Approval History	
	Drug history at FDA

## User Reviews & Ratings

6.2 / 10

[121 Reviews](#)

## Images



# Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.



## About

[About Drugs.com](#)

[Advertising policy](#)

[Attribution & citations](#)

## Terms & privacy

[Terms of use](#)

[Editorial policy](#)

[Privacy policy](#)

## Support

[Help center](#)

[Sitemap](#)

[Contact us](#)



[Subscribe to our newsletter](#) for the latest medication news, new drug approvals and FDA alerts.

**Drugs.com** provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.



Ad Choices  
ADVERTISEMENT



Copyright © 2000-2025 Drugs.com. All rights reserved.