

[Home](#)2. [Aripiprazole](#)

# Aripiprazole

**Generic name:** aripiprazole (oral) [ *AR-i-PIP-ra-zole* ]**Brand name:** [Abilify](#)**Dosage forms:** oral solution (1 mg/mL), oral tablet (10 mg; 15 mg; 2 mg; 20 mg; 30 mg; 5 mg), oral tablet, disintegrating (10 mg; 15 mg)**Drug class:** [Atypical antipsychotics](#)Medically reviewed by [Philip Thornton, DipPharm](#). Last updated on Aug 26, 2024.[Uses](#) [Warnings](#) [Before taking](#) [Dosage](#) [Side effects](#) [Interactions](#) [FAQ](#)

## What is aripiprazole?

[Aripiprazole](#) is an antipsychotic medicine that is used to treat [schizophrenia](#) in adults and children at least 13 years old.

Aripiprazole is also used in children 6 years or older who have Tourette's disorder, or symptoms of autistic disorder (irritability, aggression, mood swings, temper tantrums, and self-injury).

Aripiprazole is used alone or with a mood stabilizer medicine to treat [bipolar I disorder](#) (manic [depression](#)) in adults and children at least 10 years old.

Aripiprazole is used with antidepressant medication to treat [major depressive disorder](#) in adults.

## Warnings

Aripiprazole is not approved for use in older adults with [dementia](#)-related psychosis.

People with depression or mental illness may have thoughts about suicide. Some young people may have increased suicidal thoughts when first starting a medicine to treat depression. Tell your doctor right away if you have any sudden changes in mood or behavior, or thoughts about suicide.

Do not stop using aripiprazole suddenly, or you could have unpleasant withdrawal symptoms.

Avoid driving and doing other tasks or actions that call for you to be alert or have clear eyesight until you see how aripiprazole affects you.

High blood sugar or diabetes, high cholesterol, and weight gain have happened with drugs like this one. These may raise the chance of heart and brain blood vessel disease.

Low white blood cell counts have happened with drugs like this one. This may lead to a higher chance of infection. Rarely, infections have been deadly. Tell your doctor if you have ever had a low white blood cell count. Call your doctor right away if you have signs of infection like fever, chills, or sore throat.

## Before taking this medicine

You should not take aripiprazole if you are allergic to it.

Aripiprazole may increase the risk of death in older adults with dementia-related psychosis and is not approved for this use.

To make sure aripiprazole is safe for you, tell your doctor if you have::

- heart problems or a stroke;
- high or [low blood pressure](#);
- diabetes (in you or a family member);
- [seizures](#); or
- low white blood cell (WBC) counts.

People with depression or mental illness may have thoughts about suicide. Some young people may have increased suicidal thoughts when first starting a medicine to treat depression. Stay alert to changes in your mood or symptoms. Your family or caregivers should also watch for sudden changes in your behavior.

The liquid form (oral solution) of this medication contains sugar. Before taking aripiprazole oral solution, tell your doctor if you have diabetes.

Aripiprazole can cause high blood sugar (hyperglycemia). If you are diabetic, check your blood sugar levels carefully.

The orally disintegrating tablet form of this medication may contain over 3 milligrams of phenylalanine per tablet. Before taking Abilify Discmelt, tell your doctor if you have [phenylketonuria](#) (PKU).

**Taking antipsychotic medicine in the last 3 months of pregnancy may cause serious problems in the newborn.**

Tell your doctor right away if you get pregnant. **Do not stop the medicine without your doctor's advice.**

If you are pregnant, your name may be listed on a pregnancy registry to track the effects of aripiprazole on the baby.

Ask a doctor if it is safe to breastfeed while using aripiprazole.

 [Aripiprazole pregnancy and breastfeeding warnings](#) (more detail)

## How should I take aripiprazole?

Take aripiprazole exactly as prescribed by your doctor. Follow all directions on your prescription label and read all medication guides or instruction sheets. Your doctor may occasionally change your dose.

Aripiprazole can be taken with or without food.

Swallow the **regular tablet** whole and do not crush, chew, or break it. Do not split the **orally disintegrating tablet**.

Measure liquid medicine carefully. Use the dosing syringe provided, or use a medicine dose-measuring device (not a kitchen spoon)..

Remove an orally disintegrating tablet from the package only when you are ready to take the medicine. Place the tablet in your mouth and allow it to dissolve, without chewing. Swallow several times as the tablet dissolves. If needed, you may drink liquid to help swallow the dissolved tablet.

Your doctor will need to check your progress on a regular basis.

If you also use antidepressant medicine, do not stop using it suddenly or you could have unpleasant symptoms. Ask your doctor before stopping the antidepressant.

Store at room temperature away from moisture and heat. Aripiprazole **liquid** may be used for up to 6 months after opening, but not after the [expiration date](#) on the medicine label.

 [Detailed Aripiprazole dosage information](#)

## What happens if I miss a dose?

Take the medicine as soon as you can, but skip the missed dose if it is almost time for your next dose. Do not take two doses at one time.

Get your prescription refilled before you run out of medicine completely.

## What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222.

Overdose symptoms may include drowsiness, [vomiting](#), aggression, confusion, tremors, fast or [slow heart rate](#), seizure (convulsions), weak or shallow breathing, fainting, or coma.

## What should I avoid while taking aripiprazole?

Avoid getting up too fast from a sitting or lying position, or you may feel dizzy. Avoid driving or hazardous activity until you know how aripiprazole will affect you. [Dizziness](#) or drowsiness can cause falls, accidents, or severe injuries.

**Avoid drinking alcohol.** Dangerous side effects could occur.

While taking aripiprazole, you may be more sensitive to very hot conditions. Avoid becoming overheated or dehydrated. Drink plenty of fluids, especially in hot weather and during exercise.

## Aripiprazole side effects

Get emergency medical help if you have **signs of an allergic reaction to aripiprazole:** [hives](#); difficult breathing; swelling of your face, lips, tongue, or throat.

Tell your doctor right away if you have new or sudden changes in mood or behavior, including new or worse depression or [anxiety](#), [panic attacks](#), trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, more active or talkative, or have thoughts about suicide or hurting yourself.

Call your doctor at once if you have:

- severe [agitation](#), distress, or restless feeling;
- twitching or uncontrollable movements of your eyes, lips, tongue, face, arms, or legs;
- mask-like appearance of the face, trouble swallowing, problems with speech;

- seizure (convulsions);
- **severe nervous system reaction** - very stiff (rigid) muscles, high fever, sweating, confusion, fast or uneven heartbeats, tremors, feeling like you might pass out;
- **low blood cell counts** - fever, chills, [sore throat](#), weakness, easy bruising, unusual bleeding, purple or red spots under your skin; or
- **high blood sugar** - increased thirst, increased urination, dry mouth, fruity breath odor.

You may have increased sexual urges, unusual urges to gamble, or other intense urges while taking this medicine. Talk with your doctor if this occurs.

Common aripiprazole side effects may include:

- blurred vision;
- increased saliva or [drooling](#);
- muscle stiffness;
- uncontrolled muscle movements, shaking, anxiety, feeling restless;
- weight gain;
- [nausea](#), vomiting, [constipation](#);
- increased or decreased appetite;
- [headache](#), dizziness, drowsiness, feeling tired;
- sleep problems ([insomnia](#)); or
- [cold symptoms](#) such as stuffy nose, sneezing, sore throat;

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

 [Aripiprazole side effects](#) (more detail)

## Related/similar drugs

### Lybalvi

Lybalvi is used to treat adults with manic or mixed episodes associated with bipolar I disorder ...

#### Reviews & ratings

8.1 / 10

[16 Reviews](#)

[View more](#)

### Fanapt

Fanapt (iloperidone) is an antipsychotic medication used to treat schizophrenia. Includes Fanapt ...

#### Reviews & ratings

6.4 / 10

[30 Reviews](#)

[View more](#)

#### FEATURED

### Caplyta

Caplyta (lumateperone) may be used to treat schizophrenia or depression associated with bipolar ...

#### Reviews & ratings

6.6 / 10

[184 Reviews](#)

[View more](#)

### Quetiapine

Quetiapine is an atypical antipsychotic used to treat schizophrenia, bipolar disorder, and major ...

#### Reviews & ratings

7.2 / 10

[2,249 Reviews](#)

[View more](#)

### Fluoxetine

Fluoxetine is used to treat depression or obsessive-compulsive disorder in adults. Learn about side ...

#### Reviews & ratings

7.3 / 10

[2,232 Reviews](#)

[View more](#)

### Duloxetine

Duloxetine is a selective serotonin and norepinephrine reuptake inhibitor antidepressant used to ...

### Reviews & ratings

6.2 / 10

[2,456 Reviews](#)

[View more](#)

### Bupropion

Bupropion is an antidepressant used for depression, seasonal affective disorder, smoking cessation ...

### Reviews & ratings

7.0 / 10

[2,898 Reviews](#)

[View more](#)

### Sertraline

Sertraline (Zoloft) is an SSRI used to treat depression, anxiety, OCD, PTSD, and PMDD by increasing ...

### Reviews & ratings

7.3 / 10

[4,561 Reviews](#)

[View more](#)

### Trazodone

Trazodone is a tetracyclic antidepressant used to treat depression and anxiety disorders. Includes ...

### Reviews & ratings

6.3 / 10

[1,432 Reviews](#)

[View more](#)

## What other drugs will affect aripiprazole?

Taking aripiprazole with other drugs that make you drowsy or slow your breathing can cause dangerous side effects or death.

Sometimes it is not safe to use certain medicines at the same time. Some drugs can affect your blood levels of other drugs you use, which may increase side effects or make the medicines less effective.

Many drugs can interact with aripiprazole. This includes prescription and over-the-counter medicines, [vitamins](#), and [herbal products](#). Not all possible interactions are listed here. Tell your doctor about all other medicines you use.

 [Aripiprazole drug interactions](#) (more detail)





## Does aripiprazole interact with my other drugs?

Enter medications to view a detailed interaction report using our [Drug Interaction Checker](#).

+

Add

## Popular FAQ

- Why should you take aripiprazole in the morning? 
- What is the difference between Abilify and Abilify Maintena? 
- Does Abilify cause weight gain? 
- How does Abilify MyCite work? 

## More FAQ

- [What drugs cause tardive dyskinesia?](#)

[View more FAQ...](#)

## More about aripiprazole

- [Check interactions](#)
- [Compare alternatives](#)
- [Pricing & coupons](#)
- [Reviews \(1,540\)](#)
- [Drug images](#)
- [Latest FDA alerts \(5\)](#)
- [Side effects](#)

- [Dosage information](#)
- [Patient tips](#)
- [During pregnancy](#)
- [Support group](#)
- [Drug class: atypical antipsychotics](#)
- [Breastfeeding](#)

## Patient resources

- [Aripiprazole \(Intramuscular\) advanced reading](#)
- [Aripiprazole lauroxil \(Advanced Reading\)](#)

## Other brands

[Abilify](#), [Abilify Maintena](#), [Aristada](#), [Abilify Asimtufii](#), ... +3 more

## Professional resources

- [Aripiprazole monograph](#)
- [Aripiprazole ODT \(FDA\)](#)
- [Aripiprazole Oral Solution \(FDA\)](#)
- [Aripiprazole Tablets \(FDA\)](#)

## Other brands

[Abilify](#), [Abilify Maintena](#), [Aristada](#), [Abilify Asimtufii](#), ... +3 more

## Related treatment guides

- [Borderline Personality Disorder](#)
- [Bipolar Disorder](#)
- [Autism](#)
- [Depression](#)

## Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use aripiprazole medication only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

### [Medical Disclaimer](#)



## DRUG STATUS

### Availability

**Rx** Prescription only

### Pregnancy & Lactation


 Risk data available

### CSA Schedule\*

**N/A** Not a controlled drug



### Approval History

 Drug history at FDA



## User Reviews & Ratings

6.1 / 10

[1,540 Reviews](#)

## Images

[Aripiprazole 5 mg \(ARI 5\)](#)



[View larger images](#)

## Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.



### About

[About Drugs.com](#)

[Advertising policy](#)

[Attribution & citations](#)

### Terms & privacy

[Terms of use](#)

[Editorial policy](#)

[Privacy policy](#)

### Support

[Help center](#)

[Sitemap](#)

[Contact us](#)





[Subscribe to our newsletter](#) for the latest medication news, new drug approvals and FDA alerts.

**Drugs.com** provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.



Ad Choices  
ADVERTISEMENT



Copyright © 2000-2025 Drugs.com. All rights reserved.