

Home

Trezix

Trezix

Generic name: acetaminophen, caffeine, and dihydrocodeine [a-SEET-a-MIN-oh-fen, KAF-een, dye-HYE-droe-KOE-deen]

Brand names: Trezix, Dvorah, Panlor Tablet Drug class: Narcotic analgesic combinations



Medically reviewed by Drugs.com on Aug 4, 2023. Written by Cerner Multum.

Uses Side effects Warnings Before taking Dosage Interactions

What is Trezix?

Trezix is a combination medicine used to relieve moderate to severe pain.

Trezix may also be used for purposes not listed in this medication guide.

Trezix side effects

Get emergency medical help if you have **signs of an allergic reaction**: hives; difficult breathing; swelling of your face, lips, tongue, or throat.

In rare cases, acetaminophen may cause a severe skin reaction that can be fatal. This could occur even if you have taken acetaminophen in the past and had no reaction. Stop taking Trezix and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling.

Opioid medicine can slow or stop your breathing, and death may occur. A person caring for you should seek emergency medical attention if you have slow breathing with long pauses, blue colored lips, or if you are hard to wake up.

Trezix may cause serious side effects. Call your doctor at once if you have:

- noisy breathing, sighing, shallow breathing, breathing that stops during sleep;
- · chest pain, fast or pounding heartbeats, feeling light-headed, fainting;
- confusion, extreme drowsiness, unusual thoughts or behavior;
- painful or difficult urination;
- severe constipation;
- liver problems--nausea, upper stomach pain, itching, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes); or
- low cortisol levels-- nausea, vomiting, loss of appetite, dizziness, worsening tiredness or weakness.

Seek medical attention right away if you have symptoms of serotonin syndrome, such as: agitation, hallucinations, fever, sweating, shivering, fast heart rate, muscle stiffness, twitching, loss of coordination, nausea, vomiting, or diarrhea.

Serious side effects may be more likely in older adults and those who are overweight, malnourished, or debilitated.

Long-term use of opioid medication may affect fertility (ability to have children) in men or women. It is not known whether opioid effects on fertility are permanent.

Common side effects include:

- dizziness, drowsiness, tiredness, headache;
- nausea, vomiting, stomach pain, constipation;
- · sweating, itching; or
- · skin rash.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

i Trezix side effects (more detail)

Related/similar drugs

Oxycodone

Oxycodone is an opioid analgesic used to treat moderate to severe pain; it has a high potential for ...

Reviews & ratings

6.9 / 10

1,142 Reviews

View more

Tylenol

Tylenol is a pain reliever and a fever reducer used to treat many conditions such as headaches ...

Reviews & ratings

5.8 / 10

30 Reviews

View more

FEATURED

Repatha

Repatna (evolocumab) is a PUSK9 inhibitor used to lower high cholesterol alongside dietary changes ...

Reviews & ratings

5.3 / 10

685 Reviews

View more

Naproxen

Naproxen is a nonsteroidal antiinflammatory drug used to treat pain or inflammation caused by ...

Reviews & ratings

6.8 / 10

683 Reviews

View more

Paracetamol

Paracetamol (Panadol, Calpol, Alvedon) is a widely used over-thecounter painkiller and fever ...

Reviews & ratings

6.1 / 10

13 Reviews

View more

Hydroxyzine

Hydroxyzine is an antihistamine used to treat itching, hives, and anxiety. It also acts as a ...

Reviews & ratings

5.9 / 10

1,404 Reviews

View more

Tramadol

Tramadol is an opioid medication that may be used to treat moderate to moderately severe chronic ...

Reviews & ratings

7.0 / 10

1,876 Reviews

View more

Cyclobenzaprine

Cyclobenzaprine is a muscle relaxant and works by blocking pain sensations. Includes ...

Reviews & ratings

6.0 / 10

618 Reviews

View more

Meloxicam

Meloxicam is a nonsteroidal antiinflammatory drug used to treat pain or inflammation caused by ...

Reviews & ratings

6.3 / 10

600 Reviews

View more

Aspirin

Aspirin is used to treat mild to moderate pain and to reduce fever or inflammation. Learn about ...

Reviews & ratings

7.8 / 10

51 Reviews

View more

Warnings

MISUSE OF OPIOID MEDICINE CAN CAUSE ADDICTION, OVERDOSE, OR DEATH. Keep the medication in a place where others cannot get to it.

Do not give this medicine to anyone younger than 12 years old, or anyone under 18 who recently had surgery to remove

the tonsils or adenoids.

Fatal side effects can occur if you use opioid medicine with alcohol, or with other drugs that cause drowsiness or slow your breathing.

Stop taking this medicine and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling.

Before taking this medicine

You should not use Trezix if you are allergic to acetaminophen, caffeine, or dihydrocodeine, or if you have:

- · severe asthma or breathing problems; or
- a blockage in your stomach or intestines.

This medicine is **not approved** for use by anyone younger than 12 years old.

Do not give this medicine to anyone younger than 18 years old who recently had surgery to remove the tonsils or adenoids.

Tell your doctor if you have ever had:

- breathing problems, sleep apnea;
- liver or kidney disease;
- · alcoholism or drug addiction;
- mental illness or psychosis;
- a seizure or head injury;
- · urination problems; or
- problems with your pancreas, thyroid, or adrenal gland.

If you use opioid medicine while you are pregnant, your baby could become dependent on the drug. This can cause life-threatening withdrawal symptoms in the baby after it is born. Babies born dependent on opioids may need medical treatment for several weeks.

Do not breastfeed. Dihydrocodeine can pass into breast milk and cause drowsiness, breathing problems, or death in a nursing baby.

How should I take Trezix?

Follow the directions on your prescription label and read all medication guides. **Never use this medicine in larger amounts, or for longer than prescribed. An overdose can damage your liver or cause death.** Tell your doctor if you feel an increased urge to take more of this medicine.

Never share opioid medicine with another person, especially someone with a history of drug abuse or addiction. MISUSE CAN CAUSE ADDICTION, OVERDOSE, OR DEATH. Keep the medication in a place where others cannot get to it. Selling or giving away opioid medicine is against the law.

If you need surgery or medical tests, tell the doctor ahead of time that you are using this medicine.

Do not stop using this medicine suddenly after long-term use, or you could have unpleasant withdrawal symptoms. Ask your doctor how to safely stop using the medicine.

Store at room temperature away from moisture and heat. Keep track of your medicine. You should be aware if anyone is using it improperly or without a prescription.

Do not keep leftover opioid medication. **Just one dose can cause death in someone using this medicine accidentally or improperly.** Ask your pharmacist where to locate a drug take-back disposal program. If there is no take-back program, flush the unused medicine down the toilet.

Detailed Trezix dosage information

What happens if I miss a dose?

Since Trezix is used for pain, you are not likely to miss a dose. Skip any missed dose if it is almost time for your next dose. **Do not** use two doses at one time.

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222. **An overdose of this medicine can be fatal, especially in a child or other person using the medicine without a prescription.** Overdose can cause severe muscle weakness, pinpoint pupils, very slow breathing, extreme drowsiness, or coma.

What should I avoid while taking Trezix?

Avoid driving or operating machinery until you know how Trezix will affect you. Dizziness or drowsiness can cause falls, accidents, or severe injuries.

Do not drink alcohol. Dangerous side effects or death could occur.

Ask a doctor or pharmacist before using any other medicine that may contain acetaminophen (sometimes abbreviated as APAP). **Taking too much acetaminophen can lead to a fatal overdose.**

What other drugs will affect Trezix?

You may have breathing problems or withdrawal symptoms if you start or stop taking certain other medicines. Tell your doctor if you also use an antibiotic, antifungal medication, heart or blood pressure medication, seizure medication, or medicine to treat HIV or hepatitis C.

Opioid medication can interact with many other drugs and cause dangerous side effects or death. Be sure your doctor knows if you also use:

- other narcotic medications--opioid pain medicine or prescription cough medicine;
- a sedative like Valium--diazepam, alprazolam, lorazepam, Xanax, Klonopin, Versed, and others;
- drugs that make you sleepy or slow your breathing--a sleeping pill, muscle relaxer, medicine to treat mood disorders or mental illness; or

drugs that affect serotonin levels in your body--a stimulant, or medicine for depression, Parkinson's disease,
migraine headaches, serious infections, or nausea and vomiting.

This list is not complete. Other drugs may affect Trezix, including prescription and over-the-counter medicines, vitamins, and herbal products. **Not all possible interactions are listed here.**

Trezix drug interactions (more detail)

Does Trezix interact with my other drugs?

Enter medications to view a detailed interaction report using our Drug Interaction Checker.



More about Trezix (acetaminophen / caffeine / dihydrocodeine)

- Check interactions
- Compare alternatives
- Reviews (14)
- Drug images
- · Side effects
- Dosage information
- During pregnancy
- Drug class: narcotic analgesic combinations
- En español

Patient resources

· Trezix advanced reading

Other brands

Panlor, Dvorah, Panlor DC, Panlor SS, Zerlor

Professional resources

• Trezix prescribing information

Other brands

Dvorah

Related treatment guides

• Pain

Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use this medication only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

Medical Disclaimer

Copyright 1996-2025 Cerner Multum, Inc. Version: 10.01.

DRUG STATUS

Availability

Rx Prescription only

Pregnancy & Lactation

ন্% Risk data available

CSA Schedule*

3 Moderate abuse potential

Approval History

Drug history at FDA

User Reviews & Ratings

5.9 / 10

14 Reviews

Images

Trezix acetaminophen 356.4 mg / caffeine 30 mg / dihydrocodeine bitartrate 16 mg (TREZIX)

Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.





About

About Drugs.com

Advertising policy

Attribution & citations

Terms & privacy

Terms of use

Editorial policy

Privacy policy

Support

Help center

Sitemap

Contact us











Subscribe to our newsletter for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.







Copyright © 2000-2025 Drugs.com. All rights reserved.

