



[Home](#)

2. [Myrbetriq](#)

Myrbetriq

Generic name: [mirabegron](#) [*MIR-a-BEG-ron*]

Drug class: [Urinary antispasmodics](#)

Medically reviewed by [Kaci Durbin, MD](#). Last updated on May 1, 2024.

[Uses](#) [Warnings](#) [Before taking](#) [Dosage](#) [Side effects](#) [Interactions](#)

What is Myrbetriq?

Myrbetriq relaxes the muscles of the urinary bladder, reducing bladder spasms. It is used alone or with solifenacin succinate to treat the symptoms of overactive bladder, such as frequent or urgent urination and urinary incontinence.

Myrbetriq is used in adults to treat [overactive bladder](#) with symptoms of frequent or urgent urination and [urinary incontinence](#). It is sometimes used together with another medicine called solifenacin ([Vesicare](#)).

Myrbetriq is used in children to treat neurogenic detrusor overactivity (NDO). Urinary incontinence caused by NDO is related to permanent nerve damage from conditions such as [multiple sclerosis](#) or spinal injury.

Myrbetriq tablets may be used in children at least 3 years old. Myrbetriq granules are for use in children at least 3 years old who also weigh at least 77 pounds (35 kilograms).

Warnings

You should not use Myrbetriq if you have

- end stage renal disease (kidney failure)
- severe kidney disease; or
- severe uncontrolled high blood pressure.

A condition called angioedema can occur after taking Myrbetriq. Angioedema is a condition that leads to face, lip, tongue, throat swelling, difficulty breathing and can be fatal. Seek emergency attention immediately if these symptoms occur.

It may take up to 8 weeks before your symptoms improve. Keep using this medicine as directed and tell your doctor if your symptoms do not improve after 8 weeks of treatment.

Before taking this medicine

You should not use Myrbetriq if you are allergic to mirabegron.

To make sure this medicine is safe for you, tell your doctor if you have:

- high blood pressure;
- a bladder obstruction;
- trouble emptying your bladder (very little urine or a weak stream of urine);
- kidney disease; or
- liver disease.

Tell your doctor if you are pregnant or breastfeeding.

Myrbetriq should not be given to a child younger than 3 years old.

 [Myrbetriq pregnancy and breastfeeding warnings](#) (more detail)

How should I take Myrbetriq?

Take Myrbetriq exactly as prescribed by your doctor. Follow all directions on your prescription label and read all medication guides or instruction sheets.

If you take Myrbetriq with solifenacin, take both medicines at the same time each day.

Take the Myrbetriq **tablet** with a full glass of water. Swallow the tablet whole and do not crush, chew, or break it.

An adult may take a Myrbetriq tablet with or without food. However, a child should take the tablet with food.

Both adults and children should take Myrbetriq **granules** with food.

A pharmacist will mix the granules into a suspension (liquid) before you receive the medicine.

Shake the **suspension** before you measure a dose. Use the dosing syringe provided, or use a medicine dose-measuring device (not a kitchen spoon).

If the suspension will not be used for 2 days or longer, shake the bottle vigorously for 1 minute each day to keep the granules thoroughly mixed.

Read and carefully follow any Instructions for Use provided with your medicine. Ask your doctor or pharmacist if you do not understand these instructions.

Your dose needs may change if you switch from using tablets to using the suspension. Avoid medication errors by using only the form and strength your doctor prescribes.

Your blood pressure will need to be checked often.

It may take up to 8 weeks before your symptoms improve. Keep using the medication as directed and tell your doctor if your symptoms do not improve.

Store at room temperature away from moisture and heat. Throw away any unused Myrbetriq **suspension** after the [expiration date](#) on the medicine label has passed.

What happens if I miss a dose?

Use the medicine as soon as you can, but skip the missed dose if you are more than 12 hours late for the dose. Do not use two doses at one time.

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222.

An overdose may cause pounding heartbeats or fluttering in your chest.

What should I avoid while taking Myrbetriq?

Follow your doctor's instructions about any restrictions on food, beverages, or activity.

Myrbetriq side effects

Stop using Myrbetriq and get emergency medical help if you have **signs of an allergic reaction**: [hives](#); difficult breathing; swelling of your face, lips, tongue, or throat.

Call your doctor at once if you have:

- pain or burning when you urinate; or
- **dangerously high blood pressure** - severe [headache](#), blurred vision, pounding in your neck or ears.

Common Myrbetriq side effects may include:

- painful urination;
- increased blood pressure;
- sinus pain, [sore throat](#);
- [constipation](#);
- headache; or
- dry mouth or fast heartbeats (when taken with solifenacin).

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

Related/similar drugs

[Botox](#)

Botox is used to treat chronic

migraines, excessive sweating,
bladder conditions, eye muscle ...

Reviews & ratings

5.7 / 10

[473 Reviews](#)

[View more](#)

Detrol

Detrol is used to treat overactive
bladder symptoms of urinary
frequency, urgency, and ...

Reviews & ratings

7.5 / 10

[17 Reviews](#)

[View more](#)

FEATURED

Ozempic

Learn about Ozempic (semaglutide)
for type 2 diabetes treatment, weight
management, cardiovascular ...

Reviews & ratings

6.6 / 10

[1,504 Reviews](#)

[View more](#)

VESIcare

Vesicare is used to treat symptoms of
overactive bladder such as
incontinence and frequent ...

Reviews & ratings

5.6 / 10

[142 Reviews](#)

[View more](#)

Ditropan

Ditropan XL (oxybutynin extended
release) is used to treat the symptoms
of overactive bladder ...

Reviews & ratings

Reviews & ratings

4.7 / 10

[34 Reviews](#)

[View more](#)

Toviaz

Toviaz (fesoterodine) is used to treat overactive bladder with symptoms of urinary frequency ...

Reviews & ratings

5.2 / 10

[74 Reviews](#)

[View more](#)

Botox Cosmetic

Botox Cosmetic temporarily improves the look of moderate to severe frown lines, crow's feet ...

Reviews & ratings

4.9 / 10

[12 Reviews](#)

[View more](#)

Oxytrol

Oxytrol patches reduces muscle spasms in the urinary tract and are used to treat symptoms of ...

Reviews & ratings

7.4 / 10

[79 Reviews](#)

[View more](#)

Gelnique

Gelnique (oxybutynin) is used to treat symptoms of overactive bladder, such as frequent or urgent ...

Reviews & ratings

[7 Reviews](#)

[View more](#)

What other drugs will affect Myrbetriq?

Tell your doctor about all your other medicines, especially:

- [digoxin](#);
- [flecainide](#);
- [propafenone](#);
- solifenacin; or
- [thioridazine](#).

This list is not complete. Other drugs may interact with mirabegron, including prescription and over-the-counter medicines, [vitamins](#), and [herbal products](#). Not all possible drug interactions are listed here.

 [Myrbetriq drug interactions](#) (more detail)

Does Myrbetriq interact with my other drugs?

Enter medications to view a detailed interaction report using our [Drug Interaction Checker](#).

Myrbetriq

+

Enter a drug name

Add

References

1. [Myrbetriq Product Label](#)

More about Myrbetriq (mirabegron)

- [Check interactions](#)
- [Compare alternatives](#)
- [Pricing & coupons](#)
- [Reviews \(276\)](#)
- [Drug images](#)
- [Side effects](#)
- [Dosage information](#)
- [Patient tips](#)
- [During pregnancy](#)

- [Generic availability](#)
- [Support group](#)
- [FDA approval history](#)
- [Drug class: urinary antispasmodics](#)
- [Breastfeeding](#)
- [En español](#)

Professional resources

- [Myrbetriq prescribing information](#)
- [Mirabegron \(AHFS Monograph\)](#)

Related treatment guides

- [Urinary Frequency](#)
- [Neurogenic Detrusor Overactivity](#)
- [Urinary Incontinence](#)

Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use Myrbetriq only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

[Medical Disclaimer](#)

Copyright 1996-2025 Cerner Multum, Inc. Version: 4.01.

DRUG STATUS

| | | |
|---|-----------------------|---|
| Availability | | |
| Rx | Prescription only | |
| Pregnancy & Lactation | | |
|  | Risk data available | |
| CSA Schedule* | | |
| N/A | Not a controlled drug | ▼ |
| Approval History | | |
| 10+ years | FDA approved 2012 | ▼ |

User Reviews & Ratings

4.8 / 10

Images

Myrbetriq 50 mg (Logo 355)



[View larger images](#)

Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.



About

[About Drugs.com](#)
[Advertising policy](#)
[Attribution & citations](#)

Terms & privacy

[Terms of use](#)
[Editorial policy](#)
[Privacy policy](#)

Support

[Help center](#)
[Sitemap](#)
[Contact us](#)



[Subscribe to our newsletter](#) for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.



Ad Choices
ADVERTISEMENT

Copyright © 2000-2025 Drugs.com. All rights reserved.

