



[Home](#)

2. [Sporanox](#)

Sporanox

Generic name: [itraconazole](#) [*IT-ra-KON-a-zole*]

Drug class: [Azole antifungals](#)

Medically reviewed by [Philip Thornton, DipPharm](#). Last updated on Aug 15, 2023.

[Uses](#) [Warnings](#) [Before taking](#) [Dosage](#) [Side effects](#) [Interactions](#) [FAQ](#)

What is Sporanox?

Sporanox is an antifungal medication that fights infections caused by fungus.

Sporanox is used to treat infections in adults caused by fungus, which can invade any part of the body including the lungs, mouth or throat, [toenails](#), or [fingernails](#).

It is not known if Sporanox is safe and effective in children.

Warnings

You should not take Sporanox if you have ever had [heart failure](#).

If you have liver or kidney disease, you should not take Sporanox with colchicine, fesoterodine, or solifenacin.

Itraconazole may harm an unborn baby. Avoid getting pregnant while taking itraconazole and for 2 months after your last dose.

Stop using Sporanox and call your doctor at once if you have **signs of congestive heart failure**: feeling tired or short of breath, cough with mucus, fast heartbeats, swelling, rapid weight gain, or sleep problems.

Life-threatening side effects may occur if you take itraconazole with certain other drugs. Tell your doctor about all your current medicines and any you start or stop using.

Many drugs can interact with itraconazole, and some drugs should not be used together. This includes prescription and over-the-counter medicines, vitamins, and herbal products. Tell your doctor about all medicines you use, and those you start or stop using during your treatment with Sporanox.

Before taking this medicine

You should not take Sporanox if you are allergic to itraconazole or similar medicines such as [fluconazole](#) or [ketoconazole](#), or if you have ever had congestive heart failure.

Life-threatening side effects may occur if you take Sporanox with certain other drugs. Your doctor may change your treatment plan if you have used other medicines in the past 2 weeks, especially:

- [avanafil](#);
- [cisapride](#);
- [eliglustat](#);
- [irinotecan](#);
- [isavuconazonium](#);
- [methadone](#);
- [naloxegol](#);
- [ranolazine](#);
- [ticagrelor](#);
- [lurasidone](#) or [pimozide](#) (anti-psychotic medications);
- [lomitapide](#), [lovastatin](#), [simvastatin](#) (cholesterol-lowering medicines);
- [dihydroergotamine](#), [ergonovine](#), [ergotamine](#), or [methylergonovine](#) (ergot medicines);
- [eplerenone](#), [felodipine](#), [ivabradine](#), or [nisoldipine](#) (heart or blood pressure medicines);
- [disopyramide](#), [dofetilide](#), [dronedarone](#), or [quinidine](#) (medicines for heart rhythm disorders); or
- oral [midazolam](#), or [triazolam](#) ([Valium](#)-like sedatives).

If you have liver or kidney disease, you should not take Sporanox with colchicine, fesoterodine, solifenacin, or telithromycin.

Itraconazole may harm an unborn baby. Use effective birth control to prevent pregnancy while you are using this medicine and for at least 2 months after your last dose. Tell your doctor if you become pregnant.

To make sure Sporanox is safe for you, tell your doctor if you have ever had:

- heart problems;
- [cystic fibrosis](#) or other lung problems;
- liver or kidney disease; or
- a weak immune system.

It may not be safe to breastfeed while using itraconazole. Ask your doctor about any risk.

Itraconazole can pass into breast milk and may affect the nursing baby. Tell your doctor if you are breast-feeding.

 [Sporanox pregnancy and breastfeeding warnings](#) (more detail)

How should I take Sporanox?

Take Sporanox exactly as prescribed by your doctor. Follow all directions on your prescription label and read all medication guides or instruction sheets.

Do not share this medicine with another person, even if they have the same symptoms you have.

The Sporanox **capsule** should be taken with food.

Swallow the **capsule** whole and do not crush, chew, break, or open it.

Take the **oral solution (liquid)** on an empty stomach, at least 1 hour before or 2 hours after a meal. Swish the liquid in your mouth for several seconds before swallowing it.

Measure liquid medicine carefully. Use the dosing syringe provided, or use a medicine dose-measuring device (not a kitchen spoon).

The [Sporanox PulsePak](#) has a special dosing schedule that includes not taking the medicine for several days in a row. Follow all dosing instructions carefully.

Sporanox capsules should not be used in place of itraconazole oral solution (liquid) if that is what your doctor has prescribed. **Avoid medication errors by using only the form and strength your doctor prescribes.**

If you also take a stomach acid reducer ([Tagamet](#), [Pepcid](#), [Axid](#), [Zantac](#), and others), take Sporanox with an acidic drink such as non-diet cola.

Use this medicine for the full prescribed length of time, even if your symptoms quickly improve. Skipping doses can increase your risk of infection that is resistant to medication. Itraconazole will not treat a [viral infection](#) such as the flu or a [common cold](#).

You may need frequent medical tests.

Store at room temperature away from moisture, heat, and light.

 [Detailed Sporanox dosage information](#)

What happens if I miss a dose?

Take the medicine as soon as you can, but skip the missed dose if it is almost time for your next dose. Do not take two doses at one time.

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222.

What to avoid

Avoid driving or hazardous activity until you know how this medicine will affect you. Your reactions could be impaired.

Avoid taking [antacids](#) within 1 hour before or 2 hours after you take Sporanox. Some [antacids](#) can make it harder for your body to absorb itraconazole.

Sporanox side effects

Get emergency medical help if you have **signs of an allergic reaction to Sporanox**: [hives](#), severe [skin rash](#), tingling in

your arms or legs; difficult breathing; swelling of your face, lips, tongue, or throat.

Stop using this medicine and call your doctor at once if you have **signs of congestive heart failure**: feeling tired or short of breath, cough with mucus, fast heartbeats, swelling, rapid weight gain, or sleep problems.

Keep taking Sporanox but call your doctor at once if you have:

- confusion, a light-headed feeling (like you might pass out);
- blurred vision, double vision, ringing in your ears, problems with hearing;
- fast heartbeats;
- numbness or tingly feeling, loss of bladder control;
- little or no urinating, pain or burning when you urinate;
- **low potassium level** - leg cramps, [constipation](#), irregular heartbeats, fluttering in your chest, increased thirst or urination, muscle weakness or limp feeling;
- **pancreatitis** - severe pain in your upper stomach spreading to your back, [nausea and vomiting](#); or
- **liver problems** - [nausea](#), upper stomach pain, tiredness, loss of appetite, dark urine, clay-colored stools, [jaundice](#) (yellowing of the skin or eyes).

Common Sporanox side effects may include:

- [headache](#), [dizziness](#), drowsiness, tiredness;
- increased blood pressure;
- rash, itching;
- nausea, [vomiting](#), stomach pain, [diarrhea](#), constipation;
- swelling;
- abnormal liver function or blood tests;
- fever, muscle or joint pain;
- unusual or unpleasant taste in your mouth;
- [hair loss](#);
- [impotence](#), [erection problems](#); or
- changes in your menstrual periods.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

 [Sporanox side effects](#) (more detail)

Related/similar drugs

[Augmentin](#)

Augmentin is a prescription antibiotic combining amoxicillin and clavulanate

combining amoxicillin and clavulanate
to treat bacterial ...

Reviews & ratings

5.4 / 10

[447 Reviews](#)

[View more](#)

Diflucan

Diflucan (fluconazole) is used to treat
and prevent fungal infections. Includes
Diflucan side ...

Reviews & ratings

5.9 / 10

[144 Reviews](#)

[View more](#)

FEATURED

Ozempic

Learn about Ozempic (semaglutide)
for type 2 diabetes treatment, weight
management, cardiovascular ...

Reviews & ratings

6.6 / 10

[1,504 Reviews](#)

[View more](#)

Zosyn

Zosyn is used to treat bacterial
infections such as urinary tract and
skin infections and ...

Reviews & ratings

[5 Reviews](#)

[View more](#)

Acidophilus

Acidophilus helps maintain an acidic
environment, which can prevent the
growth of harmful bacteria ...

Reviews & ratings

9.5 / 10

[View more](#)

What other drugs will affect Sporanox?

Sometimes it is not safe to use certain medications at the same time. Some drugs can affect your blood levels of other drugs you take, which may increase side effects or make the medications less effective.

Many drugs can interact with itraconazole, and some drugs should not be used at the same time. Tell your doctor about all your current medicines and any medicine you start or stop using. This includes prescription and over-the-counter medicines, [vitamins](#), and [herbal products](#). Not all possible interactions are listed here.

 [Sporanox drug interactions](#) (more detail)

Does Sporanox interact with my other drugs?

Enter medications to view a detailed interaction report using our [Drug Interaction Checker](#).

Sporanox

+

Enter a drug name

Add

Popular FAQ

- How do I get rid of nail fungus?

▼
- What home remedies work well for toenail fungus?

▼

More FAQ

- [What is the difference between Tolsura and Sporanox?](#)

More about Sporanox (itraconazole)

- [Check interactions](#)
- [Compare alternatives](#)
- [Pricing & coupons](#)
- [Reviews \(18\)](#)
- [Drug images](#)
- [Side effects](#)

- [Dosage information](#)
- [During pregnancy](#)
- [Generic availability](#)
- [Drug class: azole antifungals](#)
- [Breastfeeding](#)
- [En español](#)

Patient resources

Other brands

[Onmel](#)

Professional resources

- [Sporanox prescribing information](#)
- [Itraconazole \(AHFS Monograph\)](#)

Other brands

[Tolsura](#), [Onmel](#)

Related treatment guides

- [Aspergillosis, Invasive](#)
- [Candida Urinary Tract Infection](#)
- [Aspergillosis, Aspergilloma](#)
- [Blastomycosis](#)

Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use Sporanox only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

[Medical Disclaimer](#)

Copyright 1996-2025 Cerner Multum, Inc. Version: 16.01.

DRUG STATUS

Availability

Rx Prescription only

Pregnancy & Lactation



Risk data available

CSA Schedule*

N/A Not a controlled drug



Approval History

10+
years

FDA approved 1992



User Reviews & Ratings

8.4 / 10

[18 Reviews](#)

Images

[Sporanox 100 mg \(JANSSEN SPORANOX 100\)](#)

Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.



About

[About Drugs.com](#)

[Advertising policy](#)

[Attribution & citations](#)

Terms & privacy

[Terms of use](#)

[Editorial policy](#)

[Privacy policy](#)

Support

[Help center](#)

[Sitemap](#)

[Contact us](#)



[Subscribe to our newsletter](#) for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and

others.



Ad Choices
ADVERTISEMENT



Copyright © 2000-2025 Drugs.com. All rights reserved.