

[Home](#)

2. Atropine/hyoscyamine/phenobarbital/scopolam..

Belladonna alkaloids and phenobarbital

Generic name: belladonna alkaloids and phenobarbital [*BEL-a-DON-a-AL-ka-loids-and-FEEN-oh-BAR-bi-tal*]

Brand names: [Donnatal](#), Phenobarbital with Belladonna Alkaloids, [Phenohydro](#), [D-Tal](#), Chardonna-2, ... [show all 23 brands](#)

Dosage forms: oral liquid (0.0194 mg-0.1037 mg-16.2 mg-0.0065 mg/5 mL), oral tablet (0.0194 mg-0.1037 mg-16.2 mg-0.0065 mg)

Drug class: [Anticholinergics / antispasmodics](#)



[Medically reviewed](#) by Drugs.com on Aug 4, 2023. Written by [Cerner Multum](#).

[Uses](#) [Side effects](#) [Warnings](#) [Before taking](#) [Dosage](#) [Interactions](#)

What is belladonna alkaloids and phenobarbital?

belladonna alkaloids and [phenobarbital](#) is made up of belladonna alkaloids (atropine, hyoscyamine, scopolamine) and phenobarbital.

Belladonna alkaloids and phenobarbital is a combination medicine used to treat [irritable bowel syndrome](#) and ulcers in the intestine.

Belladonna alkaloids and phenobarbital may also be used for purposes not listed in this medication guide.

Belladonna alkaloids and phenobarbital side effects

Get emergency medical help if you have **signs of an allergic reaction** (hives, difficult breathing, swelling in your face or throat) or a **severe skin reaction** (fever, sore throat, burning eyes, skin pain, red or purple skin rash with blistering and peeling).

This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have:

- fast or pounding heartbeats;
- blurred vision, tunnel vision, eye pain, or seeing halos around lights;
- diarrhea (especially if you have a colostomy or ileostomy);
- decreased sweating and hot or dry skin;
- painful or difficult urination;
- loss of coordination, slurred speech, fainting;
- confusion, agitation, depression, suicidal thoughts;
- fever, sore throat; or
- pale skin, easy bruising, unusual tiredness, cold hands and feet;

Side effects such as drowsiness, agitation, nervousness, and excitement may be likely to occur in older adults.

Common side effects of belladonna alkaloids and phenobarbital may include:

- constipation;
- decreased sweating or urination;
- dizziness, drowsiness, weakness;
- blurred vision;
- feeling nervous or excited;
- dry mouth, nose, or throat;
- dry skin; or
- nausea, vomiting, bloating.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

 [Atropine / hyoscyamine / phenobarbital / scopolamine side effects](#) (more detail)

Related/similar drugs

Pepcid

Pepcid is used to treat ulcers in the stomach and for conditions where the stomach produces too ...

Reviews & ratings

6.0 / 10

[10 Reviews](#)

[View more](#)

Prilosec

Prilosec (omeprazole) is used to treat symptoms of GERD and other conditions caused by excess ...

Reviews & ratings

6.9 / 10

[53 Reviews](#)

[View more](#)

FEATURED

OmvoH

OmvoH is used to treat moderate to

severe ulcerative colitis or Crohn's disease in adults. This ...

Reviews & ratings

[1 Reviews](#)

[View more](#)

Ranitidine

Ranitidine is used to treat and prevent ulcers in the stomach and intestines. Includes ranitidine ...

Reviews & ratings

7.2 / 10

[163 Reviews](#)

[View more](#)

Vancomycin

Vancomycin is used to treat infections of the intestines that cause colitis. Learn about side ...

Reviews & ratings

6.5 / 10

[57 Reviews](#)

[View more](#)

Lansoprazole

Lansoprazole is used to treat stomach and intestinal ulcers, erosive esophagitis and ...

Reviews & ratings

6.6 / 10

[143 Reviews](#)

[View more](#)

Protonix

Protonix (pantoprazole) is used to treat erosive esophagitis and other conditions involving excess ...

Reviews & ratings

7.2 / 10

[View more](#)

Warnings

belladonna alkaloids and phenobarbital may be habit-forming. **Misuse can cause addiction, overdose, or death.**

Before taking this medicine

You should not use this medicine if you are allergic to atropine, hyoscyamine, phenobarbital, or scopolamine. You also should not use this medicine if you have:

- a bladder obstruction, [enlarged prostate](#), or other urination problems;
- a stomach or [bowel obstruction](#) (including [paralytic ileus](#));
- chronic constipation or lack of bowel function (especially in older adults and those who are ill or debilitated);
- severe [ulcerative colitis](#) or toxic megacolon;
- glaucoma;
- reflux disease with a [hiatal hernia](#);
- active bleeding with fast heartbeats, [low blood pressure](#), [shortness of breath](#), and cold hands or feet;
- porphyria (a genetic enzyme disorder that causes symptoms affecting the skin or nervous system);
- an allergy to primidone; or
- a muscle disorder called [myasthenia gravis](#).

Tell your doctor if you have ever had:

- ulcerative colitis or a [stomach ulcer](#);
- a colostomy or ileostomy;
- a nerve-muscle disorder;
- heart problems, [high blood pressure](#);
- [gallbladder disease](#);
- alcoholism or [drug addiction](#);
- breathing problems;
- a thyroid disorder; or
- liver or kidney disease.

Belladonna alkaloids and phenobarbital may harm an unborn baby, but the benefit of treating your condition with this medicine may outweigh any risk. **Tell your doctor if you are pregnant or if you become pregnant.**

This medicine can make [birth control pills](#) less effective. Ask your doctor about using a non-hormonal [birth control](#)

(condom, diaphragm with spermicide) to prevent pregnancy.

It may not be safe to breast-feed while using this medicine. Ask your doctor about any risk. **This medicine can slow breast milk production.**

Do not give this medication to a child without the advice of a doctor.

How should I take belladonna alkaloids and phenobarbital?

Follow all directions on your prescription label and read all medication guides or instruction sheets. Use the medicine exactly as directed.

belladonna alkaloids and phenobarbital contains phenobarbital, a barbiturate that may be habit-forming. **Misuse can cause addiction, overdose, or death.** Keep the medication in a place where others cannot get to it. Selling or giving away this medicine is against the law.

Swallow the **extended-release tablet** whole and do not crush, chew, or break it.

Measure **liquid medicine** carefully. Use the dosing syringe provided, or use a medicine dose-measuring device (not a kitchen spoon).

Call your doctor if your symptoms do not improve, or if they get worse.

This medicine can affect the results of certain medical tests. Tell any doctor who treats you that you are using belladonna alkaloids and phenobarbital.

Store at room temperature away from moisture, heat, and light. Keep track of your medicine. You should be aware if anyone is using it improperly or without a prescription.

Do not stop using this medicine suddenly after long-term use, or you could have unpleasant withdrawal symptoms. Ask your doctor how to safely stop using this medicine.

 [Detailed Atropine / hyoscyamine / phenobarbital / scopolamine dosage information](#)

What happens if I miss a dose?

Take the medicine as soon as you can, but skip the missed dose if it is almost time for your next dose. **Do not** take two doses at one time.

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222.

Overdose symptoms may include headache, dizziness, nausea, vomiting, dry mouth, blurred vision, hot or dry skin, trouble swallowing, and feeling excited or agitated.

What should I avoid while taking belladonna alkaloids and phenobarbital?

belladonna alkaloids and phenobarbital may cause drowsiness or blurred vision and may impair your reactions. Avoid driving or hazardous activity until you know how this medicine will affect you.

Drinking alcohol with this medicine can increase certain side effects.

Avoid becoming overheated or dehydrated during exercise and in hot weather. This medicine can decrease sweating and you may be more prone to heat stroke.

What other drugs will affect belladonna alkaloids and phenobarbital?

Using this medicine with other drugs that slow your breathing can cause dangerous side effects. Ask your doctor before using opioid medication, a sleeping pill, a muscle relaxer, or medicine for anxiety or seizures.

If you take an antacid, take it at least 3 hours before or 3 hours after you take belladonna alkaloids and phenobarbital.

When you start or stop taking belladonna alkaloids and phenobarbital, your doctor may need to adjust the doses of any other medicines you take on a regular basis. Tell your doctor about all your other medicines, especially:

- potassium supplements;
- primidone;
- [St. John's wort](#);
- antibiotic or antifungal medicine;
- heart or blood pressure medication;
- antiviral medicine to treat hepatitis or HIV/AIDS; or
- a blood thinner (such as warfarin, Coumadin, Jantoven).

This list is not complete. Other drugs may affect belladonna alkaloids and phenobarbital, including prescription and over-the-counter medicines, vitamins, and [herbal products](#). Not all possible [drug interactions](#) are listed here.

 [Atropine / hyoscyamine / phenobarbital / scopolamine drug interactions](#) (more detail)

Does atropine/hyoscyamine/phenobarbital/scopolamine interact with my other drugs?

Enter medications to view a detailed interaction report using our [Drug Interaction Checker](#).

atropine/hyoscyamine/phenobarbital/scopolamine

+

Enter a drug name

Add

More about atropine / hyoscyamine / phenobarbital / scopolamine

- [Check interactions](#)
- [Compare alternatives](#)
- [Pricing & coupons](#)
- [Reviews \(62\)](#)

- [Drug images](#)
- [Side effects](#)
- [Dosage information](#)
- [During pregnancy](#)
- [Drug class: anticholinergics/antispasmodics](#)

Patient resources

- [Belladonna alkaloids and phenobarbital advanced reading](#)
- [Hyoscyamine, Atropine, Scopolamine, and Phenobarbital Elixir](#)
- [Hyoscyamine, Atropine, Scopolamine, and Phenobarbital Tablets](#)

Other brands

[Donnatal](#), [Antispasmodic](#), [Phenohydro](#), [D-Tal](#), ... +2 more

Professional resources

- [Belladonna Alkaloids with Phenobarbital prescribing information](#)
- [Phenobarbital with Belladonna Alkaloids Elixir \(FDA\)](#)
- [Phenobarbital, Hyoscyamine Sulfate, Atropine Sulfate and Scopolamine Hydrobromide Elixir \(FDA\)](#)
- [Phenobarbital, Hyoscyamine Sulfate, Atropine Sulfate and Scopolamine Hydrobromide Tablets \(FDA\)](#)

Other brands

[Donnatal](#), [Phenohydro](#)

Related treatment guides

- [Duodenal Ulcer](#)
- [Irritable Bowel Syndrome](#)
- [Enterocolitis](#)

Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use this medication only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

Medical Disclaimer

Copyright 1996-2025 Cerner Multum, Inc. Version: 7.01.

DRUG STATUS

Availability

Rx Prescription only

Pregnancy & Lactation

 Risk data available

CSA Schedule*

N/A Not a controlled drug



User Reviews & Ratings

9.2 / 10

[62 Reviews](#)

Images

Belladonna alkaloids with phenobarbital atropine 0.0194 mg / hyoscyamine 0.1037 mg / phenobarbital 16.2 mg / scopolamine 0.0065 mg (West-ward 140)



[View larger images](#)

Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.



About

[About Drugs.com](#)

[Advertising policy](#)

[Attribution & citations](#)

Terms & privacy

[Terms of use](#)

[Editorial policy](#)

[Privacy policy](#)

Support

[Help center](#)

[Sitemap](#)

[Contact us](#)





[Subscribe to our newsletter](#) for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.



Ad Choices
ADVERTISEMENT



Copyright © 2000-2025 Drugs.com. All rights reserved.