

Home

Glipizide

Glipizide 🕁

Generic name: glipizide [GLIP-i-zide]

Brand name: Glucotrol

Drug class: Sulfonylureas

Medically reviewed by Sanjai Sinha, MD. Last updated on Aug 26, 2024.

Uses Warnings Before taking Dosage Side effects Interactions

What is glipizide?

Glipizide is an oral diabetes medicine that helps control blood sugar levels by helping your pancreas produce insulin.

Glipizide is used together with diet and exercise to improve blood sugar control in adults with type 2 diabetes mellitus.

Glipizide is not for treating type 1 diabetes.

Warnings

You should not use glipizide if you have diabetic ketoacidosis (call your doctor for treatment).

Before taking this medicine

You should not use this medicine if you are allergic to glipizide, or if you have diabetic ketoacidosis (call your doctor for treatment).

Tell your doctor if you have ever had:

- liver or kidney disease;
- chronic diarrhea, or a blockage in your intestines; or
- an enzyme deficiency called glucose-6-phosphate dehydrogenase deficiency (G6PD).

Follow your doctor's instructions about using glipizide **if you are pregnant or you become pregnant.** Controlling diabetes is very important during pregnancy, and having high blood sugar may cause complications in both the mother and the baby. **You should not take glipizide during the last 2 weeks of pregnancy. Agents other than glipizide are currently recommended to treat diabetes in pregnant women.**

It may not be safe to breastfeed while using this medicine. Ask your doctor about any risk.

Glipizide pregnancy and breastfeeding warnings (more detail)

How should I take glipizide?

Take glipizide exactly as prescribed by your doctor. Follow all directions on your prescription label. Your doctor may occasionally change your dose. Do not take this medicine in larger or smaller amounts or for longer than recommended.

Take the regular tablet 30 minutes before your first meal of the day.

Take the glipizide extended-release tablet with your first meal of the day.

Swallow the tablet whole and do not crush, chew, or break it.

Your blood sugar may need to be checked often, and you may need other blood tests at your doctor's office.

Low blood sugar (hypoglycemia) can happen to anyone who has diabetes. Symptoms include headache, hunger, sweating, irritability, dizziness, nausea, fast heart rate, and feeling anxious or shaky. To quickly treat low blood sugar, always keep a fast-acting source of sugar with you such as fruit juice, hard candy, crackers, raisins, or non-diet soda.

Your doctor can prescribe a glucagon emergency injection kit to use in case you have severe hypoglycemia and cannot eat or drink. Be sure your family and close friends know how to give you this injection in an emergency.

Also watch for signs of high blood sugar (hyperglycemia) such as increased thirst or urination, blurred vision, headache, and tiredness.

Blood sugar levels can be affected by stress, illness, surgery, exercise, alcohol use, or skipping meals. **Ask your doctor** before changing your dose or medication schedule.

Some forms of glipizide are made with a shell that is not absorbed or melted in the body. Part of the tablet shell may appear in your stool. This is a normal side effect and will not make the medication less effective.

Store at room temperature away from moisture, heat, and light.

Detailed Glipizide dosage information

What happens if I miss a dose?

Take your dose as soon as you can, but only if you are getting ready to eat a meal. If you skip a meal, skip the missed dose and wait until your next meal. Do not take two doses at one time.

Get your prescription refilled before you run out of medicine completely.

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222. A glipizide overdose can cause life-threatening hypoglycemia.

Symptoms of severe hypoglycemia include extreme weakness, blurred vision, sweating, trouble speaking, tremors, stomach pain, confusion, and seizure (convulsions).

What to avoid

Avoid drinking alcohol. It lowers blood sugar and can cause side effects.

Avoid driving or operating machinery until you know how this medicine will affect you.

Glipizide side effects

Get emergency medical help if you have **signs of an allergic reaction to glipizide:** hives; difficulty breathing; swelling of your face, lips, tongue, or throat.

Call your doctor at once if you have symptoms of low blood sugar:

- · headache, irritability
- · sweating, fast heart rate;
- · dizziness, nausea; or
- hunger, feeling anxious or shaky.

Common glipizide side effects may include:

- · diarrhea, constipation, gas;
- · dizziness, drowsiness;
- tremors; or
- skin rash, redness, or itching.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

Glipizide side effects (more detail)

Related/similar drugs

Ozempic

Learn about Ozempic (semaglutide) for type 2 diabetes treatment, weight management, cardiovascular ...

Reviews & ratings

6.6 / 10

1,504 Reviews

View more

Rybelsus

Rybelsus tablets are used to improve blood sugar control in adults with type 2 diabetes, and may ...

Reviews & ratings

View more

FEATURED

Mounjaro

Mounjaro (tirzepatide) is FDAapproved to manage type 2 diabetes in adults. It lowers blood sugar ...

Reviews & ratings

8.5 / 10

1,278 Reviews

View more

Trulicity

Trulicity is an injectable diabetes medicine that is used together with diet and exercise to ...

Reviews & ratings

5.3 / 10

970 Reviews

View more

Lantus

Lantus is a long acting form of insulin used to treat type 1 or type 2 diabetes. Learn about side ...

Reviews & ratings

6.5 / 10

41 Reviews

View more

Tresiba

Tresiba (insulin degludec) is used to treat diabetes mellitus. Includes Tresiba side effects ...

Reviews & ratings

4.3 / 10

152 Reviews

Victoza

Victoza helps control blood sugar levels and reduce the risk of serious heart problems in people ...

Reviews & ratings

7.6 / 10

766 Reviews

View more

Basaglar

Basaglar (insulin glargine) is a longacting insulin that is used to improve blood sugar control in ...

Reviews & ratings

2.7 / 10

148 Reviews

View more

Levemir

Levemir (insulin detemir) is a long acting insulin used to treat diabetes in adults and children ...

Reviews & ratings

4.6 / 10

70 Reviews

View more

Novolog

NovoLog is a fast-acting insulin used to treat diabetes to control blood sugar levels in adults and ...

Reviews & ratings

5.5 / 10

32 Reviews

View more

What other drugs will affect glipizide?

Sometimes it is not safe to use certain medications at the same time. Some drugs can affect your blood levels of other drugs you take, which may increase side effects or make the medications less effective.

Many drugs can interact with glipizide. This includes prescription and over-the-counter medicines, vitamins, and herbal products. Not all possible interactions are listed here. Tell your doctor about all your current medicines and any medicine you start or stop using.

Glipizide drug interactions (more detail)

Does glipizide interact with my other drugs?

Enter medications to view a detailed interaction report using our Drug Interaction Checker.

glipizide
+
Enter a drug name
Add

More about glipizide

- · Check interactions
- Compare alternatives
- · Pricing & coupons
- Reviews (75)
- Drug images
- · Side effects
- Dosage information
- Patient tips
- During pregnancy
- Support group
- Drug class: sulfonylureas
- Breastfeeding
- En español

Patient resources

Other brands

Glucotrol, Glucotrol XL

Professional resources

- Glipizide monograph
- Glipizide (FDA)
- Glipizide ER (FDA)
- Glipizide Tablets (FDA)

Other brands

Glucotrol, GlipiZIDE XL, Glucotrol XL

Related treatment guides

• Type 2 Diabetes

Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use glipizide only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

Medical Disclaimer

Copyright 1996-2025 Cerner Multum, Inc. Version: 13.01.

Guide to Weight Loss Drugs

Find out everything you need to know about weight loss drugs in our prescription weight loss drug guide.

DRUG STATUS

Availability

Rx Prescription only

Pregnancy & Lactation

ঝ্ Risk data available

CSA Schedule*

N/A Not a controlled drug

Approval History

□ Drug history at FDA

User Reviews & Ratings

6.5 / 10

75 Reviews

Images

Glipizide 5 mg (APO GLP 5)



Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.





About

About Drugs.com
Advertising policy

Attribution & citations

Terms & privacy

Terms of use

Editorial policy

Privacy policy

Support

Help center

Sitemap

Contact us











Subscribe to our newsletter for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.







