

[Home](#)

2. Acanya

Acanya

Generic name: [benzoyl peroxide and clindamycin topical](#) [BEN-zoyl-per-OX-ide-and-clin-da-MYE-sin]**Drug class:** [Topical acne agents](#)[Medically reviewed](#) by Drugs.com. Last updated on Aug 7, 2023.[Uses](#) [Warnings](#) [Before taking](#) [Dosage](#) [Side effects](#) [Interactions](#)

What is Acanya?

Acanya Gel contains a combination of benzoyl peroxide and clindamycin. Benzoyl peroxide has an antibacterial effect. It also has a mild drying effect that allows excess oil and dirt to be washed away. Clindamycin is an antibiotic that prevents bacteria from growing on the skin.

Acanya Gel is used to treat acne vulgaris in patients 12 years or older.

Acanya may also be used for purposes not listed in this medication guide.

Warnings

Use Acanya exactly as your doctor has prescribed it for you. Using more medicine or applying it more often than prescribed will not make it work any faster, and may increase side effects. Do not use this medication for longer than your doctor has prescribed.

Avoid getting Acanya Gel in your eyes, mouth, or nose (or in the creases of your nose), or on your lips. If it does get into any of these areas, wash with water. Do not apply Acanya to sunburned, windburned, dry, chapped, irritated, or broken skin.

It may take several weeks before your symptoms improve. Keep using the medication as directed and tell your doctor if your symptoms do not improve.

Although Acanya is applied to the skin, your body may absorb enough clindamycin to cause serious side effects. You may not be able to use this medication if you have inflammation of your intestines (also called enteritis), ulcerative colitis, or if you have ever had severe diarrhea caused by antibiotic medicine.

Avoid exposure to sunlight or tanning beds. This medication can make you sunburn more easily. Wear protective clothing and use sunscreen (SPF 30 or higher) when you are outdoors.

Before using Acanya

Do not use Acanya if you are allergic to benzoyl peroxide or clindamycin (Cleocin, Clina-Derm, Clindets).

Although Acanya is applied to the skin, your body may absorb enough clindamycin to cause serious side effects. You may not be able to use this medication if you have:

- inflammation of your intestines (also called enteritis);
- ulcerative colitis; or
- if you have ever had severe diarrhea caused by antibiotic medicine.

It is not known whether Acanya will harm an unborn baby. Tell your doctor if you are pregnant or plan to become pregnant while using this medication. It is not known whether benzoyl peroxide and clindamycin passes into breast milk or if it could harm a nursing baby. You should not breast-feed while you are using Acanya.

i [Acanya pregnancy and breastfeeding warnings](#) (more detail)

How should I use Acanya?

Use Acanya exactly as prescribed by your doctor. Do not use in larger or smaller amounts or for longer than recommended. Follow the directions on your prescription label. Using more medicine or applying it more often than prescribed will not make it work any faster, and may increase side effects.

Wash your hands before and after applying Acanya Gel.

Wash your face with a mild cleanser (not soap), rinse with warm water and pat the skin dry with a clean towel.

Acanya is usually applied once daily. Apply Acanya Gel as a thin layer to the affected areas (excluding the eyes and lips).

Avoid getting this medication in your eyes, mouth, or nose (or in the creases of your nose), or on your lips. If it does get into any of these areas, wash with water. Do not apply Acanya to sunburned, windburned, dry, chapped, irritated, or broken skin.

It may take several weeks before your symptoms improve. Keep using Acanya Gel as directed and tell your doctor if your symptoms do not improve.

Store at room temperature away from moisture and heat. Do not freeze. Keep the jar tightly closed. Throw away any unused medicine 2 months after the date you filled the prescription.

i [Detailed Acanya dosage information](#)

What happens if I miss a dose?

Apply the missed dose as soon as you remember. Skip the missed dose if it is almost time for your next scheduled dose. Do not use extra medicine to make up the missed dose.

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222.

Overdose symptoms include bloody or watery diarrhea, which may result if you absorb benzoyl peroxide and clindamycin through your skin by applying too much.

What should I avoid?

Avoid using skin products that can cause irritation, such as harsh soaps, shampoos, or skin cleansers, hair coloring or permanent chemicals, hair removers or waxes, or skin products with alcohol, spices, astringents, or lime. Do not use other medicated skin products unless your doctor has told you to.

Antibiotic medicines can cause diarrhea, which may be a sign of a new infection. If you have diarrhea that is watery or has blood in it, call your doctor. Do not use any medicine to stop the diarrhea unless your doctor has told you to.

Avoid exposure to sunlight or tanning beds. Acanya Gel can make you sunburn more easily. Wear protective clothing and use sunscreen (SPF 30 or higher) when you are outdoors.

Avoid using sunscreen containing PABA on the same skin treated with Acanya, or skin discoloration may occur.

Benzoyl peroxide can bleach hair or fabrics. Do not let Acanya come into contact with clothing, hair, or colored towels or bed linens.

Acanya side effects

Get emergency medical help if you have any signs of an allergic reaction to Acanya: hives; difficulty breathing; swelling of your face, lips, tongue, or throat.

Stop using Acanya and call your doctor at once if you have any of these serious side effects:

- severe redness, burning, stinging, or peeling of treated skin areas; or
- diarrhea that is watery or bloody.

Less serious Acanya side effects may include:

- mild burning or stinging;
- itching or tingly feeling;
- dryness or peeling of treated skin; or
- redness or other irritation.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

 [Acanya side effects](#) (more detail)

Related/similar drugs

FEATURED

Ozempic

Learn about Ozempic (semaglutide) for type 2 diabetes treatment, weight management, cardiovascular ...

Reviews & ratings

6.6 / 10

[1,504 Reviews](#)

[View more](#)

Soolantra

Soolantra (ivermectin) is used to treat rosacea. Includes Soolantra side effects, interactions and ...

Reviews & ratings

7.7 / 10

[299 Reviews](#)

[View more](#)

Tazorac

Tazorac cream and gel is used to treat acne and plaque psoriasis. Includes Tazorac side effects ...

Reviews & ratings

7.9 / 10

[38 Reviews](#)

[View more](#)

What other drugs will affect Acanya?

Tell your doctor about all other medicines you use, especially:

- erythromycin topical (Akne-Mycin, Emcin Clear, Eryderm, Erygel, Erythra-Derm, Ery-Sol, and others); or
- erythromycin taken by mouth (E.E.S., EryPed, Ery-Tab, Erythrocin, Pediazole, and others).

This list is not complete and other drugs may interact with Acanya. Tell your doctor about all medications you use. This includes prescription, over-the-counter, vitamin, and herbal products. Do not start a new medication without telling your doctor.

 [Acanya drug interactions](#) (more detail)

Does Acanya interact with my other drugs?

Enter medications to view a detailed interaction report using our [Drug Interaction Checker](#).

Acanya

+

Enter a drug name

Add

More about Acanya (benzoyl peroxide / clindamycin topical)

- [Check interactions](#)
- [Compare alternatives](#)
- [Pricing & coupons](#)
- [Reviews \(56\)](#)
- [Side effects](#)
- [Dosage information](#)
- [During pregnancy](#)
- [Generic availability](#)
- [FDA approval history](#)
- [Drug class: topical acne agents](#)
- [En español](#)

Patient resources

Other brands

[Benzaclin](#)

Professional resources

- [Acanya prescribing information](#)

Other brands

[Benzaclin](#), [Duac](#), [Onexton](#), [Neuac](#)

Related treatment guides

- [Acne](#)

Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and

use Acanya Gel only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

Medical Disclaimer

Copyright 1996-2025 Cerner Multum, Inc. Version: 11.01.

Antibiotics 101

List of Common Antibiotics & Types

Antibiotics and Alcohol

Probiotics and Antibiotics

Antibiotics for Strep Throat

Antibiotics for a Tooth Infection

Antibiotics for UTI's

DRUG STATUS

Availability

Rx Prescription only

Pregnancy & Lactation

 Risk data available

CSA Schedule*

N/A Not a controlled drug



Approval History

 Drug history at FDA



User Reviews & Ratings

7.5 / 10

56 Reviews

Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.



About

About Drugs.com

[Advertising policy](#)
[Attribution & citations](#)

Terms & privacy

[Terms of use](#)
[Editorial policy](#)
[Privacy policy](#)

Support

[Help center](#)
[Sitemap](#)
[Contact us](#)



[Subscribe to our newsletter](#) for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.



Ad Choices
ADVERTISEMENT



Copyright © 2000-2025 Drugs.com. All rights reserved.