

Home

Uptravi

Uptravi

Generic name: selexipag (oral/injection) [se-LEX-i-pag]

Drug class: Agents for pulmonary hypertension

Medically reviewed by Judith Stewart, BPharm. Last updated on Oct 24, 2024.

Uses Warnings Before taking Dosage Side effects Interactions FAQ

What is Uptravi?

Uptravi lowers blood pressure in your lungs by dilating (widening) blood vessels that supply blood to the lungs.

Uptravi is used to treat pulmonary arterial hypertension (PAH).

Uptravi helps keep your condition from getting worse and helps lower your risk of needing to be hospitalized for PAH.

Warnings

Tell your doctor about all your current medicines and any you start or stop using. Many drugs can interact, and some drugs should not be used with Uptravi.

Before taking this medicine

You should not use Uptravi if you are allergic to selexipag.

Many drugs can interact and cause dangerous effects. Some drugs should not be used together with selexipag. Your doctor may change your treatment plan if you also use:

- · clopidogrel; or
- gemfibrozil.

To make sure Uptravi is safe for you, tell your doctor if you have ever had:

- a condition called pulmonary veno-occlusive disease (narrowing of the veins that carry blood from the lungs to the heart); or
- liver disease.

It is not known if selexipag will harm an unborn baby. Tell your doctor if you are pregnant or plan to become pregnant.

You should not breastfeed while using Uptravi.

How should I take Uptravi?

Take Uptravi exactly as prescribed by your doctor. Follow all directions on your prescription label and read all medication guides or instruction sheets. Your doctor may occasionally change your dose.

Uptravi tablets are taken by mouth. You may take the tablets with or without food.

Swallow the tablet whole and do not crush, chew, or break it.

Uptravi injection is given in a vein by a healthcare provider if you are unable to take medicine by mouth.

You should not stop taking Uptravi without your doctor's advice. Stopping suddenly may make your condition worse.

Store tablets at room temperature away from moisture and heat.

Dosing information

Usual Adult Dose for Pulmonary Hypertension:

Initial dose: 200 mcg orally twice a day

Maintenance dose: Increase in increments of 200 mcg orally twice a day at weekly intervals to the highest tolerated dose

Maximum dose: 1600 mcg orally twice a day

Comment:

-If the patient reaches a dose that is not well tolerated, reduce to a previously tolerated dose.

Use: For the treatment of pulmonary arterial hypertension (PAH) to delay disease progression and reduce the risk of hospitalization for PAH

Detailed Uptravi dosage information

What happens if I miss a dose?

Use the medicine as soon as you can, but skip the missed dose if your next dose is due in less than 6 hours. Do not use two doses at one time.

If you miss doses for 3 or more days, call your doctor before taking another dose. You may need to restart the medicine at a lower dose than you took before.

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222.

What should I avoid while taking Uptravi?

Follow your doctor's instructions about any restrictions on food, beverages, or activity.



Uptravi side effects

Get emergency medical help if you have **signs of an allergic reaction to Uptravi:** hives; difficult breathing; swelling of your face, lips, tongue, or throat.

Call your doctor at once if you have:

- low red blood cells (anemia) pale skin, tiredness, feeling light-headed or short of breath, cold hands and feet; or
- **new lung problems** anxiety, sweating, pale skin, severe shortness of breath, wheezing, gasping for breath, cough with foamy mucus, chest pain, fast or uneven heart rate.

Common Uptravi side effects may include:

- · headache;
- jaw pain;
- · muscle pain, pain in your arms or legs;
- · joint pain;
- flushing (warmth, redness, or tingly feeling);
- rash;
- anemia;
- · nausea, vomiting, loss of appetite, diarrhea; or
- pain, redness, or swelling where a Uptravi injection was given.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

Uptravi side effects (more detail)

Related/similar drugs

Cialis

Cialis (tadalafil) increase increases blood flow to a certain area of the body and is used to treat ...

Reviews & ratings

8.6 / 10

554 Reviews

View more

Viagra

Viagra (sildenafil) is used to treat erectile dysfunction or impotence in men. Includes Viagra side ...

Reviews & ratings

8.5 / 10

253 Reviews

View more

FEATURED

Ozempic

Learn about Ozempic (semaglutide) for type 2 diabetes treatment, weight management, cardiovascular ...

Reviews & ratings

6.6 / 10

1,504 Reviews

View more

Adcirca

Addirca is used to treat pulmonary arterial hypertension. Learn about side effects, interactions ...

Reviews & ratings

6 Reviews

View more

Opsumit

Opsumit (macitentan) is used to treat pulmonary arterial hypertension. Includes Opsumit side ...

Reviews & ratings

8.2 / 10

10 Reviews

View more

Revatio

Revatio (sildenafil) is used to treat pulmonary arterial hypertension. Includes Revatio side ...

Reviews & ratings

6.9 / 10

10 Reviews

View more

Adempas

Adempas (riociguat) is used to treat pulmonary hypertension. Includes Adempas side effects ...

Reviews & ratings

4 Reviews

View more

What other drugs will affect Uptravi?

Sometimes it is not safe to use certain medicines at the same time. Some drugs can affect your blood levels of other drugs you use, which may increase side effects or make the medicines less effective.

Other drugs may interact with selexipag, including prescription and over-the-counter medicines, vitamins, and herbal products. Tell your doctor about all other medicines you use.

Uptravi drug interactions (more detail)

Does Uptravi interact with my other drugs?

Enter medications to view a detailed interaction report using our Drug Interaction Checker.

Uptravi
+
Enter a drug name
Add

Popular FAQ

What is the drug Uptravi used for?

More about Uptravi (selexipag)

- · Check interactions
- Compare alternatives
- Pricing & coupons
- Reviews (27)

- Drug images
- · Side effects
- Dosage information
- During pregnancy
- · Generic availability
- FDA approval history
- · Drug class: agents for pulmonary hypertension
- Breastfeeding
- En español

Patient resources

- Uptravi (Selexipag Intravenous) advanced reading
- Uptravi (Selexipag Oral) (Advanced Reading)

Professional resources

- Uptravi prescribing information
- Selexipag (AHFS Monograph)

Related treatment guides

• Pulmonary Arterial Hypertension

Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use Uptravi only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

Medical Disclaimer

Copyright 1996-2025 Cerner Multum, Inc. Version: 4.01.

DRUG STATUS

Availability

Rx Prescription only

Pregnancy & Lactation

& Risk data available

CSA Schedule*

N/A Not a controlled drug

Approval History 9 years FDA approved 2015

User Reviews & Ratings

5.4 / 10

27 Reviews

Images

Uptravi 1000 mcg (10)





View larger images

Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.





About

About Drugs.com

Advertising policy

Attribution & citations

Terms & privacy

Terms of use

Editorial policy

Privacy policy

Support

Help center

Sitemap

Contact us













Subscribe to our newsletter for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.









